

7.5: Average Value of a Function

The average value of finitely many numbers y_1, y_2, \dots, y_n :

The **average value of a function** $y = f(x)$ over the interval $[a, b]$:

$$f_{ave} = \frac{1}{b-a} \int_a^b f(x) dx.$$


EXAMPLE 1. Determine the average value of $f(x) = x^2 - 4x + 7 \sin(\pi x)$ over the interval $[-\frac{1}{2}; \frac{1}{2}]$.

EXAMPLE 2. The temperature of a metal rod, 10 m long, is $5x$ (in $^{\circ}C$) at a distance x meters from one end of the rod. What is the average temperature of the rod?

EXAMPLE 3. Find the value(s) b s. t. the average value of $f(x) = 3 + 2x - 3x^2$ on the interval $[0, b]$ is equal to 1.

MEAN VALUE THEOREM FOR INTEGRALS: *If f is continuous on $[a, b]$, then there exists a number c on $[a, b]$ s.t.*

$$\int_a^b f(x) dx = f(c)(b - a).$$



The geometric interpretation of the Mean Value Theorem for Integrals: for *positive* functions f , there is a number c s.t. the rectangle with base $[a, b]$ and height $f(c)$ has the same area as the region under the graph of f from a to b .

EXAMPLE 4. *If g is continuous and $\int_{-1}^7 g(x) dx = 24$ show that g takes on the value 3 at least once on the interval $[-1, 7]$.*

EXAMPLE 5. *Determine the number c that satisfies the Mean Value Theorem for Integrals for the function $f(x) = x^2 - 2x - 2$ on the interval $[1, 4]$*