

## 10.6: Representation of Functions as Power Series

Problem: Find the sum of the following series:  $\sum_{n=0}^{\infty} x^n$

EXAMPLE 1. Find a power series representation for  $f(x)$  and determine the interval of convergence.

(a)  $f(x) = \frac{1}{5-x}$

(b)  $f(x) = \frac{x}{5-x}$

(c)  $f(x) = \frac{1}{1+x^2}$

### Differentiation and Integration of power series

THEOREM 2. If the power series  $\sum_{n=0}^{\infty} c_n(x-a)^n$  has radius of convergence  $R > 0$ , then the function  $f$  defined by

$$f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n$$

is differentiable (and therefore continuous) on the interval  $(a-R, a+R)$  and

- $f'(x) = \sum_{n=0}^{\infty} n c_n(x-a)^{n-1}$
- $\int f(x) dx = C + \sum_{n=0}^{\infty} \frac{c_n}{n+1} (x-a)^{n+1}$

The radii of convergence of the power series for  $f'(x)$  and  $\int f(x) dx$  are both  $R$ .

EXAMPLE 3. Find a power series representation for  $f(x)$  and determine the radius of convergence.

(a)  $f(x) = \frac{1}{(5-x)^2}$

**(b)**  $f(x) = \ln(2 + x)$