


# Trace Test

Special Session on Applied Algebraic Geometry  
12 November 2016



Frank Sottile

sottile@math.tamu.edu



Work with Anton Leykin and Jose Israel Rodriguez.

# Witness sets in Numerical Algebraic Geometry

Numerical algebraic geometry uses the ability to solve systems of polynomial equations to study algebraic varieties on a computer.

It represents a variety  $V \subset \mathbb{C}^n$  as a *witness set*  $W = V \cap L$ , where  $L$  is a general affine linear space of codimension  $m = \dim V$ .

Basic problem: Given a subset  $W' \subset W$  how to certify that  $W' = W$ ?

Trace Test: Suppose that  $L(t)$  for  $t \in \mathbb{C}$  is a pencil of affine-linear spaces with  $L(0) = L$ . Use continuation to follow points of  $W'$  along  $t$ , obtaining sets  $W'(t)$ . Then the trace of points in  $W'(t)$ ,

$$\text{Tr}(W'(t)) := \sum \{w \mid w \in W'(t)\},$$

is an affine function of  $t$  if and only if  $W' = W$ .

# Proof of Trace Test

A general irreducible curve in  $\mathbb{C}^2$  is defined by a dense irreducible polynomial  $f \in \mathbb{C}[x, t]$  of degree  $d$ . Normalize  $f$  so that 1 = coefficient of  $x^d$ .

$f \in \mathbb{C}(t)[x]$  is irreducible and monic. The negative sum of its roots is its coefficient of  $x^{d-1}$ . Thus

$$\text{trace}(K/\mathbb{C}(t))(x) = c_0 t + c_1 \quad c_0, c_1 \in \mathbb{C}, \quad (1)$$

where  $K$  contains the roots of  $f$ .

A general pencil  $L(t)$  spans a codimension  $m-1$  plane  $M$  with  $M \cap V$  a curve, and  $M$  has coordinates  $(\underline{x}, t)$ . By (1),  $\text{Tr}(W(t))$  is an affine function when  $W$  is a witness set.

This does not hold for  $\text{Tr}(W'(t))$  if  $W' \subsetneq W$ , as the monodromy in  $t$  is the full symmetric group. **explain**

# Multihomogeneous Witness Sets

A subvariety  $V \subset \mathbb{P}^A \times \mathbb{P}^B$  of dimension  $m$  has *multidegrees*  $d_{a,b}$  for  $a+b = m$ : For a general codimension  $a$  plane  $L \subset \mathbb{P}^A$  and a general codimension  $b$  plane  $M \subset \mathbb{P}^B$ ,

$$d_{a,b}(V) = \#V \cap (L \times M).$$

Definition (Hauenstein-Rodriguez) An intersection  $W_{a,b} = V \cap (L \times M)$  is a *multihomogeneous witness set* of bidimension  $(a, b)$  for  $V$ .

Advantages:

- (1) Reflects the structure of  $V$  in  $\mathbb{P}^A \times \mathbb{P}^B$ .
- (2) Smaller than alternatives. Embedding  $V$  into  $\mathbb{P}^{AB+A+B}$  via Segre  $\sigma$ ,

$$\deg(\sigma(V)) = \sum_{a+b=m} \binom{m}{a} d_{a,b}.$$

This is yuge.

# Using Multihomogeneous Witness Sets

Hauenstein and Rodriguez showed that many algorithms in numerical algebraic geometry work well with multihomogeneous witness sets.

These include regeneration, membership, and using a multihomogeneous witness set in one bidimension to populate another.

What does not work well is the trace test.

Fact. If  $L(t) \subset \mathbb{P}^A$  and  $M(s) \subset \mathbb{P}^B$  are pencils of affine spaces of codimensions  $a$  and  $b$ , respectively, then  $\text{Tr}(V \cap (L(t) \times M(s)))$  is **not** a bilinear function in  $s$  and  $t$ .

We cannot even fix  $t$  and let  $s$  vary for irreducible decomposition, for  $V \cap L$  could be reducible even if  $V$  is irreducible.

# Dimension Reduction

Let  $V \subset \mathbb{P}^A \times \mathbb{P}^B$  be irreducible of dimension  $m \geq 2$ ,  $a+b = m$  with  $d_{a,b}(V) \neq 0$ ,  $L' \subset \mathbb{P}^A$  a general linear space of codimension  $a-1$ , and  $M' \subset \mathbb{P}^B$  a general linear space of codimension  $b-1$ .

$U := V \cap (L' \times M')$  is irreducible of dimension 2 with multidegrees

$$d_{0,2} = d_{a-1,b+1}(V), \quad d_{1,1} = d_{a,b}(V), \quad d_{2,0} = d_{a+1,b-1}(V).$$

Either (1)  $d_{0,2} = d_{2,0} = 0 \Rightarrow U$  is a product of curves. Then  $V$  is also a product and we may treat each factor separately.

Or (2) a further linear slice is possible, reducing  $V$  to a curve in a product of projective spaces.

The two cases are detected from the tangent space of  $V$  or of  $U$ .

# A Multihomogeneous Trace Test

Assume that  $V$  is not a product. Given nonzero adjacent multidegrees  $d_{\alpha+1,\beta}$  and  $d_{\alpha,\beta+1}$ ,  $L' \subset \mathbb{P}^A$  and  $M' \subset \mathbb{P}^B$  of codimensions  $\alpha$  and  $\beta$  containing hyperplanes  $L \subset L'$  and  $M \subset M'$ , then

$$W_{10} := V \cap (L \times M') \text{ and } W_{01} := V \cap (L' \times M)$$



are the corresponding multihomogeneous witness sets.

Then  $C := V \cap (L' \times M')$  is an irreducible curve with multidegrees  $d_{10} = d_{\alpha+1,\beta}$  and  $d_{01} = d_{\alpha,\beta+1}$  with witness sets  $W_{10}$  and  $W_{01}$ .

Working in an affine patch  $\mathbb{C}^n \oplus \mathbb{C}^m$  on  $L' \times M'$ ,  $C$  has degree  $d_{10} + d_{01}$  and  $W_{01} \cup W_{10}$  can be used to get a witness set  $W = C \cap H$ , which we may use for a trace test in the affine space  $\mathbb{C}^n \oplus \mathbb{C}^m$ .

# Example

Suppose that  $C \subset \mathbb{P}^1 \times \mathbb{P}^1$  is defined locally by  $y^2x = 1$ .



Left: Linear spaces  $x = x_0$  and  $y = y_0$ , line  $H : h = 0$ , and the curve  $g(\frac{1}{2})$ , where  $g(t) := (x-x_0)(y-y_0)(1-t) + th$ . These are  $g(t)$  at  $t = 0, \frac{1}{2}, 1$ .

Right: the parallel slices  $h = \tau$  are in green, and the averages of witness points ( $\frac{1}{3}$  of the trace) lies on the brown line.