

Name _____

MATH 221

Exam 1, Version B

Spring 2024

501

Solutions

P. Yasskin

Multiple Choice: (6 points each. No part credit.)

1-8	/48	11	/12
9	/20	12	/12
10	/12	Total	/104






1. A point has spherical coordinates $(\rho, \phi, \theta) = \left(6, \frac{\pi}{6}, \frac{\pi}{4}\right)$. Find its cylindrical coordinates.

- a. $(r, \theta, z) = \left(3, \frac{\pi}{6}, 3\sqrt{3}\right)$
- b. $(r, \theta, z) = \left(3, \frac{\pi}{6}, 6\sqrt{3}\right)$
- c. $(r, \theta, z) = \left(3, \frac{\pi}{4}, 3\sqrt{3}\right)$ Correct
- d. $(r, \theta, z) = \left(6\sqrt{3}, \frac{\pi}{6}, 3\right)$
- e. $(r, \theta, z) = \left(3\sqrt{3}, \frac{\pi}{4}, 3\right)$

Solution:

$$z = \rho \cos \phi = 6 \cos \frac{\pi}{6} = 6 \frac{\sqrt{3}}{2} = 3\sqrt{3} \quad r = \rho \sin \phi = 6 \frac{1}{2} = 3 \quad (r, \theta, z) = \left(3, \frac{\pi}{4}, 3\sqrt{3}\right)$$

2. Which of the following is the contour plot of the function $f(x, y) = (x - 2)^4 + (y - 3)^4$?



Solution: The function is everywhere positive except at $(x, y) = (2, 3)$ where it is 0. So $(2, 3)$ is a local minimum. Contours circle around a minimum or maximum. So **(b)** is the answer.

3. A hiker starts at the point $P = (4, 2)$, travels along the vector $\vec{a} = \langle 2, -2 \rangle$, then along the vector $\vec{b} = \langle 1, 3 \rangle$ and finally along the vector $\vec{c} = \langle -1, 2 \rangle$. Along what vector should the hiker travel to get back to the starting point P ?
- $\langle 2, 3 \rangle$
 - $\langle -6, -5 \rangle$
 - $\langle 6, 5 \rangle$
 - $\langle -2, -3 \rangle$ Correct
 - $\langle 2, -1 \rangle$

Solution: In total the hiker travels along the vector $\vec{v} = \vec{a} + \vec{b} + \vec{c} = \langle 2, -2 \rangle + \langle 1, 3 \rangle + \langle -1, 2 \rangle = \langle 2, 3 \rangle$. To get back the hiker must travel along $-\vec{v} = \langle -2, -3 \rangle$. The point P is irrelevant.

4. For what value of p is $\vec{u} = \langle p, 5, 3 \rangle$ perpendicular to $\vec{v} = \langle 2, 1, p \rangle$?
- $p = 2$
 - $p = 1$
 - $p = 0$
 - $p = -1$ Correct
 - $p = -2$

Solution: They are perpendicular if their dot product is 0.

$$\vec{u} \cdot \vec{v} = \langle p, 5, 3 \rangle \cdot \langle 2, 1, p \rangle = 2p + 5 + 3p = 5p + 5 = 0 \quad \text{for } p = -1$$

5. Find the volume of the parallelepiped with edge vectors $\vec{a} = \langle 4, 2, 0 \rangle$, $\vec{b} = \langle 1, 0, -3 \rangle$ and $\vec{c} = \langle 0, -1, 2 \rangle$.
- 16 Correct
 - 12
 - 8
 - 12
 - 16

$$\text{Solution: } \vec{a} \cdot \vec{b} \times \vec{c} = \begin{vmatrix} 4 & 2 & 0 \\ 1 & 0 & -3 \\ 0 & -1 & 2 \end{vmatrix} = 4(0 - 3) - 2(2 - 0) + 0(-1 - 0) = -16$$

$$\text{Volume} = |\vec{a} \cdot \vec{b} \times \vec{c}| = |-16| = 16$$

6. If \hat{T} points Up and \hat{B} points NorthEast, in what direction does \hat{N} point?

- a. Down
- b. NorthWest
- c. SouthEast Correct
- d. SouthWest
- e.

Solution: $\hat{N} = \hat{B} \times \hat{T}$ If the fingers of your right hand point NorthEast along \hat{B} and the palm faces Up toward \hat{T} , then your thumb points SouthEast along \hat{N} .

7. Which of the following is a plane perpendicular to the line $(x,y,z) = (1 + 3t, 3 + 2t, 4 - t)$?

- a. $3x + 2y - z = 7$ Correct
- b. $-3x + 2y + z = 2$
- c. $x + 3y + 4z = 5$
- d. $3x - 2y - z = 3$
- e. $x - 3y + 4z = 5$

Solution: The normal to the plane is the direction of the line: $\vec{N} = \vec{v} = \langle 3, 2, -1 \rangle$. Any plane with this normal has the form $\vec{N} \cdot X = \vec{N} \cdot P$, or $3x + 2y - z = D$.

8. Classify the quadratic surface: $-x^2 + 2x + y^2 + 4y - 2z^2 + 12z = 14$

- a. Hyperboloid of 1 sheet Correct
- b. Hyperboloid of 2 sheets
- c. Hyperbolic Paraboloid opening up in the x -direction and down in the y -direction
- d. Hyperbolic Paraboloid opening up in the y -direction and down in the x -direction
- e. Cone

Solution: We complete the squares on x , y and z :

$$-(x^2 - 2x) + (y^2 + 4y) - 2(z^2 - 6z) = 14$$

$$-(x^2 - 2x + 1) + (y^2 + 4y + 4) - 2(z^2 - 6z + 9) = 14 - 1 + 4 - 18 = -1$$

$$-(x - 1)^2 + (y + 2)^2 - 2(z - 3)^2 = -1$$

To get the standard form (with a 1 on the right), we multiply by -1 :

$$(x - 1)^2 - (y + 2)^2 + 2(z - 3)^2 = 1$$

This is a hyperboloid of 1 sheet.

Work Out: (Points indicated. Part credit possible. Show all work.)

9. (20 pts) Consider the twisted cubic $\vec{r} = (6t, 3t^2, t^3)$. Compute each of the following.

Note: $4 + 4t^2 + t^4 = (2 + t^2)^2$

- a. (6 pts) Arc length between $(0, 0, 0)$ and $(6, 3, 1)$.

Solution: $\vec{v} = \langle 6, 6t, 3t^2 \rangle$ $|\vec{v}| = \sqrt{36 + 36t^2 + 9t^4} = 3\sqrt{4 + 4t^2 + t^4} = 3\sqrt{(2 + t^2)^2} = 3(2 + t^2)$

$$L = \int_0^1 |\vec{v}| dt = \int_0^1 3(2 + t^2) dt = 3 \left[2t + \frac{t^3}{3} \right]_0^1 = 3 \left[2 + \frac{1}{3} \right] = 7$$

- b. (6 pts) Curvature $\kappa = \frac{|\vec{v} \times \vec{a}|}{|\vec{v}|^3}$.

HINT: Factor out an 18^2 .

Solution: $\vec{a} = \langle 0, 6, 6t \rangle$ $\vec{v} \times \vec{a} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 6 & 6t & 3t^2 \\ 0 & 6 & 6t \end{vmatrix} = \langle 36t^2 - 18t^2, -36t, 36 \rangle = \langle 18t^2, -36t, 36 \rangle$

$$|\vec{v} \times \vec{a}| = \sqrt{18^2 t^4 + 36^2 t^2 + 36^2} = 18\sqrt{t^4 + 4t^2 + 4} = 18\sqrt{(2 + t^2)^2} = 18(2 + t^2)$$

$$\kappa = \frac{|\vec{v} \times \vec{a}|}{|\vec{v}|^3} = \frac{18(2 + t^2)}{3^3(2 + t^2)^3} = \frac{2}{3(2 + t^2)^2}$$

- c. (4 pts) Tangential acceleration, a_T .

HINT: You do NOT need to compute \hat{T} , \hat{N} or \hat{B} .

Solution: $a_T = \frac{d}{dt} |\vec{v}| = \frac{d}{dt} 3(2 + t^2) = 6t$

- d. (4 pts) Normal acceleration, a_N .

HINT: You do NOT need to compute \hat{T} , \hat{N} or \hat{B} .

Solution: $a_n = \kappa |\vec{v}|^2 = \frac{2}{3(2 + t^2)^2} 3^2(2 + t^2)^2 = 6$

10. (12 pts) Write the vector, $\vec{a} = \langle 5, 1, -3 \rangle$, as a sum of two vectors \vec{p} and \vec{q} , where \vec{p} is parallel to $\vec{b} = \langle 6, 4, 2 \rangle$ and \vec{q} is perpendicular to \vec{b} .

Solution: $\vec{p} = \text{proj}_{\vec{b}} \vec{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \vec{b} = \frac{30 + 4 - 6}{36 + 16 + 4} \langle 6, 4, 2 \rangle = \frac{28}{56} \langle 6, 4, 2 \rangle = \frac{1}{2} \langle 6, 4, 2 \rangle = \langle 3, 2, 1 \rangle$

$$\vec{q} = \vec{a} - \vec{p} = \langle 5, 1, -3 \rangle - \langle 3, 2, 1 \rangle = \langle 2, -1, -4 \rangle$$

We check $\vec{p} + \vec{q} = \langle 3, 2, 1 \rangle + \langle 2, -1, -4 \rangle = \langle 5, 1, -3 \rangle = \vec{a}$

$\vec{p} \parallel \vec{b}$ because $\vec{p} = \langle 3, 2, 1 \rangle$ is a multiple of $\vec{b} = \langle 6, 4, 2 \rangle$.

$\vec{q} \perp \vec{b}$ because $\vec{q} \cdot \vec{b} = 12 - 4 - 8 = 0$

11. (12 pts) Consider the helix $\vec{r}(\theta) = \langle 3 \cos \theta, 3 \sin \theta, 4\theta \rangle$ for $0 \leq \theta \leq 2\pi$.

a. Find its mass, if its linear density is $\delta(x, y, z) = z$.

Solution: The velocity is $\vec{v} = \langle -3 \sin \theta, 3 \cos \theta, 4 \rangle$.

The speed is $|\vec{v}| = \sqrt{9 \sin^2 \theta + 9 \cos^2 \theta + 16} = 5$.

The density along the curve is $\delta = z = 4\theta$. So the mass is

$$M = \int_0^{2\pi} \delta |\vec{v}| dt = \int_0^{2\pi} 4\theta 5 d\theta = 20 \left[\frac{\theta^2}{2} \right]_0^{2\pi} = 10(4\pi^2) = 40\pi^2$$

b. Find the work done to push a bead along the helix if the force is $\vec{F} = \langle -2y, 2x, 0 \rangle$.

Solution: Along the curve the force is $\vec{F} = \langle -6 \sin \theta, 6 \cos \theta, 0 \rangle$. So the work is

$$W = \int_0^{2\pi} \vec{F} \cdot d\vec{s} = \int_0^{2\pi} \vec{F} \cdot \vec{v} d\theta = \int_0^{2\pi} (18 \sin^2 \theta + 18 \cos^2 \theta) d\theta = \int_0^{2\pi} 18 d\theta = 36\pi$$

12. (12 pts) Consider the planes:

$$P_1 : \quad x + 3y + 3z = 5$$

$$P_2 : \quad x + y - z = 3$$

Determine if they are parallel or intersecting. If they intersect, find the line of intersection. You MUST explain why they are or are not parallel.

Solution: The normal vectors are $\vec{N}_1 = \langle 1, 3, 3 \rangle$ and $\vec{N}_2 = \langle 1, 1, -1 \rangle$.

Since these are not multiples of each other, the planes are not parallel.

The direction of the line of intersection is the cross product of the normals:

$$\vec{v} = \vec{N}_1 \times \vec{N}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 3 & 3 \\ 1 & 1 & -1 \end{vmatrix} = \hat{i}(-3-3) - \hat{j}(-1-3) + \hat{k}(1-3) = \langle -6, 4, -2 \rangle$$

To find a point of intersection, we pick $z = 0$ and solve:

$$\begin{aligned} x + 3y &= 5 \\ x + y &= 3 \end{aligned} \quad \Rightarrow \quad 2y = 2 \quad y = 1 \quad x = 2 \quad P = (2, 1, 0)$$

The line is: $X = P + t\vec{v} \quad (x, y, z) = (2, 1, 0) + t\langle -6, 4, -2 \rangle = (2 - 6t, 1 + 4t, -2t)$

Other answers are possible.