

Name \_\_\_\_\_ Sec \_\_\_\_\_

MATH 221 Exam 1 Fall 2009

Section 503 P. Yasskin

Multiple Choice: (6 points each. No part credit.)

1-11	/66	13	/10
12	/20	14	/10
		Total	/106

1. If  $f(x,y) = x^2 \cos(y^2)$ , which of the following is FALSE?

- a.  $f_x(x,y) = 2x \cos(y^2)$
- b.  $f_y(x,y) = -2x^2 y \sin(y^2)$
- c.  $f_{xx}(x,y) = 2 \cos(y^2)$
- d.  $f_{yy}(x,y) = -4x^2 y \cos(y^2)$
- e.  $f_{xy}(x,y) = -4xy \sin(y^2)$

2. The quadratic surface  $x^2 - y^2 + z^2 - 4x - 6y - 10z + 16 = 0$  is a

- a. hyperboloid of 1 sheet and center  $(2, 3, 5)$
- b. hyperboloid of 1 sheet and center  $(2, -3, 5)$
- c. hyperboloid of 2 sheets and center  $(2, 3, 5)$
- d. hyperboloid of 2 sheets and center  $(2, -3, 5)$
- e. cone with vertex  $(2, 3, 5)$

3. An airplane is travelling due North at constant speed and a constant altitude as it crosses the equator. In what direction does the  $\hat{B}$  vector point?

HINTS: Remember the Earth is curved. Ignore the rotation of the Earth.

- a. East
- b. West
- c. South
- d. Up
- e. Down

4. A triangle has edge vectors  $\vec{AB} = (2, 1, -2)$  and  $\vec{AC} = (-2, -2, 4)$ . Find the altitude of the triangle if  $\overline{AB}$  is the base.

- a.  $\frac{2\sqrt{5}}{3}$
- b.  $\frac{\sqrt{5}}{3}$
- c.  $2\sqrt{5}$
- d.  $\sqrt{5}$
- e.  $3\sqrt{5}$

5. A box slides down the helical ramp  $\vec{r}(t) = (4\cos t, 4\sin t, 9 - 3t)$  starting at height  $z = 9$  and ending at height  $z = 0$ . How far does the box slide?

- a. 3
- b. 5
- c. 15
- d. 25
- e. 75

6. A box slides down the helical ramp  $\vec{r}(t) = (4\cos t, 4\sin t, 9 - 3t)$  starting at height  $z = 9$  and ending at height  $z = 0$  under the action of the force  $\vec{F} = (-yz, xz, 5z)$ .

Find the work done on the box.






- a.  $\frac{9}{2}$
- b. 9
- c.  $\frac{25}{2}$
- d.  $\frac{27}{2}$
- e. 27

7. The diameter and height of a cylindrical trash can (no lid) are measured as  $D = 30$  cm and  $h = 40$  cm. The metal is  $0.2$  cm thick. Use differentials to estimate the volume of metal used to make the can.
- $165\pi \text{ cm}^3$
  - $210\pi \text{ cm}^3$
  - $285\pi \text{ cm}^3$
  - $330\pi \text{ cm}^3$
  - $525\pi \text{ cm}^3$
8. Find the equation of the plane tangent to the surface  $z = x^3y^2$  at the point  $(2, 1)$ . Then the  $z$ -intercept is  $z =$
- $-40$
  - $8$
  - $-8$
  - $32$
  - $-32$
9. Find the equation of the plane tangent to the surface  $12xyz - z^3 = 45$  at the point  $(1, 2, 3)$ . Then the  $z$ -intercept is  $z =$
- $135$
  - $45$
  - $-\sqrt[3]{6}$
  - $-45$
  - $-135$

10. Starting from the point  $(1, -2)$ , find the maximum rate at which the function  $f(x, y) = x^2y^3$  increases.

- a. 20
- b. 25
- c. 400
- d.  $(-16, 12)$
- e.  $(16, -12)$

11. Which of the following is the plot of the vector field  $F(x, y) = (x + y, x - y)$ ?



Work Out: (Points indicated. Part credit possible. Show all work.)

12. (20 points) Find the point on the curve  $\vec{r}(t) = (e^t, \sqrt{2}t, e^{-t})$  where the curvature is a local maximum or local minimum. Is it a local maximum or local minimum?

HINTS: First find the curvature  $\kappa = \frac{|\vec{v} \times \vec{a}|}{|\vec{v}|^3}$ . Then find the critical point and apply the first or second derivative test.

13. (10 points) The pressure,  $P$ , density,  $D$ , and temperature,  $T$ , of a certain ideal gas are related by  $P = 4DT$ . A fly is currently at the point  $\vec{r}(t_0) = (3, 2, 4)$  and has velocity  $\vec{v}(t_0) = (2, 1, 2)$ . At the point  $(3, 2, 4)$ , the density and temperature and their gradients are

$$D = 50 \quad \vec{\nabla}D = \left( \frac{\partial D}{\partial x}, \frac{\partial D}{\partial y}, \frac{\partial D}{\partial z} \right) = (0.1, 0.4, 0.2)$$

$$T = 300 \quad \vec{\nabla}T = \left( \frac{\partial T}{\partial x}, \frac{\partial T}{\partial y}, \frac{\partial T}{\partial z} \right) = (2, 3, 1)$$

Find the time rate of change of the pressure,  $\frac{dP}{dt}$ , as seen by the fly.

14. (10 points) Determine whether or not each of these limits exists. If it exists, find its value.

a.  $\lim_{(x,y) \rightarrow (0,0)} \frac{3x^2y^2}{x^6 + 3y^3}$

b.  $\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^2}$