

Name \_\_\_\_\_

MATH 221 Final Exam Version A Fall 2019

Section 505 Solutions P. Yasskin

Multiple Choice: (5 points each. No part credit.)

1-11	/55	13	/10
12	/15	14	/25
		Total	/105

1. A triangle has vertices  $A = (3, 2, 1)$ ,  $B = (3, 3, 2)$  and  $C = (4, 4, 2)$ . Find the angle at  $A$ .

- a.  $90^\circ$
- b.  $60^\circ$
- c.  $45^\circ$
- d.  $30^\circ$  Correct Choice
- e.  $0^\circ$

**Solution:**  $\overrightarrow{AB} = B - A = (0, 1, 1)$      $\overrightarrow{AC} = C - A = (1, 2, 1)$      $|\overrightarrow{AB}| = \sqrt{2}$      $|\overrightarrow{AC}| = \sqrt{6}$   
 $\overrightarrow{AB} \cdot \overrightarrow{AC} = 0 + 2 + 1 = 3$      $\cos \theta = \frac{\overrightarrow{AB} \cdot \overrightarrow{AC}}{|\overrightarrow{AB}| |\overrightarrow{AC}|} = \frac{3}{\sqrt{2} \sqrt{6}} = \frac{\sqrt{3}}{2}$      $\theta = 30^\circ$

2. Find the area of the triangle with vertices  $A = (3, 2, 1)$ ,  $B = (3, 3, 2)$  and  $C = (4, 4, 2)$ .

- a.  $\frac{\sqrt{3}}{2}$  Correct Choice
- b.  $\sqrt{3}$
- c.  $\frac{\sqrt{6}}{2}$
- d.  $\sqrt{6}$
- e.  $2\sqrt{6}$

**Solution:**  $\overrightarrow{AB} = B - A = \langle 0, 1, 1 \rangle$      $\overrightarrow{AC} = C - A = \langle 1, 2, 1 \rangle$   
 $\overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 1 & 1 \\ 1 & 2 & 1 \end{vmatrix} = \hat{i}(1 - 2) - \hat{j}(0 - 1) + \hat{k}(0 - 1) = \langle -1, 1, -1 \rangle$      $A = \frac{\sqrt{3}}{2}$

3. Find the arc length of the curve  $\vec{r}(t) = (t^2, 2t, \ln t)$  between  $(1, 2, 0)$  and  $(e^2, 2e, 1)$ .

- a.  $e^2 + 1$
- b.  $e^2 - 1$
- c.  $e^2$       Correct Choice
- d.  $\sqrt{e^2 + 1}$
- e.  $\sqrt{e^2 + 1} - 1$

**Solution:**  $|\vec{v}| = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} = \sqrt{(2t)^2 + (2)^2 + (t^{-1})^2} = \sqrt{4t^2 + 4 + \frac{1}{t^2}} = 2t + \frac{1}{t}$   
 $L = \int_1^e ds = \int_1^e |\vec{v}| dt = \int_1^e \left(2t + \frac{1}{t}\right) dt = \left[t^2 + \ln t\right]_1^e = (e^2 + 1) - (1 - 0) = e^2$

4. Find the average value of the function  $f(x, y, z) = x$  along the curve  $\vec{r}(t) = (t^2, 2t, \ln t)$  between  $(1, 2, 0)$  and  $(e^2, 2e, 1)$ .

- a.  $\frac{e^2}{2} + \frac{1}{2} - \frac{1}{e^2}$       Correct Choice
- b.  $\frac{e^4}{2} + \frac{e^2}{2} - 1$
- c.  $\frac{e^2}{2} - \frac{1}{2} - \frac{1}{e^2}$
- d.  $\frac{e^4}{2} - \frac{e^2}{2} - 1$
- e.  $\frac{e^2}{2} - \frac{1}{e^2}$

**Solution:**  $f_{\text{ave}} = \frac{1}{L} \int_1^e f ds = \frac{1}{e^2} \int_1^e x |\vec{v}| dt = \frac{1}{e^2} \int_1^e t^2 \left(2t + \frac{1}{t}\right) dt = \frac{1}{e^2} \int_1^e (2t^3 + t) dt = \frac{1}{e^2} \left[ \frac{t^4}{2} + \frac{t^2}{2} \right]_1^e$   
 $= \left(\frac{e^2}{2} + \frac{1}{2}\right) - \frac{1}{e^2} \left(\frac{1}{2} + \frac{1}{2}\right) = \frac{e^2}{2} + \frac{1}{2} - \frac{1}{e^2}$

5. Find the plane tangent to  $x \sin y + 2z \cos y = 2$  at  $(x, y, z) = \left(\sqrt{2}, \frac{\pi}{4}, \frac{1}{\sqrt{2}}\right)$ . The  $z$ -intercept is:

- a. 0
- b.  $\frac{1}{2\sqrt{2}}$
- c.  $\frac{1}{\sqrt{2}}$
- d.  $\sqrt{2}$       Correct Choice
- e. 2

**Solution:** Let  $f = x \sin y + 2z \cos y$  and  $P = \left(\sqrt{2}, \frac{\pi}{4}, \frac{1}{\sqrt{2}}\right)$ . Then

$\vec{\nabla}f = \langle \sin y, x \cos y - 2z \sin y, 2 \cos y \rangle$  and

$$\vec{N} = \vec{\nabla}f|_P = \left\langle \frac{1}{\sqrt{2}}, \sqrt{2} \frac{1}{\sqrt{2}} - 2 \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}}, 2 \frac{1}{\sqrt{2}} \right\rangle = \left\langle \frac{1}{\sqrt{2}}, 0, \sqrt{2} \right\rangle.$$

So the tangent plane is  $\vec{N} \cdot X = \vec{N} \cdot P$  or  $\frac{1}{\sqrt{2}}x + \sqrt{2}z = \frac{1}{\sqrt{2}}\sqrt{2} + \sqrt{2} \frac{1}{\sqrt{2}} = 2$ .

The  $z$ -intercept occurs when  $x = y = 0$  or  $\sqrt{2}z = 2$  or  $z = \sqrt{2}$ .

6. The velocity field of the water in a sink is  $\vec{V} = \langle -x^2y, xy^2 \rangle$ . Find the circulation of the water,  $Circ = \oint \vec{V} \cdot d\vec{s}$ , counterclockwise around the circle  $x^2 + y^2 = 4$ .

HINT: Use Green's Theorem.

- a.  $64\pi$
- b.  $32\pi$
- c.  $8\pi$       Correct Choice
- d.  $\frac{16}{3}\pi$
- e.  $2\pi$

**Solution:**  $Circ = \oint \vec{V} \cdot d\vec{s} = \oint P dx + Q dy$  where  $P = -x^2y$  and  $Q = xy^2$ .

By Green's Theorem,  $Circ = \iint (\partial_x Q - \partial_y P) dx dy = \iint (y^2 - -x^2) dx dy = \iint (x^2 + y^2) dx dy$

In polar coordinates,  $Circ = \int_0^{2\pi} \int_0^2 r^2 r dr d\theta = 2\pi \left[ \frac{r^4}{4} \right]_0^2 = 8\pi$

7. Compute  $\int \vec{F} \cdot d\vec{s}$  for  $\vec{F} = \langle 4x^3, 3y^2, 2z \rangle$  along the curve

$$\vec{r}(t) = \left( t \sin\left(\frac{\pi}{2}t\right), 2^t \cos\left(\frac{\pi}{2}t\right), 2^t \sin\left(\frac{\pi}{2}t\right) \right) \text{ from } t = 0 \text{ to } t = 1.$$

HINT: Find a scalar potential.

- a. 4      Correct Choice
- b. 3
- c. 2
- d. 1
- e. 0

**Solution:** A scalar potential is  $f = x^4 + y^3 + z^2$ .

The endpoints are  $A = \vec{r}(0) = (0, 1, 0)$  and  $B = \vec{r}(1) = (1, 0, 2)$ .

By the FTCC,  $\int_A^B \vec{F} \cdot d\vec{s} = \int_A^B \nabla f \cdot d\vec{s} = f(B) - f(A) = (1^4 + 2^2) - (1^3) = 4$

8. Compute  $\iint_S \vec{\nabla} \times \vec{G} \cdot d\vec{S}$  for  $\vec{G} = \langle y, -x, z \rangle$  over the surface  $\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, 16 - r^4)$

for  $0 \leq r \leq 2$  and  $0 \leq \theta \leq 2\pi$  oriented up and out.

HINT: Use a theorem.

- a.  $4\pi$
- b.  $2\pi$
- c.  $-2\pi$
- d.  $-4\pi$
- e.  $-8\pi$       Correct Choice

**Solution:** The boundary curve ( $r = 2$ ) is  $\vec{r}(\theta) = (2 \cos \theta, 2 \sin \theta, 0)$ .

This correctly points counterclockwise. The velocity is  $\vec{v} = \langle -2 \sin \theta, 2 \cos \theta, 0 \rangle$ .

On the circle,  $\vec{G} = \langle y, -x, z \rangle = \langle 2 \sin \theta, -2 \cos \theta, 0 \rangle$ . By Stokes' Theorem:

$$\iint_S \vec{\nabla} \times \vec{G} \cdot d\vec{S} = \oint_{\partial S} \vec{G} \cdot d\vec{s} = \int_0^{2\pi} \vec{G} \cdot \vec{v} d\theta = \int_0^{2\pi} -4 \sin^2 \theta - 4 \cos^2 \theta d\theta = -8\pi$$

9. Find the centroid of the solid hemisphere  $0 \leq z \leq \sqrt{4 - x^2 - y^2}$ .

- a.  $(0, 0, 2\pi)$
- b.  $(0, 0, 4\pi)$
- c.  $\left(0, 0, \frac{3}{4}\right)$     Correct Choice
- d.  $\left(0, 0, \frac{4}{3}\right)$
- e.  $\left(0, 0, \frac{1}{2}\right)$

**Solution:** By symmetry  $\bar{x} = \bar{y} = 0$ .

In spherical coordinates,  $z = \rho \cos \varphi$  and  $dV = \rho^2 \sin \varphi d\rho d\varphi d\theta$ .

$$V_z = \iiint z dV = \int_0^{2\pi} \int_0^{\pi/2} \int_0^2 \rho \cos \varphi \rho^2 \sin \varphi d\rho d\varphi d\theta = 2\pi \left[ \frac{\sin^2 \varphi}{2} \right]_0^{\pi/2} \left[ \frac{\rho^4}{4} \right]_0^2 = 4\pi$$

$$V = \frac{1}{2} \cdot \frac{4}{3} \pi 2^3 = \frac{16}{3} \pi \quad \bar{z} = \frac{V_z}{V} = \frac{3 \cdot 4\pi}{16\pi} = \frac{3}{4}$$

10. Find the area of the surface  $\vec{R}(u, v) = (u, v, u + v)$  for  $0 \leq u \leq 2$  and  $0 \leq v \leq 3$ .

- a.  $3\sqrt{2}$
- b.  $6\sqrt{3}$     Correct Choice
- c.  $6\sqrt{2}$
- d.  $12\sqrt{3}$
- e.  $12\sqrt{2}$

**Solution:**  $\vec{e}_u = \langle 1, 0, 1 \rangle \quad \vec{e}_v = \langle 0, 1, 1 \rangle \quad \vec{N} = \vec{e}_u \times \vec{e}_v = \langle -1, -1, 1 \rangle \quad |\vec{N}| = \sqrt{3}$

$$A = \iint |\vec{N}| du dv = \int_0^3 \int_0^2 \sqrt{3} du dv = 6\sqrt{3}$$

11. Find the flux of  $\vec{F} = \langle y, -z, x \rangle$  upward thru the surface  $\vec{R}(u, v) = (u, v, u + v)$  for  $0 \leq u \leq 2$  and  $0 \leq v \leq 3$ .

- a. 0
- b. 12    Correct Choice
- c. 18
- d. 24
- e. 30

**Solution:**  $\vec{N} = \vec{e}_u \times \vec{e}_v = \langle -1, -1, 1 \rangle$  which is correctly upward.  $\vec{F} = \langle y, -z, x \rangle = \langle v, -u - v, u \rangle$

$$\iint \vec{F} \cdot d\vec{S} = \iint \vec{F} \cdot \vec{N} du dv = \int_0^3 \int_0^2 (-v - (-u - v) + u) du dv = \int_0^3 \int_0^2 2u du dv = [u^2]_0^3 [v]_0^3 = 12$$

Work Out: (Points indicated. Part credit possible. Show all work.)

12. (15 points) The Ideal Gas Law says the Pressure,  $P$ , Density,  $\delta$ , and Temperature,  $T$ , are related by  $P = k\delta T$  for some constant  $k$ . We will assume the atmosphere is an ideal gas with  $k = 2$ . A weather balloon measures the Density and Temperature to be

$$\delta = 0.1 \frac{\text{gm}}{\text{m}^3} \quad T = 270^\circ\text{K}$$

and their gradients to be

$$\vec{\nabla}\delta = \langle 0.03, 0.02, 0.01 \rangle \quad \vec{\nabla}T = \langle -1, 1, 3 \rangle$$

Find the gradient of the Pressure.

**Solution:**  $\frac{\partial P}{\partial \delta} = kT = 2 \cdot 270 = 540 \quad \frac{\partial P}{\partial T} = k\delta = 2 \cdot 0.1 = 0.2$

By the chain rule,

$$\begin{aligned} \frac{\partial P}{\partial x} &= \frac{\partial P}{\partial \delta} \frac{\partial \delta}{\partial x} + \frac{\partial P}{\partial T} \frac{\partial T}{\partial x} = 540 \cdot 0.03 + 0.2 \cdot (-1) = 16 \\ \frac{\partial P}{\partial y} &= \frac{\partial P}{\partial \delta} \frac{\partial \delta}{\partial y} + \frac{\partial P}{\partial T} \frac{\partial T}{\partial y} = 540 \cdot 0.02 + 0.2 \cdot 1 = 11 \\ \frac{\partial P}{\partial z} &= \frac{\partial P}{\partial \delta} \frac{\partial \delta}{\partial z} + \frac{\partial P}{\partial T} \frac{\partial T}{\partial z} = 540 \cdot 0.01 + 0.2 \cdot 3 = 6 \end{aligned}$$

So

$$\vec{\nabla}P = \langle 16, 11, 6 \rangle$$

13. (10 points) Find the largest value of  $f = xy^2z^3$  on the plane  $6x + 3y + 2z = 72$ .

**Solution 1:** Lagrange Multipliers: Let  $g = 6x + 3y + 2z$ .

Then  $\vec{\nabla}f = \langle y^2z^3, 2xyz^3, 3xy^2z^2 \rangle$  and  $\vec{\nabla}g = \langle 6, 3, 2 \rangle$ .

The Lagrange equations are:  $y^2z^3 = 6\lambda$ ,  $2xyz^3 = 3\lambda$ ,  $3xy^2z^2 = 2\lambda$

Then  $xy^2z^3 = 6\lambda x = \frac{3}{2}\lambda y = \frac{2}{3}\lambda z$  or  $y = 4x$  and  $z = 9x$

From the constraint,  $72 = 6x + 3y + 2z = 6x + 12x + 18x = 36x$ .

So  $x = 2$ ,  $y = 8$ ,  $z = 18$  and  $f = xy^2z^3 = 2 \cdot 8^2 \cdot 18^3$

**Solution 2:** Eliminate  $x = \frac{72 - 3y - 2z}{6} = 12 - \frac{y}{2} - \frac{z}{3}$ . Then

$$f = xy^2z^3 = \left(12 - \frac{y}{2} - \frac{z}{3}\right)y^2z^3 = 12y^2z^3 - \frac{1}{2}y^3z^3 - \frac{1}{3}y^2z^4$$

We find critical points:

$$f_y = 24yz^3 - \frac{3}{2}y^2z^3 - \frac{2}{3}yz^4 = \frac{1}{6}yz^3(144 - 9y - 4z) = 0$$

$$f_z = 36y^2z^2 - \frac{3}{2}y^3z^2 - \frac{4}{3}y^2z^3 = \frac{1}{6}y^2z^2(216 - 9y - 8z) = 0$$

Since  $y \neq 0$  and  $z \neq 0$ , we need to solve  $9y + 4z = 144$  and  $9y + 8z = 216$

We subtract  $4z = 72$  or  $z = 18$


Substituting back gives:  $9y + 72 = 144 \quad 9y = 72 \quad y = 8$

and  $x = 12 - \frac{y}{2} - \frac{z}{3} = 12 - 4 - 6 = 2$ . So  $f = xy^2z^3 = 2 \cdot 8^2 \cdot 18^3$

14. (25 points) Verify Gauss' Theorem  $\iiint_V \vec{\nabla} \cdot \vec{F} dV = \iint_{\partial V} \vec{F} \cdot d\vec{S}$

for the vector field  $\vec{F} = (xz, yz, 3z\sqrt{x^2 + y^2})$  and the solid cone  $\sqrt{x^2 + y^2} \leq z \leq 2$ .

Be careful with orientations. Use the following steps:



**First the Left Hand Side:**

- a. Compute the divergence:

$$\vec{\nabla} \cdot \vec{F} = z + z + 3\sqrt{x^2 + y^2}$$

- b. Express the divergence and the volume element in the appropriate coordinate system:

$$\vec{\nabla} \cdot \vec{F} = 2z + 3r \quad dV = r dr d\theta dz$$

- c. Compute the left hand side:

$$\begin{aligned} \iiint_V \vec{\nabla} \cdot \vec{F} dV &= \int_0^2 \int_0^{2\pi} \int_r^2 (2z + 3r) r dz d\theta dr = 2\pi \int_0^2 [z^2 + 3rz]_{z=r}^2 r dr \\ &= 2\pi \int_0^2 [(4 + 6r) - (r^2 + 3r^2)] r dr = 2\pi \int_0^2 (4r + 6r^2 - 4r^3) dr = 2\pi [2r^2 + 2r^3 - r^4]_0^2 \\ &= 2\pi(8 + 16 - 16) = 16\pi \end{aligned}$$

**Second the Right Hand Side:**

The boundary surface consists of a hemisphere  $H$  and a disk  $D$  with appropriate orientations.

The disk  $D$  may be parametrized as:  $\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, 2)$

- d. Compute the tangent vectors:

$$\vec{e}_r = (\cos \theta, \sin \theta, 0)$$

$$\vec{e}_\theta = (-r \sin \theta, r \cos \theta, 0)$$

- e. Compute the normal vector:

$$\vec{N} = \hat{i}(0) - \hat{j}(0) + \hat{k}(r \cos^2 \theta - r \sin^2 \theta) = (0, 0, r)$$

This is up as required.

- f. Evaluate  $\vec{F} = (xz, yz, 3z\sqrt{x^2 + y^2})$  on the disk:

$$\vec{F}|_{\vec{R}(r,\theta)} = (2r \cos \theta, 2r \sin \theta, 6r)$$

- g. Compute the dot product:

$$\vec{F} \cdot \vec{N} = 6r^2$$

- h. Compute the flux through  $D$ :

$$\iint_D \vec{F} \cdot d\vec{S} = \int_0^{2\pi} \int_0^2 6r^2 dr d\theta = 2\pi [2r^3]_0^2 = 32\pi$$

The cone  $C$  may be parametrized as:  $\vec{R}(r, \theta) = (r\cos\theta, r\sin\theta, r)$

- i. Compute the tangent vectors:

$$\vec{e}_r = (\cos\theta, \sin\theta, 1)$$

$$\vec{e}_\theta = (-r\sin\theta, r\cos\theta, 0)$$

- j. Compute the normal vector:

$$\begin{aligned}\vec{N} &= \hat{i}(-r\cos\theta) - \hat{j}(r\sin\theta) + \hat{k}(r\cos^2\theta + r\sin^2\theta) \\ &= (-r\cos\theta, -r\sin\theta, r)\end{aligned}$$

This is oriented up and in. We need down and out. Reverse it:

$$\vec{N} = (r\cos\theta, r\sin\theta, -r)$$

- k. Evaluate  $\vec{F} = (xz, yz, 3z\sqrt{x^2 + y^2})$  on the cone:

$$\vec{F} \Big|_{\vec{R}(r,\theta)} = (r^2\cos\theta, r^2\sin\theta, 3r^2)$$

- l. Compute the dot product:

$$\vec{F} \cdot \vec{N} = r^3\cos^2\theta + r^3\sin^2\theta - 3r^3 = -2r^3$$

- m. Compute the flux through  $C$ :

$$\iint_C \vec{F} \cdot d\vec{S} = \int_0^{2\pi} \int_0^2 -2r^3 dr d\theta = -2\pi \left[ \frac{r^4}{2} \right]_0^2 = -16\pi = -16\pi$$

- n. Compute the **TOTAL** right hand side:

$$\iint_{\partial V} \vec{F} \cdot d\vec{S} = \iint_D \vec{F} \cdot d\vec{S} + \iint_C \vec{F} \cdot d\vec{S} = 32\pi - 16\pi = 16\pi \quad \text{which agrees with (c).}$$