

## Vector Analysis Theorems

1. The **Fundamental Theorem of Calculus for Curves** states that if  $\vec{r}(t)$  is a nice curve in  $\mathbf{R}^n$  and  $f$  is a nice function in  $\mathbf{R}^n$  then

$$\int_{\vec{r}(A)}^{\vec{r}(B)} \vec{\nabla} f \cdot d\vec{s} = f(B) - f(A)$$

2. **Green's Theorem** states that if  $R$  is a nice region in  $\mathbf{R}^2$  and  $\partial R$  is its boundary curve traversed counterclockwise and  $P$  and  $Q$  are nice functions on  $R$  then

$$\iint_R \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy = \oint_{\partial R} P dx + Q dy$$

- a. **2D Stokes' Theorem** states that if  $R$  is a nice region in  $\mathbf{R}^2$  and  $\partial R$  is its boundary curve traversed counterclockwise and  $\vec{F} = (P(x,y), Q(x,y), 0)$  is a nice vector field on  $R$  then

$$\iint_R \vec{\nabla} \times \vec{F} \cdot \hat{k} dx dy = \oint_{\partial R} \vec{F} \cdot d\vec{s}$$

- b. **2D Gauss' Theorem** states that if  $R$  is a nice region in  $\mathbf{R}^2$  and  $\partial R$  is its boundary curve traversed counterclockwise and  $\vec{G} = (Q(x,y), -P(x,y), 0)$  is a nice vector field on  $R$  then

$$\iint_R \vec{\nabla} \cdot \vec{G} dx dy = \oint_{\partial R} \vec{G} \cdot d\vec{n}$$

3. **Stokes' Theorem** states that if  $S$  is a nice surface in  $\mathbf{R}^3$  and  $\partial S$  is its boundary curve traversed counterclockwise as seen from the tip of the normal to  $S$  and  $\vec{F}$  is a nice vector field on  $S$  then

$$\iint_S \vec{\nabla} \times \vec{F} \cdot d\vec{S} = \oint_{\partial S} \vec{F} \cdot d\vec{s}$$

4. **Gauss' Theorem** states that if  $V$  is a volume in  $\mathbf{R}^3$  and  $\partial V$  is its boundary surface oriented outward from  $V$  and  $\vec{F}$  is a nice vector field on  $V$  then

$$\iiint_V \vec{\nabla} \cdot \vec{F} dV = \iint_{\partial V} \vec{F} \cdot d\vec{S}$$