

Name _____ ID _____

1-3	/15
4	/10
Total	/25

MATH 251 Quiz 5 Spring 2007
Sections 509 Solutions P. Yasskin

Multiple Choice: (5 points each)


1. Find the volume of the solid below the paraboloid
- $z = 9 - x^2 - y^2$
- above the
- xy
- plane.

- a. $\frac{3}{2}\pi$
 b. 3π
 c. $\frac{9}{2}\pi$
 d. $\frac{27}{2}\pi$
 e. $\frac{81}{2}\pi$ Correct Choice

$$V = \iint_R (9 - x^2 - y^2) dA = \int_0^{2\pi} \int_0^3 (9 - r^2) r dr d\theta = 2\pi \int_0^3 (9r - r^3) dr = 2\pi \left[\frac{9r^2}{2} - \frac{r^4}{4} \right]_{r=0}^3 = 2\pi \left(\frac{81}{2} - \frac{81}{4} \right) = \frac{81}{2}\pi$$

2. Find the center of mass of the circle
- $r = 3 \sin \theta$
-
- if the mass surface density is
- $\rho = y$
- .

- a. $(0, \frac{8}{15})$
 b. $(0, \frac{15}{8})$ Correct Choice
 c. $(0, \frac{9}{4})$
 d. $(0, \frac{4}{9})$
 e. $(0, \frac{405}{64}\pi)$



HINTS: $\int_0^\pi \sin^4 \theta d\theta = \int_0^\pi \cos^4 \theta d\theta = \frac{3}{8}\pi$ $\int_0^{2\pi} \sin^4 \theta d\theta = \int_0^{2\pi} \cos^4 \theta d\theta = \frac{3}{4}\pi$
 $\int_0^\pi \sin^6 \theta d\theta = \int_0^\pi \cos^6 \theta d\theta = \frac{5}{16}\pi$ $\int_0^{2\pi} \sin^6 \theta d\theta = \int_0^{2\pi} \cos^6 \theta d\theta = \frac{5}{8}\pi$

$$\rho = y = r \sin \theta \quad \bar{x} = 0 \text{ by symmetry.}$$

$$M = \iint_R \rho dA = \int_0^\pi \int_0^{3 \sin \theta} (r \sin \theta) r dr d\theta = \int_0^\pi \left[\frac{r^3}{3} \right]_{r=0}^{3 \sin \theta} \sin \theta d\theta = 9 \int_0^\pi \sin^4 \theta d\theta = 9 \cdot \frac{3}{8}\pi = \frac{27}{8}\pi$$

$$\begin{aligned} y\text{-mom} &= \iint_R y\rho dA = \int_0^\pi \int_0^{3 \sin \theta} r \sin \theta (r \sin \theta) r dr d\theta = \int_0^\pi \left[\frac{r^4}{4} \right]_{r=0}^{3 \sin \theta} \sin^2 \theta d\theta \\ &= \frac{81}{4} \int_0^\pi \sin^6 \theta d\theta = \frac{81}{4} \cdot \frac{5}{16}\pi = \frac{405}{64}\pi \end{aligned}$$

$$\bar{y} = \frac{y\text{-mom}}{M} = \frac{405\pi}{64} \cdot \frac{8}{27\pi} = \frac{15}{8}$$

3. Compute $\int_0^2 \int_0^{\sqrt{4-x^2}} e^{x^2+y^2} dy dx$


a. $\frac{\pi}{2}(e^4 - 1)$

b. $\frac{\pi}{2}e^4$

c. $\frac{\pi}{4}(e^4 - 1)$ Correct Choice

d. $\frac{\pi}{4}e^4$

e. $\frac{\pi}{2}e^3$




$$\int_0^{\pi/2} \int_0^2 e^{r^2} r dr d\theta = \int_0^{\pi/2} \left[\frac{e^{r^2}}{2} \right]_0^2 d\theta = \int_0^{\pi/2} \frac{e^4 - 1}{2} d\theta = \frac{\pi}{4}(e^4 - 1)$$

4. Compute $\iint_R y^2 dx dy$ over the diamond shaped region R bounded by

$$y = \frac{1}{x}, \quad y = \frac{6}{x}, \quad y = x, \quad y = 2x$$

FULL CREDIT for integrating in the curvilinear coordinates (u, v) where $u^2 = xy$ and $v^2 = \frac{y}{x}$.
(Solve for x and y .)

HALF CREDIT for integrating in rectangular coordinates.



$$\begin{cases} u^2 = xy \\ v^2 = \frac{y}{x} \end{cases} \Rightarrow \begin{cases} u^2 v^2 = y^2 \\ \frac{u^2}{v^2} = x^2 \end{cases} \Rightarrow \begin{cases} x = \frac{u}{v} \\ y = uv \end{cases}$$

$$J = \left| \left| \frac{\partial(x,y)}{\partial(u,v)} \right| \right| = \left| \begin{array}{cc} \frac{1}{v} & -\frac{u}{v^2} \\ v & u \end{array} \right| = \left| \frac{u}{v} - \frac{u}{v} \right| = \frac{2u}{v}$$

$$xy = 1 \Rightarrow u^2 = 1 \Rightarrow u = 1$$

$$xy = 6 \Rightarrow u^2 = 6 \Rightarrow u = \sqrt{6}$$

$$\text{So: } 1 \leq u \leq \sqrt{6}$$

$$\frac{y}{x} = 1 \Rightarrow v^2 = 1 \Rightarrow v = 1$$

$$\frac{y}{x} = 2 \Rightarrow v^2 = 2 \Rightarrow v = \sqrt{2}$$

$$\text{So: } 1 \leq v \leq \sqrt{2}$$

$$\begin{aligned} \iint_R y^2 dx dy &= \int_1^{\sqrt{2}} \int_1^{\sqrt{6}} u^2 v^2 \frac{2u}{v} du dv = 2 \int_1^{\sqrt{2}} \int_1^{\sqrt{6}} u^3 v du dv \\ &= 2 \left[\frac{u^4}{4} \right]_{u=1}^{\sqrt{6}} \left[\frac{v^2}{2} \right]_{v=1}^{\sqrt{2}} = 2 \left[\frac{36}{4} - \frac{1}{4} \right] \left[\frac{2}{2} - \frac{1}{2} \right] = \frac{35}{4} \end{aligned}$$