

Name _____

Math 311 Exam 3 Version A Spring 2015

Section 503 P. Yasskin

Points indicated. Show all work.

1	/20	3	/30
2	/36	4	/26
		Total	/112

1. (20 points) Compute $\iint_S \vec{\nabla} \times \vec{F} \cdot d\vec{S}$ for $\vec{F} = (-y, x, z)$


over the "clam shell" surface, S , parametrized by

$$\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, r \sin(5\theta))$$

for $r \leq 3$ oriented upward.

HINTS: Use Stokes Theorem.

What is the value of r on the boundary?



2. (36 points) Let $V = \text{Span}(e^{2x} + e^{-2x}, e^{2x} - e^{-2x})$ be the vector space of functions spanned by the basis

$$e_1 = e^{2x} + e^{-2x}, \quad e_2 = e^{2x} - e^{-2x}$$

Consider the linear operator $L : V \rightarrow V$ given by $L(f) = 3 \frac{df}{dx}$. Our goals are to compute the eigenvalues and eigenfunctions of the linear operator L , to find the similarity transformation which diagonalizes the matrix of L and use this similarity transformation to compute a matrix power.

- a. (5 pts) Find the matrix of L relative to the (e_1, e_2) basis. Call it $A_{e \leftarrow e}$.

- b. (3 pts) Find the characteristic polynomial for $A_{e \leftarrow e}$.

Factor it and identify the eigenvalues of $A_{e \leftarrow e}$. These are also the eigenvalues of L .

- c. (8 pts) Find the eigenvector(s) of $A_{e \leftarrow e}$ for each eigenvalue, as vectors in \mathbb{R}^2 .

Name them \vec{v}_1 and \vec{v}_2 .

- d. (6 pts) Convert the eigenvectors of $A_{e \leftarrow e}$ into eigenfunctions of L as functions in V .

Name them f_1 and f_2 and simplify them.

Then compute $L(f_1)$ and $L(f_2)$ to verify f_1 and f_2 are eigenfunctions.

Hint: Remember that the components of \vec{v}_1 and \vec{v}_2 are components of f_1 and f_2 relative to the (e_1, e_2) basis.

- e. (3 pts) Using the eigenfunctions as a new (f_1, f_2) basis for V , find the matrix of L relative to the (f_1, f_2) basis. Call it D .
- f. (5 pts) Find the change of basis matrices $C_{e \leftarrow f}$ and $C_{f \leftarrow e}$ between the (e_1, e_2) basis to the (f_1, f_2) bases. Be sure to identify which is which.
- g. (2 pts) A and D are related by a similarity transformation $A = S^{-1}DS$. Identify S as $C_{e \leftarrow f}$ or $C_{f \leftarrow e}$.
- h. (4 pts) Compute A^{12} and A^{25} .

3. (30 points) The density, ρ , of an ideal gas is related to its pressure, P , and its absolute temperature, T , by the equation $\rho = \frac{P}{kT}$ where k is a constant which depends on the particular ideal gas. We are considering an ideal gas for which $k = 10^{-4} \text{ atm}\cdot\text{m}^3/\text{kg}/^\circ\text{K}$. At the current time, $t = t_0$, a flying robotic nanobot is located at $(x, y, z) = (2, 1, 3)^\top \text{ m}$ and has velocity $\vec{v} = (.4, .5, .2)^\top \text{ m/sec}$. The nanobot measures the current pressure is $P = 2 \text{ atm}$ while its gradient is $\vec{\nabla}P = (-.03, .01, .02) \text{ atm/m}$. Similarly, the nanobot measures the current temperature is $T = 250 \text{ }^\circ\text{K}$ while its gradient is $\vec{\nabla}T = (3, -2, -4) \text{ }^\circ\text{K/m}$.

a. (2 pts) Find the current density, ρ .

b. (6 pts) Find the Jacobian matrix of the density $\frac{D(\rho)}{D(P, T)}$ in general (in terms of symbols like $\frac{\partial \rho}{\partial T}$), then in terms of P and T , and finally at the current time $t = t_0$.


c. (4 pts) Find the Jacobian matrix $\frac{D(P, T)}{D(x, y, z)}$ in general (in terms of symbols like $\frac{\partial P}{\partial y}$) and then at the current time $t = t_0$.

d. (4 pts) Find the Jacobian matrix $\frac{D(x, y, z)}{D(t)}$ in general and then at $t = t_0$.

e. (6 pts) Find the time rate of change of the pressure as seen by the nanobot, at the current time $t = t_0$. Is the pressure currently increasing or decreasing?

f. (8 pts) Find the time rate of change of the density as seen by the nanobot, at the current time $t = t_0$. Is the density currently increasing or decreasing?

4. (26 points) Compute the integral $\iint x dA$ over the region in the first quadrant bounded by $y = 1 + x^2$, $y = 3 + x^2$, $y = 4 - x^2$, and $y = 5 - x^2$.



- a. (4 pts) Define the curvilinear coordinates u and v by $y = u + x^2$ and $y = v - x^2$. What are the 4 boundaries in terms of u and v ?
- b. (4 pts) Solve for x and y in terms of u and v . Express the results as a position vector.
- c. (4 pts) Find the coordinate tangent vectors:
- d. (8 pts) Compute the Jacobian factor:
- e. (6 pts) Compute the integral: