

MATH 304
Linear Algebra

Lecture 31:
Eigenvalues and eigenvectors (continued).

Eigenvalues and eigenvectors of a matrix

Definition. Let A be an $n \times n$ matrix. A number $\lambda \in \mathbb{R}$ is called an **eigenvalue** of the matrix A if $A\mathbf{v} = \lambda\mathbf{v}$ for a nonzero column vector $\mathbf{v} \in \mathbb{R}^n$.

The vector \mathbf{v} is called an **eigenvector** of A belonging to (or associated with) the eigenvalue λ .

If λ is an eigenvalue of A then the nullspace $N(A - \lambda I)$, which is nontrivial, is called the **eigenspace** of A corresponding to λ . The eigenspace consists of all eigenvectors belonging to the eigenvalue λ plus the zero vector.

Characteristic equation

Definition. Given a square matrix A , the equation $\det(A - \lambda I) = 0$ is called the **characteristic equation** of A .

Eigenvalues λ of A are roots of the characteristic equation.

If A is an $n \times n$ matrix then $p(\lambda) = \det(A - \lambda I)$ is a polynomial of degree n . It is called the **characteristic polynomial** of A .

Theorem Any $n \times n$ matrix has at most n eigenvalues.

Example. $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$

$$\begin{aligned} \det(A - \lambda I) &= \begin{vmatrix} a - \lambda & b \\ c & d - \lambda \end{vmatrix} \\ &= (a - \lambda)(d - \lambda) - bc \\ &= \lambda^2 - (a + d)\lambda + (ad - bc). \end{aligned}$$

Example. $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}.$

$$\det(A - \lambda I) = \begin{vmatrix} a_{11} - \lambda & a_{12} & a_{13} \\ a_{21} & a_{22} - \lambda & a_{23} \\ a_{31} & a_{32} & a_{33} - \lambda \end{vmatrix}$$
$$= -\lambda^3 + c_1\lambda^2 - c_2\lambda + c_3,$$

where $c_1 = a_{11} + a_{22} + a_{33}$ (the *trace* of A),

$$c_2 = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} + \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} + \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix},$$

$$c_3 = \det A.$$

Example. $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$.

Characteristic equation: $\begin{vmatrix} 2 - \lambda & 1 \\ 1 & 2 - \lambda \end{vmatrix} = 0$.

$$(2 - \lambda)^2 - 1 = 0 \implies \lambda_1 = 1, \lambda_2 = 3.$$

$$(A - I)\mathbf{x} = \mathbf{0} \iff \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\iff \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \iff x + y = 0.$$

The general solution is $(-t, t) = t(-1, 1)$, $t \in \mathbb{R}$.

Thus $\mathbf{v}_1 = (-1, 1)$ is an eigenvector associated with the eigenvalue 1. The corresponding eigenspace is the line spanned by \mathbf{v}_1 .

$$(A - 3I)\mathbf{x} = \mathbf{0} \iff \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$
$$\iff \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \iff x - y = 0.$$

The general solution is $(t, t) = t(1, 1)$, $t \in \mathbb{R}$.

Thus $\mathbf{v}_2 = (1, 1)$ is an eigenvector associated with the eigenvalue 3. The corresponding eigenspace is the line spanned by \mathbf{v}_2 .

Summary. $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$.

- The matrix A has two eigenvalues: 1 and 3.
- The eigenspace of A associated with the eigenvalue 1 is the line $t(-1, 1)$.
- The eigenspace of A associated with the eigenvalue 3 is the line $t(1, 1)$.
- Eigenvectors $\mathbf{v}_1 = (-1, 1)$ and $\mathbf{v}_2 = (1, 1)$ of the matrix A form an orthogonal basis for \mathbb{R}^2 .
- Geometrically, the mapping $\mathbf{x} \mapsto A\mathbf{x}$ is a stretch by a factor of 3 away from the line $x + y = 0$ in the orthogonal direction.

Example. $A = \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix}.$

Characteristic equation:

$$\begin{vmatrix} 1 - \lambda & 1 & -1 \\ 1 & 1 - \lambda & 1 \\ 0 & 0 & 2 - \lambda \end{vmatrix} = 0.$$

Expand the determinant by the 3rd row:

$$(2 - \lambda) \begin{vmatrix} 1 - \lambda & 1 \\ 1 & 1 - \lambda \end{vmatrix} = 0.$$

$$\begin{aligned} ((1 - \lambda)^2 - 1)(2 - \lambda) &= 0 \iff -\lambda(2 - \lambda)^2 = 0 \\ \implies \lambda_1 &= 0, \quad \lambda_2 = 2. \end{aligned}$$

$$A\mathbf{x} = \mathbf{0} \iff \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

Convert the matrix to reduced row echelon form:

$$\begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & -1 \\ 0 & 0 & 2 \\ 0 & 0 & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$A\mathbf{x} = \mathbf{0} \iff \begin{cases} x + y = 0, \\ z = 0. \end{cases}$$

The general solution is $(-t, t, 0) = t(-1, 1, 0)$, $t \in \mathbb{R}$. Thus $\mathbf{v}_1 = (-1, 1, 0)$ is an eigenvector associated with the eigenvalue 0. The corresponding eigenspace is the line spanned by \mathbf{v}_1 .

$$(A - 2I)\mathbf{x} = \mathbf{0} \iff \begin{pmatrix} -1 & 1 & -1 \\ 1 & -1 & 1 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\iff \begin{pmatrix} 1 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \iff x - y + z = 0.$$

The general solution is $x = t - s$, $y = t$, $z = s$, where $t, s \in \mathbb{R}$. Equivalently,

$$\mathbf{x} = (t - s, t, s) = t(1, 1, 0) + s(-1, 0, 1).$$

Thus $\mathbf{v}_2 = (1, 1, 0)$ and $\mathbf{v}_3 = (-1, 0, 1)$ are eigenvectors associated with the eigenvalue 2.

The corresponding eigenspace is the plane spanned by \mathbf{v}_2 and \mathbf{v}_3 .

Summary. $A = \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix}.$

- The matrix A has two eigenvalues: 0 and 2.
- The eigenvalue 0 is *simple*: the corresponding eigenspace is a line.
- The eigenvalue 2 is of *multiplicity* 2: the corresponding eigenspace is a plane.
- Eigenvectors $\mathbf{v}_1 = (-1, 1, 0)$, $\mathbf{v}_2 = (1, 1, 0)$, and $\mathbf{v}_3 = (-1, 0, 1)$ of the matrix A form a basis for \mathbb{R}^3 .
- Geometrically, the map $\mathbf{x} \mapsto A\mathbf{x}$ is the projection on the plane $\text{Span}(\mathbf{v}_2, \mathbf{v}_3)$ along the lines parallel to \mathbf{v}_1 with the subsequent scaling by a factor of 2.

Eigenvalues and eigenvectors of an operator

Definition. Let V be a vector space and $L : V \rightarrow V$ be a linear operator. A number λ is called an **eigenvalue** of the operator L if $L(\mathbf{v}) = \lambda\mathbf{v}$ for a nonzero vector $\mathbf{v} \in V$. The vector \mathbf{v} is called an **eigenvector** of L associated with the eigenvalue λ . (If V is a functional space then eigenvectors are also called **eigenfunctions**.)

If $V = \mathbb{R}^n$ then the linear operator L is given by $L(\mathbf{x}) = A\mathbf{x}$, where A is an $n \times n$ matrix.

In this case, eigenvalues and eigenvectors of the operator L are precisely eigenvalues and eigenvectors of the matrix A .

Eigenspaces

Let $L : V \rightarrow V$ be a linear operator.

For any $\lambda \in \mathbb{R}$, let V_λ denotes the set of all solutions of the equation $L(\mathbf{x}) = \lambda\mathbf{x}$.

Then V_λ is a *subspace* of V since V_λ is the *kernel* of a linear operator given by $\mathbf{x} \mapsto L(\mathbf{x}) - \lambda\mathbf{x}$.

V_λ minus the zero vector is the set of all eigenvectors of L associated with the eigenvalue λ . In particular, $\lambda \in \mathbb{R}$ is an eigenvalue of L if and only if $V_\lambda \neq \{\mathbf{0}\}$.

If $V_\lambda \neq \{\mathbf{0}\}$ then it is called the **eigenspace** of L corresponding to the eigenvalue λ .

Example. $V = C^\infty(\mathbb{R})$, $D : V \rightarrow V$, $Df = f'$.

A function $f \in C^\infty(\mathbb{R})$ is an eigenfunction of the operator D belonging to an eigenvalue λ if $f'(x) = \lambda f(x)$ for all $x \in \mathbb{R}$.

It follows that $f(x) = ce^{\lambda x}$, where c is a nonzero constant.

Thus each $\lambda \in \mathbb{R}$ is an eigenvalue of D .

The corresponding eigenspace is spanned by $e^{\lambda x}$.

Example. $V = C^\infty(\mathbb{R})$, $L : V \rightarrow V$, $Lf = f''$.

$$Lf = \lambda f \iff f''(x) - \lambda f(x) = 0 \text{ for all } x \in \mathbb{R}.$$

It follows that each $\lambda \in \mathbb{R}$ is an eigenvalue of L and the corresponding eigenspace V_λ is two-dimensional.

If $\lambda > 0$ then $V_\lambda = \text{Span}(\exp(\sqrt{\lambda}x), \exp(-\sqrt{\lambda}x))$.

If $\lambda < 0$ then $V_\lambda = \text{Span}(\sin(\sqrt{-\lambda}x), \cos(\sqrt{-\lambda}x))$.

If $\lambda = 0$ then $V_\lambda = \text{Span}(1, x)$.

Let V be a vector space and $L : V \rightarrow V$ be a linear operator.

Proposition 1 If $\mathbf{v} \in V$ is an eigenvector of the operator L then the associated eigenvalue is unique.

Proof: Suppose that $L(\mathbf{v}) = \lambda_1\mathbf{v}$ and $L(\mathbf{v}) = \lambda_2\mathbf{v}$. Then $\lambda_1\mathbf{v} = \lambda_2\mathbf{v} \implies (\lambda_1 - \lambda_2)\mathbf{v} = \mathbf{0} \implies \lambda_1 - \lambda_2 = 0 \implies \lambda_1 = \lambda_2$.

Proposition 2 Suppose \mathbf{v}_1 and \mathbf{v}_2 are eigenvectors of L associated with different eigenvalues λ_1 and λ_2 . Then \mathbf{v}_1 and \mathbf{v}_2 are linearly independent.

Proof: For any scalar $t \neq 0$ the vector $t\mathbf{v}_1$ is also an eigenvector of L associated with the eigenvalue λ_1 . Since $\lambda_2 \neq \lambda_1$, it follows that $\mathbf{v}_2 \neq t\mathbf{v}_1$. That is, \mathbf{v}_2 is not a scalar multiple of \mathbf{v}_1 . Similarly, \mathbf{v}_1 is not a scalar multiple of \mathbf{v}_2 .

Let $L : V \rightarrow V$ be a linear operator.

Proposition 3 If \mathbf{v}_1 , \mathbf{v}_2 , and \mathbf{v}_3 are eigenvectors of L associated with distinct eigenvalues λ_1 , λ_2 , and λ_3 , then they are linearly independent.

Proof: Suppose that $t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + t_3\mathbf{v}_3 = \mathbf{0}$ for some $t_1, t_2, t_3 \in \mathbb{R}$. Then

$$\begin{aligned}L(t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + t_3\mathbf{v}_3) &= \mathbf{0}, \\t_1L(\mathbf{v}_1) + t_2L(\mathbf{v}_2) + t_3L(\mathbf{v}_3) &= \mathbf{0}, \\t_1\lambda_1\mathbf{v}_1 + t_2\lambda_2\mathbf{v}_2 + t_3\lambda_3\mathbf{v}_3 &= \mathbf{0}.\end{aligned}$$

It follows that

$$\begin{aligned}t_1\lambda_1\mathbf{v}_1 + t_2\lambda_2\mathbf{v}_2 + t_3\lambda_3\mathbf{v}_3 - \lambda_3(t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + t_3\mathbf{v}_3) &= \mathbf{0} \\ \implies t_1(\lambda_1 - \lambda_3)\mathbf{v}_1 + t_2(\lambda_2 - \lambda_3)\mathbf{v}_2 &= \mathbf{0}.\end{aligned}$$

By the above, \mathbf{v}_1 and \mathbf{v}_2 are linearly independent.

Hence $t_1(\lambda_1 - \lambda_3) = t_2(\lambda_2 - \lambda_3) = 0$
 $\implies t_1 = t_2 = 0 \implies t_3 = 0$.

Theorem If $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are eigenvectors of a linear operator L associated with distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_k$, then $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly independent.

Corollary Let A be an $n \times n$ matrix such that the characteristic equation $\det(A - \lambda I) = 0$ has n distinct real roots. Then \mathbb{R}^n has a basis consisting of eigenvectors of A .

Proof: Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be distinct real roots of the characteristic equation. Any λ_i is an eigenvalue of A , hence there is an associated eigenvector \mathbf{v}_i . By the theorem, vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are linearly independent. Therefore they form a basis for \mathbb{R}^n .

Theorem If $\lambda_1, \lambda_2, \dots, \lambda_k$ are distinct real numbers, then the functions $e^{\lambda_1 x}, e^{\lambda_2 x}, \dots, e^{\lambda_k x}$ are linearly independent.

Proof: Consider a linear operator

$D : C^\infty(\mathbb{R}) \rightarrow C^\infty(\mathbb{R})$ given by $Df = f'$.

Then $e^{\lambda_1 x}, \dots, e^{\lambda_k x}$ are eigenfunctions of D associated with distinct eigenvalues $\lambda_1, \dots, \lambda_k$.