

MATH 304
Linear Algebra

Lecture 13:
Review for Test 1.

Topics for Test 1

Part I: Elementary linear algebra (Leon 1.1–1.5, 2.1–2.2)

- Systems of linear equations: elementary operations, Gaussian elimination, back substitution.
- Matrix of coefficients and augmented matrix. Elementary row operations, row echelon form and reduced row echelon form.
- Matrix algebra. Inverse matrix.
- Determinants: explicit formulas for 2×2 and 3×3 matrices, row and column expansions, elementary row and column operations.

Topics for Test 1

Part II: Abstract linear algebra (Leon 3.1–3.4, 3.6)

- Vector spaces (vectors, matrices, polynomials, functional spaces).
- Subspaces. Nullspace, column space, and row space of a matrix.
- Span, spanning set. Linear independence.
- Basis and dimension.
- Rank and nullity of a matrix.

Sample problems for Test 1

Problem 1 (15 pts.) Find a quadratic polynomial $p(x)$ such that $p(1) = 1$, $p(2) = 3$, and $p(3) = 7$.

Problem 2 (25 pts.) Let $A = \begin{pmatrix} 1 & -2 & 4 & 1 \\ 2 & 3 & 2 & 0 \\ 2 & 0 & -1 & 1 \\ 2 & 0 & 0 & 1 \end{pmatrix}$.

- (i) Evaluate the determinant of the matrix A .
- (ii) Find the inverse matrix A^{-1} .

Problem 3 (20 pts.) Determine which of the following subsets of \mathbb{R}^3 are subspaces. Briefly explain.

- (i) The set S_1 of vectors $(x, y, z) \in \mathbb{R}^3$ such that $xyz = 0$.
- (ii) The set S_2 of vectors $(x, y, z) \in \mathbb{R}^3$ such that $x + y + z = 0$.
- (iii) The set S_3 of vectors $(x, y, z) \in \mathbb{R}^3$ such that $y^2 + z^2 = 0$.
- (iv) The set S_4 of vectors $(x, y, z) \in \mathbb{R}^3$ such that $y^2 - z^2 = 0$.

Problem 4 (30 pts.) Let $B = \begin{pmatrix} 0 & -1 & 4 & 1 \\ 1 & 1 & 2 & -1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix}$.

- (i) Find the rank and the nullity of the matrix B .
- (ii) Find a basis for the row space of B , then extend this basis to a basis for \mathbb{R}^4 .
- (iii) Find a basis for the nullspace of B .

Bonus Problem 5 (15 pts.) Show that the functions $f_1(x) = x$, $f_2(x) = xe^x$, and $f_3(x) = e^{-x}$ are linearly independent in the vector space $C^\infty(\mathbb{R})$.

Bonus Problem 6 (15 pts.) Let V be a finite-dimensional vector space and V_0 be a proper subspace of V (where proper means that $V_0 \neq V$). Prove that $\dim V_0 < \dim V$.

Problem 1. Find a quadratic polynomial $p(x)$ such that $p(1) = 1$, $p(2) = 3$, and $p(3) = 7$.

Let $p(x) = ax^2 + bx + c$. Then $p(1) = a + b + c$, $p(2) = 4a + 2b + c$, and $p(3) = 9a + 3b + c$.

The coefficients a , b , and c have to be chosen so that

$$\begin{cases} a + b + c = 1, \\ 4a + 2b + c = 3, \\ 9a + 3b + c = 7. \end{cases}$$

We solve this system of linear equations using elementary operations:

$$\begin{cases} a + b + c = 1 \\ 4a + 2b + c = 3 \\ 9a + 3b + c = 7 \end{cases} \iff \begin{cases} a + b + c = 1 \\ 3a + b = 2 \\ 9a + 3b + c = 7 \end{cases}$$

$$\Leftrightarrow \begin{cases} a + b + c = 1 \\ 3a + b = 2 \\ 9a + 3b + c = 7 \end{cases} \Leftrightarrow \begin{cases} a + b + c = 1 \\ 3a + b = 2 \\ 8a + 2b = 6 \end{cases}$$

$$\Leftrightarrow \begin{cases} a + b + c = 1 \\ 3a + b = 2 \\ 4a + b = 3 \end{cases} \Leftrightarrow \begin{cases} a + b + c = 1 \\ 3a + b = 2 \\ a = 1 \end{cases}$$

$$\Leftrightarrow \begin{cases} a + b + c = 1 \\ b = -1 \\ a = 1 \end{cases} \Leftrightarrow \begin{cases} c = 1 \\ b = -1 \\ a = 1 \end{cases}$$

Thus the desired polynomial is $p(x) = x^2 - x + 1$.

Problem 2. Let $A = \begin{pmatrix} 1 & -2 & 4 & 1 \\ 2 & 3 & 2 & 0 \\ 2 & 0 & -1 & 1 \\ 2 & 0 & 0 & 1 \end{pmatrix}$.

(i) Evaluate the determinant of the matrix A .

Subtract the 4th row of A from the 3rd row:

$$\begin{vmatrix} 1 & -2 & 4 & 1 \\ 2 & 3 & 2 & 0 \\ 2 & 0 & -1 & 1 \\ 2 & 0 & 0 & 1 \end{vmatrix} = \begin{vmatrix} 1 & -2 & 4 & 1 \\ 2 & 3 & 2 & 0 \\ 0 & 0 & -1 & 0 \\ 2 & 0 & 0 & 1 \end{vmatrix}.$$

Expand the determinant by the 3rd row:

$$\begin{vmatrix} 1 & -2 & 4 & 1 \\ 2 & 3 & 2 & 0 \\ 0 & 0 & -1 & 0 \\ 2 & 0 & 0 & 1 \end{vmatrix} = (-1) \begin{vmatrix} 1 & -2 & 1 \\ 2 & 3 & 0 \\ 2 & 0 & 1 \end{vmatrix}.$$

Expand the determinant by the 3rd column:

$$(-1) \begin{vmatrix} 1 & -2 & 1 \\ 2 & 3 & 0 \\ 2 & 0 & 1 \end{vmatrix} = (-1) \left(\begin{vmatrix} 2 & 3 \\ 2 & 0 \end{vmatrix} + \begin{vmatrix} 1 & -2 \\ 2 & 3 \end{vmatrix} \right) = -1.$$

Problem 2. Let $A = \begin{pmatrix} 1 & -2 & 4 & 1 \\ 2 & 3 & 2 & 0 \\ 2 & 0 & -1 & 1 \\ 2 & 0 & 0 & 1 \end{pmatrix}$.

(ii) Find the inverse matrix A^{-1} .

First we merge the matrix A with the identity matrix into one 4×8 matrix

$$(A|I) = \left(\begin{array}{cccc|cccc} 1 & -2 & 4 & 1 & 1 & 0 & 0 & 0 \\ 2 & 3 & 2 & 0 & 0 & 1 & 0 & 0 \\ 2 & 0 & -1 & 1 & 0 & 0 & 1 & 0 \\ 2 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right).$$

Then we apply elementary row operations to this matrix until the left part becomes the identity matrix.

Subtract 2 times the 1st row from the 2nd row:

$$\left(\begin{array}{cccc|cccc} 1 & -2 & 4 & 1 & 1 & 0 & 0 & 0 \\ 0 & 7 & -6 & -2 & -2 & 1 & 0 & 0 \\ 2 & 0 & -1 & 1 & 0 & 0 & 1 & 0 \\ 2 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right)$$

Subtract 2 times the 1st row from the 3rd row:

$$\left(\begin{array}{cccc|cccc} 1 & -2 & 4 & 1 & 1 & 0 & 0 & 0 \\ 0 & 7 & -6 & -2 & -2 & 1 & 0 & 0 \\ 0 & 4 & -9 & -1 & -2 & 0 & 1 & 0 \\ 2 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right)$$

Subtract 2 times the 1st row from the 4th row:

$$\left(\begin{array}{cccc|cccc} 1 & -2 & 4 & 1 & 1 & 0 & 0 & 0 \\ 0 & 7 & -6 & -2 & -2 & 1 & 0 & 0 \\ 0 & 4 & -9 & -1 & -2 & 0 & 1 & 0 \\ 0 & 4 & -8 & -1 & -2 & 0 & 0 & 1 \end{array} \right)$$

Subtract 2 times the 4th row from the 2nd row:

$$\left(\begin{array}{cccc|cccc} 1 & -2 & 4 & 1 & 1 & 0 & 0 & 0 \\ 0 & -1 & 10 & 0 & 2 & 1 & 0 & -2 \\ 0 & 4 & -9 & -1 & -2 & 0 & 1 & 0 \\ 0 & 4 & -8 & -1 & -2 & 0 & 0 & 1 \end{array} \right)$$

Subtract the 4th row from the 3rd row:

$$\left(\begin{array}{cccc|cccc} 1 & -2 & 4 & 1 & 1 & 0 & 0 & 0 \\ 0 & -1 & 10 & 0 & 2 & 1 & 0 & -2 \\ 0 & 0 & -1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 4 & -8 & -1 & -2 & 0 & 0 & 1 \end{array} \right)$$

Add 4 times the 2nd row to the 4th row:

$$\left(\begin{array}{cccc|cccc} 1 & -2 & 4 & 1 & 1 & 0 & 0 & 0 \\ 0 & -1 & 10 & 0 & 2 & 1 & 0 & -2 \\ 0 & 0 & -1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 32 & -1 & 6 & 4 & 0 & -7 \end{array} \right)$$

Add 32 times the 3rd row to the 4th row:

$$\left(\begin{array}{cccc|cccc} 1 & -2 & 4 & 1 & 1 & 0 & 0 & 0 \\ 0 & -1 & 10 & 0 & 2 & 1 & 0 & -2 \\ 0 & 0 & -1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & -1 & 6 & 4 & 32 & -39 \end{array} \right)$$

Add 10 times the 3rd row to the 2nd row:

$$\left(\begin{array}{cccc|cccc} 1 & -2 & 4 & 1 & 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 2 & 1 & 10 & -12 \\ 0 & 0 & -1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & -1 & 6 & 4 & 32 & -39 \end{array} \right)$$

Add the 4th row to the 1st row:

$$\left(\begin{array}{cccc|cccc} 1 & -2 & 4 & 0 & 7 & 4 & 32 & -39 \\ 0 & -1 & 0 & 0 & 2 & 1 & 10 & -12 \\ 0 & 0 & -1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & -1 & 6 & 4 & 32 & -39 \end{array} \right)$$

Add 4 times the 3rd row to the 1st row:

$$\left(\begin{array}{cccc|cccc} 1 & -2 & 0 & 0 & 7 & 4 & 36 & -43 \\ 0 & -1 & 0 & 0 & 2 & 1 & 10 & -12 \\ 0 & 0 & -1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & -1 & 6 & 4 & 32 & -39 \end{array} \right)$$

Subtract 2 times the 2nd row from the 1st row:

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 3 & 2 & 16 & -19 \\ 0 & -1 & 0 & 0 & 2 & 1 & 10 & -12 \\ 0 & 0 & -1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & -1 & 6 & 4 & 32 & -39 \end{array} \right)$$

Multiply the 2nd, the 3rd, and the 4th rows by -1 :

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 3 & 2 & 16 & -19 \\ 0 & 1 & 0 & 0 & -2 & -1 & -10 & 12 \\ 0 & 0 & 1 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 1 & -6 & -4 & -32 & 39 \end{array} \right)$$

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 3 & 2 & 16 & -19 \\ 0 & 1 & 0 & 0 & -2 & -1 & -10 & 12 \\ 0 & 0 & 1 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 1 & -6 & -4 & -32 & 39 \end{array} \right) = (I | A^{-1})$$

Finally the left part of our 4×8 matrix is transformed into the identity matrix. Therefore the current right part is the inverse matrix of A . Thus

$$A^{-1} = \begin{pmatrix} 1 & -2 & 4 & 1 \\ 2 & 3 & 2 & 0 \\ 2 & 0 & -1 & 1 \\ 2 & 0 & 0 & 1 \end{pmatrix}^{-1} = \begin{pmatrix} 3 & 2 & 16 & -19 \\ -2 & -1 & -10 & 12 \\ 0 & 0 & -1 & 1 \\ -6 & -4 & -32 & 39 \end{pmatrix}.$$

Problem 2. Let $A = \begin{pmatrix} 1 & -2 & 4 & 1 \\ 2 & 3 & 2 & 0 \\ 2 & 0 & -1 & 1 \\ 2 & 0 & 0 & 1 \end{pmatrix}$.

(i) Evaluate the determinant of the matrix A .

Alternative solution: We have transformed A into the identity matrix using elementary row operations. These included no row exchanges and three row multiplications, each time by -1 .

It follows that $\det I = (-1)^3 \det A$.

$$\implies \det A = -\det I = -1.$$

Problem 3. Determine which of the following subsets of \mathbb{R}^3 are subspaces. Briefly explain.

A subset of \mathbb{R}^3 is a subspace if it is closed under addition and scalar multiplication. Besides, the subset must not be empty.

(i) The set S_1 of vectors $(x, y, z) \in \mathbb{R}^3$ such that $xyz = 0$.

$(0, 0, 0) \in S_1 \implies S_1$ is not empty.

$xyz = 0 \implies (rx)(ry)(rz) = r^3xyz = 0$.

That is, $\mathbf{v} = (x, y, z) \in S_1 \implies r\mathbf{v} = (rx, ry, rz) \in S_1$.

Hence S_1 is closed under scalar multiplication.

However S_1 is not closed under addition.

Counterexample: $(1, 1, 0) + (0, 0, 1) = (1, 1, 1)$.

Problem 3. Determine which of the following subsets of \mathbb{R}^3 are subspaces. Briefly explain.

A subset of \mathbb{R}^3 is a subspace if it is closed under addition and scalar multiplication. Besides, the subset must not be empty.

(ii) The set S_2 of vectors $(x, y, z) \in \mathbb{R}^3$ such that $x + y + z = 0$.

$(0, 0, 0) \in S_2 \implies S_2$ is not empty.

$x + y + z = 0 \implies rx + ry + rz = r(x + y + z) = 0$.

Hence S_2 is closed under scalar multiplication.

$x + y + z = x' + y' + z' = 0 \implies$

$(x + x') + (y + y') + (z + z') = (x + y + z) + (x' + y' + z') = 0$.

That is, $\mathbf{v} = (x, y, z)$, $\mathbf{v}' = (x', y', z') \in S_2$

$\implies \mathbf{v} + \mathbf{v}' = (x + x', y + y', z + z') \in S_2$.

Hence S_2 is closed under addition.

(iii) The set S_3 of vectors $(x, y, z) \in \mathbb{R}^3$ such that $y^2 + z^2 = 0$.

$$y^2 + z^2 = 0 \iff y = z = 0.$$

S_3 is a nonempty set closed under addition and scalar multiplication.

(iv) The set S_4 of vectors $(x, y, z) \in \mathbb{R}^3$ such that $y^2 - z^2 = 0$.

S_4 is a nonempty set closed under scalar multiplication. However S_4 is not closed under addition.

Counterexample: $(0, 1, 1) + (0, 1, -1) = (0, 2, 0)$.

Problem 4. Let $B = \begin{pmatrix} 0 & -1 & 4 & 1 \\ 1 & 1 & 2 & -1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix}$.

(i) Find the rank and the nullity of the matrix B .

The rank (= dimension of the row space) and the nullity (= dimension of the nullspace) of a matrix are preserved under elementary row operations. We apply such operations to convert the matrix B into row echelon form.

Interchange the 1st row with the 2nd row:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & -1 & 4 & 1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix}$$

Add 3 times the 1st row to the 3rd row, then subtract 2 times the 1st row from the 4th row:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & -1 & 4 & 1 \\ 0 & 3 & 5 & -3 \\ 2 & -1 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & -1 & 4 & 1 \\ 0 & 3 & 5 & -3 \\ 0 & -3 & -4 & 3 \end{pmatrix}$$

Multiply the 2nd row by -1 :

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 3 & 5 & -3 \\ 0 & -3 & -4 & 3 \end{pmatrix}$$

Add the 4th row to the 3rd row:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -3 & -4 & 3 \end{pmatrix}$$

Add 3 times the 2nd row to the 4th row:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -16 & 0 \end{pmatrix}$$

Add 16 times the 3rd row to the 4th row:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Now that the matrix is in row echelon form, its rank equals the number of nonzero rows, which is 3. Since

$(\text{rank of } B) + (\text{nullity of } B) = (\text{the number of columns of } B) = 4$,
it follows that the nullity of B equals 1.

Problem 4. Let $B = \begin{pmatrix} 0 & -1 & 4 & 1 \\ 1 & 1 & 2 & -1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix}$.

(ii) Find a basis for the row space of B , then extend this basis to a basis for \mathbb{R}^4 .

The row space of a matrix is invariant under elementary row operations. Therefore the row space of the matrix B is the same as the row space of its row echelon form:

$$\begin{pmatrix} 0 & -1 & 4 & 1 \\ 1 & 1 & 2 & -1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

The nonzero rows of the latter matrix are linearly independent so that they form a basis for its row space:

$$\mathbf{v}_1 = (1, 1, 2, -1), \quad \mathbf{v}_2 = (0, 1, -4, -1), \quad \mathbf{v}_3 = (0, 0, 1, 0).$$

To extend the basis $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ to a basis for \mathbb{R}^4 , we need a vector $\mathbf{v}_4 \in \mathbb{R}^4$ that is not a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$.

It is known that at least one of the vectors $\mathbf{e}_1 = (1, 0, 0, 0)$, $\mathbf{e}_2 = (0, 1, 0, 0)$, $\mathbf{e}_3 = (0, 0, 1, 0)$, and $\mathbf{e}_4 = (0, 0, 0, 1)$ can be chosen as \mathbf{v}_4 .

In particular, the vectors $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{e}_4$ form a basis for \mathbb{R}^4 .

This follows from the fact that the 4×4 matrix whose rows are these vectors is not singular:

$$\begin{vmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{vmatrix} = 1 \neq 0.$$

Problem 4. Let $B = \begin{pmatrix} 0 & -1 & 4 & 1 \\ 1 & 1 & 2 & -1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix}$.

(iii) Find a basis for the nullspace of B .

The nullspace of B is the solution set of the system of linear homogeneous equations with B as the coefficient matrix. To solve the system, we convert B to reduced row echelon form:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\implies x_1 = x_2 - x_4 = x_3 = 0$$

General solution: $(x_1, x_2, x_3, x_4) = (0, t, 0, t) = t(0, 1, 0, 1)$.

Thus the vector $(0, 1, 0, 1)$ forms a basis for the nullspace of B .

Bonus Problem 5. Show that the functions $f_1(x) = x$, $f_2(x) = xe^x$, and $f_3(x) = e^{-x}$ are linearly independent in the vector space $C^\infty(\mathbb{R})$.

The functions f_1, f_2, f_3 are linearly independent whenever the Wronskian $W[f_1, f_2, f_3]$ is not identically zero.

$$\begin{aligned} W[f_1, f_2, f_3](x) &= \begin{vmatrix} f_1(x) & f_2(x) & f_3(x) \\ f_1'(x) & f_2'(x) & f_3'(x) \\ f_1''(x) & f_2''(x) & f_3''(x) \end{vmatrix} = \begin{vmatrix} x & xe^x & e^{-x} \\ 1 & e^x + xe^x & -e^{-x} \\ 0 & 2e^x + xe^x & e^{-x} \end{vmatrix} \\ &= e^{-x} \begin{vmatrix} x & xe^x & 1 \\ 1 & e^x + xe^x & -1 \\ 0 & 2e^x + xe^x & 1 \end{vmatrix} = \begin{vmatrix} x & x & 1 \\ 1 & 1+x & -1 \\ 0 & 2+x & 1 \end{vmatrix} \\ &= x \begin{vmatrix} 1+x & -1 \\ 2+x & 1 \end{vmatrix} - \begin{vmatrix} x & 1 \\ 2+x & 1 \end{vmatrix} = x(2x+3) + 2 = 2x^2 + 3x + 2. \end{aligned}$$

The polynomial $2x^2 + 3x + 2$ is never zero.

Bonus Problem 5. Show that the functions $f_1(x) = x$, $f_2(x) = xe^x$, and $f_3(x) = e^{-x}$ are linearly independent in the vector space $C^\infty(\mathbb{R})$.

Alternative solution: Suppose that $af_1(x) + bf_2(x) + cf_3(x) = 0$ for all $x \in \mathbb{R}$, where a, b, c are constants. We have to show that $a = b = c = 0$.

Let us differentiate this identity:

$$ax + bxe^x + ce^{-x} = 0,$$

$$a + be^x + bxe^x - ce^{-x} = 0,$$

$$2be^x + bxe^x + ce^{-x} = 0,$$

$$3be^x + bxe^x - ce^{-x} = 0,$$

$$4be^x + bxe^x + ce^{-x} = 0.$$

(the 5th identity) – (the 3rd identity): $2be^x = 0 \implies b = 0$.

Substitute $b = 0$ in the 3rd identity: $ce^{-x} = 0 \implies c = 0$.

Substitute $b = c = 0$ in the 2nd identity: $a = 0$.

Bonus Problem 5. Show that the functions $f_1(x) = x$, $f_2(x) = xe^x$, and $f_3(x) = e^{-x}$ are linearly independent in the vector space $C^\infty(\mathbb{R})$.

Alternative solution: Suppose that $ax + bxe^x + ce^{-x} = 0$ for all $x \in \mathbb{R}$, where a, b, c are constants. We have to show that $a = b = c = 0$.

For any $x \neq 0$ divide both sides of the identity by xe^x :

$$ae^{-x} + b + cx^{-1}e^{-2x} = 0.$$

The left-hand side approaches b as $x \rightarrow +\infty$. $\implies b = 0$

Now $ax + ce^{-x} = 0$ for all $x \in \mathbb{R}$. For any $x \neq 0$ divide both sides of the identity by x :

$$a + cx^{-1}e^{-x} = 0.$$

The left-hand side approaches a as $x \rightarrow +\infty$. $\implies a = 0$

Now $ce^{-x} = 0 \implies c = 0$.

Bonus Problem 6. Let V be a finite-dimensional vector space and V_0 be a proper subspace of V (where proper means that $V_0 \neq V$). Prove that $\dim V_0 < \dim V$.

Any vector space has a basis. Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ be a basis for V_0 .

Vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly independent in V since they are linearly independent in V_0 . Therefore we can extend this collection of vectors to a basis for V by adding some vectors $\mathbf{w}_1, \dots, \mathbf{w}_m$. As $V_0 \neq V$, we do need to add some vectors, i.e., $m \geq 1$.

Thus $\dim V_0 = k$ and $\dim V = k + m > k$.