

MATH 304

Linear Algebra

Lecture 17:

Basis and dimension (continued).

Rank of a matrix.

Basis

Definition. Let V be a vector space. A linearly independent spanning set for V is called a **basis**.

Equivalently, a subset $S \subset V$ is a basis for V if any vector $\mathbf{v} \in V$ is *uniquely represented* as a linear combination

$$\mathbf{v} = r_1\mathbf{v}_1 + r_2\mathbf{v}_2 + \cdots + r_k\mathbf{v}_k,$$

where $\mathbf{v}_1, \dots, \mathbf{v}_k$ are distinct vectors from S and $r_1, \dots, r_k \in \mathbb{R}$.

Dimension

Theorem 1 Any vector space has a basis.

Theorem 2 If a vector space V has a finite basis, then all bases for V are finite and have the same number of elements.

Definition. The **dimension** of a vector space V , denoted $\dim V$, is the number of elements in any of its bases.

Examples. • $\dim \mathbb{R}^n = n$

- $\mathcal{M}_{m,n}(\mathbb{R})$: the space of $m \times n$ matrices; $\dim \mathcal{M}_{m,n} = mn$
- \mathcal{P}_n : polynomials of degree less than n ; $\dim \mathcal{P}_n = n$
- \mathcal{P} : the space of all polynomials; $\dim \mathcal{P} = \infty$
- $\{\mathbf{0}\}$: the trivial vector space; $\dim \{\mathbf{0}\} = 0$

How to find a basis?

Theorem Let V be a vector space. Then

- (i) any spanning set for V contains a basis;
- (ii) any linearly independent subset of V is contained in a basis.

Approach 1. Get a spanning set for the vector space, then reduce this set to a basis.

Approach 2. Get a linearly independent set, then extend it to a basis.

Problem. Find a basis for the vector space V spanned by vectors $\mathbf{w}_1 = (1, 1, 0)$, $\mathbf{w}_2 = (0, 1, 1)$, $\mathbf{w}_3 = (2, 3, 1)$, and $\mathbf{w}_4 = (1, 1, 1)$.

To pare this spanning set, we need to find a relation of the form $r_1\mathbf{w}_1 + r_2\mathbf{w}_2 + r_3\mathbf{w}_3 + r_4\mathbf{w}_4 = \mathbf{0}$, where $r_i \in \mathbb{R}$ are not all equal to zero. Equivalently,

$$\begin{pmatrix} 1 & 0 & 2 & 1 \\ 1 & 1 & 3 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} r_1 \\ r_2 \\ r_3 \\ r_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

To solve this system of linear equations for r_1, r_2, r_3, r_4 , we apply row reduction.

$$\begin{pmatrix} 1 & 0 & 2 & 1 \\ 1 & 1 & 3 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\
 \rightarrow \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (\text{reduced row echelon form})$$

$$\begin{cases} r_1 + 2r_3 = 0 \\ r_2 + r_3 = 0 \\ r_4 = 0 \end{cases} \iff \begin{cases} r_1 = -2r_3 \\ r_2 = -r_3 \\ r_4 = 0 \end{cases}$$

General solution: $(r_1, r_2, r_3, r_4) = (-2t, -t, t, 0)$, $t \in \mathbb{R}$.

Particular solution: $(r_1, r_2, r_3, r_4) = (2, 1, -1, 0)$.

Problem. Find a basis for the vector space V spanned by vectors $\mathbf{w}_1 = (1, 1, 0)$, $\mathbf{w}_2 = (0, 1, 1)$, $\mathbf{w}_3 = (2, 3, 1)$, and $\mathbf{w}_4 = (1, 1, 1)$.

We have obtained that $2\mathbf{w}_1 + \mathbf{w}_2 - \mathbf{w}_3 = \mathbf{0}$.

Hence any of vectors $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3$ can be dropped.

For instance, $V = \text{Span}(\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_4)$.

Let us check whether vectors $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_4$ are linearly independent:

$$\begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 1 & 0 \end{vmatrix} = \begin{vmatrix} 1 & 1 \\ 0 & 1 \end{vmatrix} = 1 \neq 0.$$

They are!!! It follows that $V = \mathbb{R}^3$ and $\{\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_4\}$ is a basis for V .

Row space of a matrix

Definition. The **row space** of an $m \times n$ matrix A is the subspace of \mathbb{R}^n spanned by rows of A .

The dimension of the row space is called the **rank** of the matrix A .

Theorem 1 The rank of a matrix A is the maximal number of linearly independent rows in A .

Theorem 2 Elementary row operations do not change the row space of a matrix.

Theorem 3 If a matrix A is in row echelon form, then the nonzero rows of A are linearly independent.

Corollary The rank of a matrix is equal to the number of nonzero rows in its row echelon form.

Theorem Elementary row operations do not change the row space of a matrix.

Proof: Suppose that A and B are $m \times n$ matrices such that B is obtained from A by an elementary row operation. Let $\mathbf{a}_1, \dots, \mathbf{a}_m$ be the rows of A and $\mathbf{b}_1, \dots, \mathbf{b}_m$ be the rows of B . We have to show that $\text{Span}(\mathbf{a}_1, \dots, \mathbf{a}_m) = \text{Span}(\mathbf{b}_1, \dots, \mathbf{b}_m)$.

Observe that any row \mathbf{b}_i of B belongs to $\text{Span}(\mathbf{a}_1, \dots, \mathbf{a}_m)$. Indeed, either $\mathbf{b}_i = \mathbf{a}_j$ for some $1 \leq j \leq m$, or $\mathbf{b}_i = r\mathbf{a}_i$ for some scalar $r \neq 0$, or $\mathbf{b}_i = \mathbf{a}_i + r\mathbf{a}_j$ for some $j \neq i$ and $r \in \mathbb{R}$.

It follows that $\text{Span}(\mathbf{b}_1, \dots, \mathbf{b}_m) \subset \text{Span}(\mathbf{a}_1, \dots, \mathbf{a}_m)$.

Now the matrix A can also be obtained from B by an elementary row operation. By the above,

$$\text{Span}(\mathbf{a}_1, \dots, \mathbf{a}_m) \subset \text{Span}(\mathbf{b}_1, \dots, \mathbf{b}_m).$$

Problem. Find the rank of the matrix

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 1 \end{pmatrix}.$$

Elementary row operations do not change the row space. Let us convert A to row echelon form:

$$\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

Vectors $(1, 1, 0)$, $(0, 1, 1)$, and $(0, 0, 1)$ form a basis for the row space of A . Thus the rank of A is 3.

It follows that the row space of A is the entire space \mathbb{R}^3 .

Problem. Find a basis for the vector space V spanned by vectors $\mathbf{w}_1 = (1, 1, 0)$, $\mathbf{w}_2 = (0, 1, 1)$, $\mathbf{w}_3 = (2, 3, 1)$, and $\mathbf{w}_4 = (1, 1, 1)$.

The vector space V is the row space of a matrix

$$\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 1 \end{pmatrix}.$$

According to the solution of the previous problem, vectors $(1, 1, 0)$, $(0, 1, 1)$, and $(0, 0, 1)$ form a basis for V .

Column space of a matrix

Definition. The **column space** of an $m \times n$ matrix A is the subspace of \mathbb{R}^m spanned by columns of A .

Theorem 1 The column space of a matrix A coincides with the row space of the transpose matrix A^T .

Theorem 2 Elementary column operations do not change the column space of a matrix.

Theorem 3 Elementary row operations do not change the dimension of the column space of a matrix (although they can change the column space).

Theorem 4 For any matrix, the row space and the column space have the same dimension.

Problem. Find a basis for the column space of the matrix

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 1 \end{pmatrix}.$$

The column space of A coincides with the row space of A^T . To find a basis, we convert A^T to row echelon form:

$$A^T = \begin{pmatrix} 1 & 0 & 2 & 1 \\ 1 & 1 & 3 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} \mathbf{1} & 0 & 2 & 1 \\ 0 & \mathbf{1} & 1 & 0 \\ 0 & 0 & 0 & \mathbf{1} \end{pmatrix}$$

Vectors $(1, 0, 2, 1)$, $(0, 1, 1, 0)$, and $(0, 0, 0, 1)$ form a basis for the column space of A .

Problem. Find a basis for the column space of the matrix

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 1 \end{pmatrix}.$$

Alternative solution: We already know from a previous problem that the rank of A is 3. It follows that the columns of A are linearly independent. Therefore these columns form a basis for the column space.