

MATH 304

Linear Algebra

Lecture 18:

Rank and nullity of a matrix.

Basis and coordinates.

Change of coordinates.

Rank of a matrix

Definition. The **row space** of an $m \times n$ matrix A is the subspace of \mathbb{R}^n spanned by rows of A . The **column space** of A is a subspace of \mathbb{R}^m spanned by columns of A .

The row space and the column space of A have the same dimension, which is called the **rank** of A .

Theorem 1 Elementary row operations do not change the row space of a matrix.

Theorem 2 If a matrix A is in row echelon form, then the nonzero rows of A form a basis for the row space.

Theorem 3 The rank of a matrix is equal to the number of nonzero rows in its row echelon form.

Nullspace of a matrix

Let $A = (a_{ij})$ be an $m \times n$ matrix.

Definition. The **nullspace** of the matrix A , denoted $N(A)$, is the set of all n -dimensional column vectors \mathbf{x} such that $\boxed{A\mathbf{x} = \mathbf{0}}$.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

The nullspace $N(A)$ is the solution set of a system of linear homogeneous equations (with A as the coefficient matrix).

Let A be an $m \times n$ matrix. Then the nullspace $N(A)$ is the solution set of a system of linear homogeneous equations in n variables.

Theorem The nullspace $N(A)$ is a subspace of the vector space \mathbb{R}^n .

Definition. The dimension of the nullspace $N(A)$ is called the **nullity** of the matrix A .

Problem. Find the nullity of the matrix

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 2 & 3 & 4 & 5 \end{pmatrix}.$$

Elementary row operations do not change the nullspace.

Let us convert A to reduced row echelon form:

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 2 & 3 & 4 & 5 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 3 \end{pmatrix}$$

$$\begin{cases} x_1 - x_3 - 2x_4 = 0 \\ x_2 + 2x_3 + 3x_4 = 0 \end{cases} \iff \begin{cases} x_1 = x_3 + 2x_4 \\ x_2 = -2x_3 - 3x_4 \end{cases}$$

General element of $N(A)$:

$$\begin{aligned} (x_1, x_2, x_3, x_4) &= (t + 2s, -2t - 3s, t, s) \\ &= t(1, -2, 1, 0) + s(2, -3, 0, 1), \quad t, s \in \mathbb{R}. \end{aligned}$$

Vectors $(1, -2, 1, 0)$ and $(2, -3, 0, 1)$ form a basis for $N(A)$.

Thus the nullity of the matrix A is 2.

rank + nullity

Theorem The rank of a matrix A plus the nullity of A equals the number of columns in A .

Sketch of the proof: The rank of A equals the number of nonzero rows in the row echelon form, which equals the number of leading entries.

The nullity of A equals the number of free variables in the corresponding system, which equals the number of columns without leading entries in the row echelon form.

Consequently, rank+nullity is the number of all columns in the matrix A .

Problem. Find the nullity of the matrix

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 2 & 3 & 4 & 5 \end{pmatrix}.$$

Alternative solution: Clearly, the rows of A are linearly independent. Therefore the rank of A is 2. Since

$$(\text{rank of } A) + (\text{nullity of } A) = 4,$$

it follows that the nullity of A is 2.

Basis and dimension

Definition. Let V be a vector space. A linearly independent spanning set for V is called a **basis**.

Theorem Any vector space V has a basis. If V has a finite basis, then all bases for V are finite and have the same number of elements (called the *dimension* of V).

Example. Vectors $\mathbf{e}_1 = (1, 0, 0, \dots, 0, 0)$, $\mathbf{e}_2 = (0, 1, 0, \dots, 0, 0), \dots, \mathbf{e}_n = (0, 0, 0, \dots, 0, 1)$ form a basis for \mathbb{R}^n (called *standard*) since

$$(x_1, x_2, \dots, x_n) = x_1\mathbf{e}_1 + x_2\mathbf{e}_2 + \cdots + x_n\mathbf{e}_n.$$

Basis and coordinates

If $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a basis for a vector space V , then any vector $\mathbf{v} \in V$ has a unique representation

$$\mathbf{v} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \dots + x_n\mathbf{v}_n,$$

where $x_i \in \mathbb{R}$. The coefficients x_1, x_2, \dots, x_n are called the **coordinates** of \mathbf{v} with respect to the ordered basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$.

The mapping

$$\text{vector } \mathbf{v} \mapsto \text{its coordinates } (x_1, x_2, \dots, x_n)$$

is a one-to-one correspondence between V and \mathbb{R}^n . This correspondence respects linear operations in V and in \mathbb{R}^n .

Examples. • Coordinates of a vector

$\mathbf{v} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ relative to the standard basis $\mathbf{e}_1 = (1, 0, \dots, 0, 0)$, $\mathbf{e}_2 = (0, 1, \dots, 0, 0), \dots$, $\mathbf{e}_n = (0, 0, \dots, 0, 1)$ are (x_1, x_2, \dots, x_n) .

• Coordinates of a matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathcal{M}_{2,2}(\mathbb{R})$

relative to the basis $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ are (a, c, b, d) .

• Coordinates of a polynomial

$p(x) = a_0 + a_1x + \dots + a_{n-1}x^{n-1} \in \mathcal{P}_n$ relative to the basis $1, x, x^2, \dots, x^{n-1}$ are $(a_0, a_1, \dots, a_{n-1})$.

Vectors $\mathbf{u}_1=(3, 1)$ and $\mathbf{u}_2=(2, 1)$ form a basis for \mathbb{R}^2 .

Problem 1. Find coordinates of the vector $\mathbf{v} = (7, 4)$ with respect to the basis $\mathbf{u}_1, \mathbf{u}_2$.

The desired coordinates x, y satisfy

$$\mathbf{v} = x\mathbf{u}_1 + y\mathbf{u}_2 \iff \begin{cases} 3x + 2y = 7 \\ x + y = 4 \end{cases} \iff \begin{cases} x = -1 \\ y = 5 \end{cases}$$

Problem 2. Find the vector \mathbf{w} whose coordinates with respect to the basis $\mathbf{u}_1, \mathbf{u}_2$ are $(7, 4)$.

$$\mathbf{w} = 7\mathbf{u}_1 + 4\mathbf{u}_2 = 7(3, 1) + 4(2, 1) = (29, 11)$$

Change of coordinates

Given a vector $\mathbf{v} \in \mathbb{R}^2$, let (x, y) be its standard coordinates, i.e., coordinates with respect to the standard basis $\mathbf{e}_1 = (1, 0)$, $\mathbf{e}_2 = (0, 1)$, and let (x', y') be its coordinates with respect to the basis $\mathbf{u}_1 = (3, 1)$, $\mathbf{u}_2 = (2, 1)$.

Problem. Find a relation between (x, y) and (x', y') .

By definition, $\mathbf{v} = x\mathbf{e}_1 + y\mathbf{e}_2 = x'\mathbf{u}_1 + y'\mathbf{u}_2$.

In standard coordinates,

$$\begin{aligned} \begin{pmatrix} x \\ y \end{pmatrix} &= x' \begin{pmatrix} 3 \\ 1 \end{pmatrix} + y' \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x' \\ y' \end{pmatrix} \\ \implies \begin{pmatrix} x' \\ y' \end{pmatrix} &= \begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & -2 \\ -1 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \end{aligned}$$

Change of coordinates in \mathbb{R}^n

The usual (standard) coordinates of a vector

$\mathbf{v} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ are coordinates relative to the standard basis $\mathbf{e}_1 = (1, 0, \dots, 0, 0)$, $\mathbf{e}_2 = (0, 1, \dots, 0, 0), \dots$, $\mathbf{e}_n = (0, 0, \dots, 0, 1)$.

Let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ be another basis for \mathbb{R}^n and $(x'_1, x'_2, \dots, x'_n)$ be the coordinates of the same vector \mathbf{v} with respect to this basis.

Problem 1. Given the standard coordinates (x_1, x_2, \dots, x_n) , find the nonstandard coordinates $(x'_1, x'_2, \dots, x'_n)$.

Problem 2. Given the nonstandard coordinates $(x'_1, x'_2, \dots, x'_n)$, find the standard coordinates (x_1, x_2, \dots, x_n) .

It turns out that

$$\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ u_{21} & u_{22} & \dots & u_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ u_{n1} & u_{n2} & \dots & u_{nn} \end{pmatrix} \begin{pmatrix} x'_1 \\ x'_2 \\ \vdots \\ x'_n \end{pmatrix}.$$

The matrix $U = (u_{ij})$ does not depend on the vector \mathbf{v} .

Columns of U are coordinates of vectors

$\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ with respect to the standard basis.

U is called the **transition matrix** from the basis $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ to the standard basis $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$.

This solves Problem 2. To solve Problem 1, we have to use the inverse matrix U^{-1} , which is the transition matrix from $\mathbf{e}_1, \dots, \mathbf{e}_n$ to $\mathbf{u}_1, \dots, \mathbf{u}_n$.