

MATH 304

Linear Algebra

**Lecture 20:**

**Change of coordinates (continued).**

**Linear transformations.**

## Basis and coordinates

If  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  is a basis for a vector space  $V$ , then any vector  $\mathbf{v} \in V$  has a unique representation

$$\mathbf{v} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \dots + x_n\mathbf{v}_n,$$

where  $x_i \in \mathbb{R}$ . The coefficients  $x_1, x_2, \dots, x_n$  are called the **coordinates** of  $\mathbf{v}$  with respect to the ordered basis  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ .

The mapping

$$\text{vector } \mathbf{v} \mapsto \text{its coordinates } (x_1, x_2, \dots, x_n)$$

is a one-to-one correspondence between  $V$  and  $\mathbb{R}^n$ . This correspondence respects linear operations in  $V$  and in  $\mathbb{R}^n$ .

*Examples.* • Coordinates of a vector

$\mathbf{v} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$  relative to the standard basis  $\mathbf{e}_1 = (1, 0, \dots, 0, 0)$ ,  $\mathbf{e}_2 = (0, 1, \dots, 0, 0), \dots$ ,  $\mathbf{e}_n = (0, 0, \dots, 0, 1)$  are  $(x_1, x_2, \dots, x_n)$ .

• Coordinates of a matrix  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathcal{M}_{2,2}(\mathbb{R})$

relative to the basis  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$  are  $(a, b, c, d)$ .

• Coordinates of a polynomial

$p(x) = a_0 + a_1x + \dots + a_{n-1}x^{n-1} \in \mathcal{P}_n$  relative to the basis  $1, x, x^2, \dots, x^{n-1}$  are  $(a_0, a_1, \dots, a_{n-1})$ .

## Change of coordinates in $\mathbb{R}^n$

The usual (standard) coordinates of a vector  $\mathbf{v} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$  are coordinates relative to the standard basis  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ . Let  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$  be another basis for  $\mathbb{R}^n$  and  $(x'_1, x'_2, \dots, x'_n)$  be the coordinates of the same vector  $\mathbf{v}$  with respect to this basis. Then

$$\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ u_{21} & u_{22} & \dots & u_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ u_{n1} & u_{n2} & \dots & u_{nn} \end{pmatrix} \begin{pmatrix} x'_1 \\ x'_2 \\ \vdots \\ x'_n \end{pmatrix},$$

where the matrix  $U = (u_{ij})$  does not depend on the vector  $\mathbf{v}$ . Namely, columns of  $U$  are coordinates of vectors  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$  with respect to the standard basis.  $U$  is called the **transition matrix** from the basis  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$  to the standard basis  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ . The inverse matrix  $U^{-1}$  is called the **transition matrix** from  $\mathbf{e}_1, \dots, \mathbf{e}_n$  to  $\mathbf{u}_1, \dots, \mathbf{u}_n$ .

**Problem.** Find coordinates of the vector  $\mathbf{v} = (1, 2, 3)$  with respect to the basis  $\mathbf{u}_1 = (1, 1, 0)$ ,  $\mathbf{u}_2 = (0, 1, 1)$ ,  $\mathbf{u}_3 = (1, 1, 1)$ .

The nonstandard coordinates  $(x', y', z')$  of  $\mathbf{v}$  satisfy

$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = U \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix},$$

where  $U$  is the transition matrix from the standard basis  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  to the basis  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ .

The transition matrix from  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  to  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  is

$$U_0 = (\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3) = \left( \begin{array}{c|c|c} 1 & 0 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{array} \right).$$

The transition matrix from  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  to  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  is the inverse matrix:  $U = U_0^{-1}$ .

The inverse matrix can be computed using row reduction.

$$(U_0 | I) = \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{array} \right)$$

$$\rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{array} \right) \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & 1 & -1 & 1 \end{array} \right)$$

$$\rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & 1 & -1 & 1 \end{array} \right) = (I | U_0^{-1})$$

Thus


$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 0 & 1 & -1 \\ -1 & 1 & 0 \\ 1 & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}.$$

## Change of coordinates: general case

Let  $V$  be a vector space of dimension  $n$ .

Let  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  be a basis for  $V$  and  $g_1 : V \rightarrow \mathbb{R}^n$  be the coordinate mapping corresponding to this basis.

Let  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$  be another basis for  $V$  and  $g_2 : V \rightarrow \mathbb{R}^n$  be the coordinate mapping corresponding to this basis.



The composition  $g_2 \circ g_1^{-1}$  is a transformation of  $\mathbb{R}^n$ .

It has the form  $\mathbf{x} \mapsto U\mathbf{x}$ , where  $U$  is an  $n \times n$  matrix.

$U$  is called the **transition matrix** from  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  to  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ . Columns of  $U$  are coordinates of the vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  with respect to the basis  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ .

**Problem.** Find the transition matrix from the basis  $p_1(x) = 1$ ,  $p_2(x) = x + 1$ ,  $p_3(x) = (x + 1)^2$  to the basis  $q_1(x) = 1$ ,  $q_2(x) = x$ ,  $q_3(x) = x^2$  for the vector space  $\mathcal{P}_3$ .

We have to find coordinates of the polynomials  $p_1, p_2, p_3$  with respect to the basis  $q_1, q_2, q_3$ :

$$p_1(x) = 1 = q_1(x),$$

$$p_2(x) = x + 1 = q_1(x) + q_2(x),$$

$$p_3(x) = (x+1)^2 = x^2 + 2x + 1 = q_1(x) + 2q_2(x) + q_3(x).$$

Hence the transition matrix is 
$$\begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix}.$$



Thus the polynomial identity

$$a_1 + a_2(x + 1) + a_3(x + 1)^2 = b_1 + b_2x + b_3x^2$$

is equivalent to the relation

$$\begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}.$$

**Problem.** Find the transition matrix from the basis  $\mathbf{v}_1 = (1, 2, 3)$ ,  $\mathbf{v}_2 = (1, 0, 1)$ ,  $\mathbf{v}_3 = (1, 2, 1)$  to the basis  $\mathbf{u}_1 = (1, 1, 0)$ ,  $\mathbf{u}_2 = (0, 1, 1)$ ,  $\mathbf{u}_3 = (1, 1, 1)$ .

It is convenient to make a two-step transition: first from  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  to  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ , and then from  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  to  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ .

Let  $U_1$  be the transition matrix from  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  to  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  and  $U_2$  be the transition matrix from  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  to  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ :

$$U_1 = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 0 & 2 \\ 3 & 1 & 1 \end{pmatrix}, \quad U_2 = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}.$$

Basis  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3 \implies$  coordinates  $\mathbf{x}$

Basis  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3 \implies$  coordinates  $U_1\mathbf{x}$

Basis  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3 \implies$  coordinates  $U_2^{-1}(U_1\mathbf{x}) = (U_2^{-1}U_1)\mathbf{x}$

Thus the transition matrix from  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  to  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  is  $U_2^{-1}U_1$ .

$$\begin{aligned} U_2^{-1}U_1 &= \begin{pmatrix} 1 & 0 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 & 1 & 1 \\ 2 & 0 & 2 \\ 3 & 1 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 0 & 1 & -1 \\ -1 & 1 & 0 \\ 1 & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 2 & 0 & 2 \\ 3 & 1 & 1 \end{pmatrix} = \begin{pmatrix} -1 & -1 & 1 \\ 1 & -1 & 1 \\ 2 & 2 & 0 \end{pmatrix}. \end{aligned}$$

## Linear mapping = linear transformation = linear function

*Definition.* Given vector spaces  $V_1$  and  $V_2$ , a mapping  $L : V_1 \rightarrow V_2$  is **linear** if

$$L(\mathbf{x} + \mathbf{y}) = L(\mathbf{x}) + L(\mathbf{y}),$$

$$L(r\mathbf{x}) = rL(\mathbf{x})$$

for any  $\mathbf{x}, \mathbf{y} \in V_1$  and  $r \in \mathbb{R}$ .

A linear mapping  $\ell : V \rightarrow \mathbb{R}$  is called a **linear functional** on  $V$ .

If  $V_1 = V_2$  (or if both  $V_1$  and  $V_2$  are functional spaces) then a linear mapping  $L : V_1 \rightarrow V_2$  is called a **linear operator**.

## Linear mapping = linear transformation = linear function

*Definition.* Given vector spaces  $V_1$  and  $V_2$ , a mapping  $L : V_1 \rightarrow V_2$  is **linear** if

$$L(\mathbf{x} + \mathbf{y}) = L(\mathbf{x}) + L(\mathbf{y}),$$

$$L(r\mathbf{x}) = rL(\mathbf{x})$$

for any  $\mathbf{x}, \mathbf{y} \in V_1$  and  $r \in \mathbb{R}$ .

*Remark.* A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) = ax + b$  is a linear transformation of the vector space  $\mathbb{R}$  if and only if  $b = 0$ .

## Examples of linear mappings

- *Scaling*  $L : V \rightarrow V$ ,  $L(\mathbf{v}) = s\mathbf{v}$ , where  $s \in \mathbb{R}$ .

$$L(\mathbf{x} + \mathbf{y}) = s(\mathbf{x} + \mathbf{y}) = s\mathbf{x} + s\mathbf{y} = L(\mathbf{x}) + L(\mathbf{y}),$$

$$L(r\mathbf{x}) = s(r\mathbf{x}) = r(s\mathbf{x}) = rL(\mathbf{x}).$$

- *Dot product with a fixed vector*

$$\ell : \mathbb{R}^n \rightarrow \mathbb{R}, \ell(\mathbf{v}) = \mathbf{v} \cdot \mathbf{v}_0, \text{ where } \mathbf{v}_0 \in \mathbb{R}^n.$$

$$\ell(\mathbf{x} + \mathbf{y}) = (\mathbf{x} + \mathbf{y}) \cdot \mathbf{v}_0 = \mathbf{x} \cdot \mathbf{v}_0 + \mathbf{y} \cdot \mathbf{v}_0 = \ell(\mathbf{x}) + \ell(\mathbf{y}),$$

$$\ell(r\mathbf{x}) = (r\mathbf{x}) \cdot \mathbf{v}_0 = r(\mathbf{x} \cdot \mathbf{v}_0) = r\ell(\mathbf{x}).$$

- *Cross product with a fixed vector*

$$L : \mathbb{R}^3 \rightarrow \mathbb{R}^3, L(\mathbf{v}) = \mathbf{v} \times \mathbf{v}_0, \text{ where } \mathbf{v}_0 \in \mathbb{R}^3.$$

- *Multiplication by a fixed matrix*

$$L : \mathbb{R}^n \rightarrow \mathbb{R}^m, L(\mathbf{v}) = A\mathbf{v}, \text{ where } A \text{ is an } m \times n \text{ matrix and all vectors are column vectors.}$$

## Linear mappings of functional vector spaces

- *Evaluation at a fixed point*

$$\ell : F(\mathbb{R}) \rightarrow \mathbb{R}, \quad \ell(f) = f(a), \quad \text{where } a \in \mathbb{R}.$$

- *Multiplication by a fixed function*

$$L : F(\mathbb{R}) \rightarrow F(\mathbb{R}), \quad L(f) = gf, \quad \text{where } g \in F(\mathbb{R}).$$

- *Differentiation*  $D : C^1(\mathbb{R}) \rightarrow C(\mathbb{R}), \quad L(f) = f'.$

$$D(f + g) = (f + g)' = f' + g' = D(f) + D(g),$$

$$D(rf) = (rf)' = rf' = rD(f).$$

- *Integration over a finite interval*

$$\ell : C(\mathbb{R}) \rightarrow \mathbb{R}, \quad \ell(f) = \int_a^b f(x) dx, \quad \text{where}$$

$$a, b \in \mathbb{R}, \quad a < b.$$

## Linear differential operators

- an ordinary differential operator

$$L : C^\infty(\mathbb{R}) \rightarrow C^\infty(\mathbb{R}), \quad L = g_0 \frac{d^2}{dx^2} + g_1 \frac{d}{dx} + g_2,$$

where  $g_0, g_1, g_2$  are smooth functions on  $\mathbb{R}$ .

That is,  $L(f) = g_0 f'' + g_1 f' + g_2 f$ .

- Laplace's operator  $\Delta : C^\infty(\mathbb{R}^2) \rightarrow C^\infty(\mathbb{R}^2)$ ,

$$\Delta f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}$$

(a.k.a. the Laplacian; also denoted by  $\nabla^2$ ).



## Linear integral operators

- anti-derivative

$$L : C[a, b] \rightarrow C^1[a, b], \quad (Lf)(x) = \int_a^x f(y) dy.$$

- Hilbert-Schmidt operator

$$L : C[a, b] \rightarrow C[c, d], \quad (Lf)(x) = \int_a^b K(x, y)f(y) dy,$$

where  $K \in C([c, d] \times [a, b])$ .

- Laplace transform

$$\mathcal{L} : BC(0, \infty) \rightarrow C(0, \infty), \quad (\mathcal{L}f)(x) = \int_0^{\infty} e^{-xy} f(y) dy.$$