

MATH 304
Linear Algebra

Lecture 27:
Diagonalization (continued).
Euclidean structure in \mathbb{R}^n .

Diagonalization

Theorem 1 Let L be a linear operator on a finite-dimensional vector space V . Then the following conditions are equivalent:

- the matrix of L with respect to some basis is diagonal;
- there exists a basis for V formed by eigenvectors of L .

The operator L is **diagonalizable** if it satisfies these conditions.

Theorem 2 Let A be an $n \times n$ matrix. Then the following conditions are equivalent:

- A is the matrix of a diagonalizable operator;
- A is similar to a diagonal matrix, i.e., it is represented as $A = UBU^{-1}$, where the matrix B is diagonal;
- there exists a basis for \mathbb{R}^n formed by eigenvectors of A .

The matrix A is **diagonalizable** if it satisfies these conditions.

To *diagonalize* an $n \times n$ matrix A is to find a diagonal matrix B and an invertible matrix U such that $A = UBU^{-1}$.

Suppose there exists a basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ for \mathbb{R}^n consisting of eigenvectors of A . That is, $A\mathbf{v}_k = \lambda_k\mathbf{v}_k$, where $\lambda_k \in \mathbb{R}$.

Then $A = UBU^{-1}$, where $B = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$ and U is a transition matrix whose columns are vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$.

Example. $A = \begin{pmatrix} 4 & 3 \\ 0 & 1 \end{pmatrix}$. $\det(A - \lambda I) = (4 - \lambda)(1 - \lambda)$.

Eigenvalues: $\lambda_1 = 4$, $\lambda_2 = 1$.

Associated eigenvectors: $\mathbf{v}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\mathbf{v}_2 = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$.

Thus $A = UBU^{-1}$, where

$$B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Suppose we have a problem that involves a square matrix A in the context of matrix multiplication.

Also, suppose that the case when A is a diagonal matrix is simple. Then the diagonalization may help in solving this problem (or may not). Namely, it may reduce the case of a diagonalizable matrix to that of a diagonal one.

An example of such a problem is, given a square matrix A , to find its power A^k :

$$A = \begin{pmatrix} s_1 & & & 0 \\ & s_2 & & \\ & & \ddots & \\ 0 & & & s_n \end{pmatrix} \implies A^k = \begin{pmatrix} s_1^k & & & 0 \\ & s_2^k & & \\ & & \ddots & \\ 0 & & & s_n^k \end{pmatrix}$$

Problem. Let $A = \begin{pmatrix} 4 & 3 \\ 0 & 1 \end{pmatrix}$. Find A^5 .

We know that $A = UBU^{-1}$, where

$$B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Then $A^5 = UBU^{-1}UBU^{-1}UBU^{-1}UBU^{-1}UBU^{-1}$

$$= UB^5U^{-1} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1024 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 1024 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1024 & 1023 \\ 0 & 1 \end{pmatrix}.$$

Problem. Let $A = \begin{pmatrix} 4 & 3 \\ 0 & 1 \end{pmatrix}$. Find A^k ($k \geq 1$).

We know that $A = UBU^{-1}$, where

$$B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Then

$$\begin{aligned} A^k &= UB^kU^{-1} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 4^k & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 4^k & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 4^k & 4^k - 1 \\ 0 & 1 \end{pmatrix}. \end{aligned}$$

Problem. Let $A = \begin{pmatrix} 4 & 3 \\ 0 & 1 \end{pmatrix}$. Find a matrix C such that $C^2 = A$.

We know that $A = UBU^{-1}$, where

$$B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Suppose that $D^2 = B$ for some matrix D . Let $C = UDU^{-1}$. Then $C^2 = UDU^{-1}UDU^{-1} = UD^2U^{-1} = UBU^{-1} = A$.

We can take $D = \begin{pmatrix} \sqrt{4} & 0 \\ 0 & \sqrt{1} \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$.

Then $C = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$.

Initial value problem for a system of linear ODEs:

$$\begin{cases} \frac{dx}{dt} = 4x + 3y, \\ \frac{dy}{dt} = y, \end{cases} \quad x(0) = 1, \quad y(0) = 1.$$

The system can be rewritten in vector form:

$$\frac{d\mathbf{v}}{dt} = A\mathbf{v}, \quad \text{where } A = \begin{pmatrix} 4 & 3 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} x \\ y \end{pmatrix}.$$

Matrix A is diagonalizable: $A = UBU^{-1}$, where

$$B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Let $\mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$ be coordinates of the vector \mathbf{v} relative to the basis $\mathbf{v}_1 = (1, 0)$, $\mathbf{v}_2 = (-1, 1)$ of eigenvectors of A . Then $\mathbf{v} = U\mathbf{w} \implies \mathbf{w} = U^{-1}\mathbf{v}$.

It follows that

$$\frac{d\mathbf{w}}{dt} = \frac{d}{dt}(U^{-1}\mathbf{v}) = U^{-1}\frac{d\mathbf{v}}{dt} = U^{-1}A\mathbf{v} = U^{-1}AU\mathbf{w}.$$

$$\text{Hence } \frac{d\mathbf{w}}{dt} = B\mathbf{w} \iff \begin{cases} \frac{dw_1}{dt} = 4w_1, \\ \frac{dw_2}{dt} = w_2. \end{cases}$$

General solution: $w_1(t) = c_1 e^{4t}$, $w_2(t) = c_2 e^t$, where $c_1, c_2 \in \mathbb{R}$.

Initial condition:

$$\mathbf{w}(0) = U^{-1}\mathbf{v}(0) = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}.$$

Thus $w_1(t) = 2e^{4t}$, $w_2(t) = e^t$. Then

$$\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = U\mathbf{w}(t) = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2e^{4t} \\ e^t \end{pmatrix} = \begin{pmatrix} 2e^{4t} - e^t \\ e^t \end{pmatrix}.$$

There are **two obstructions** to existence of a basis consisting of eigenvectors. They are illustrated by the following examples.

Example 1. $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

$\det(A - \lambda I) = (\lambda - 1)^2$. Hence $\lambda = 1$ is the only eigenvalue. The associated eigenspace is the line $t(1, 0)$.


Example 2. $A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

$\det(A - \lambda I) = \lambda^2 + 1$.

\implies no real eigenvalues or eigenvectors


(However there are *complex* eigenvalues/eigenvectors.)

Classical vectors: geometric approach



- A vector is represented by a directed segment.
- Directed segment is drawn as an arrow.
- Different arrows represent the same vector if they are of the same length and direction.


Classical vectors: geometric approach



\overrightarrow{AB} denotes a vector represented by the arrow with tip (endpoint) at B and tail (beginning) at A .

For any vector \mathbf{v} and a point A there is a unique point B such that $\overrightarrow{AB} = \mathbf{v}$.

Vectors: geometric approach




If $\mathbf{v} = \overrightarrow{AB}$ then \overrightarrow{BA} is called the *negative vector* of \mathbf{v} and denoted $-\mathbf{v}$.

\overrightarrow{AA} is called the *zero vector* and denoted $\mathbf{0}$.

Linear structure: vector addition


Given vectors \mathbf{a} and \mathbf{b} , their *sum* $\mathbf{a} + \mathbf{b}$ is defined by the rule $\overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC}$.

That is, choose points A, B, C so that $\overrightarrow{AB} = \mathbf{a}$ and $\overrightarrow{BC} = \mathbf{b}$. Then $\mathbf{a} + \mathbf{b} = \overrightarrow{AC}$.



Linear structure: scalar multiplication


Let \mathbf{v} be a vector and $r \in \mathbb{R}$. By definition, $r\mathbf{v}$ is a vector whose magnitude is $|r|$ times the magnitude of \mathbf{v} . The direction of $r\mathbf{v}$ coincides with that of \mathbf{v} if $r > 0$. If $r < 0$ then the directions of $r\mathbf{v}$ and \mathbf{v} are opposite.




Beyond linearity: Euclidean structure

The Euclidean structure includes:

- length of a vector: $|\mathbf{x}|$,
- angle between vectors: θ ,
- dot product: $\mathbf{x} \cdot \mathbf{y} = |\mathbf{x}| |\mathbf{y}| \cos \theta$.



Cartesian coordinates: geometric meets algebraic



Cartesian coordinates allow us to identify a line, a plane, and space with \mathbb{R} , \mathbb{R}^2 , and \mathbb{R}^3 , respectively.

Once we specify an *origin* O , each point A is associated a *position vector* \vec{OA} . Conversely, every vector has a unique representative with tail at O .

Length and distance

Definition. The **length** of a vector $\mathbf{v} = (v_1, v_2, \dots, v_n) \in \mathbb{R}^n$ is

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}.$$

The **distance** between vectors \mathbf{x} and \mathbf{y} is defined as $\|\mathbf{y} - \mathbf{x}\|$.

Properties of length:

$$\|\mathbf{x}\| \geq 0, \quad \|\mathbf{x}\| = 0 \text{ only if } \mathbf{x} = \mathbf{0} \quad (\text{positivity})$$

$$\|r\mathbf{x}\| = |r| \|\mathbf{x}\| \quad (\text{homogeneity})$$

$$\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\| \quad (\text{triangle inequality})$$

Scalar product

Definition. The **scalar product** of vectors $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and $\mathbf{y} = (y_1, y_2, \dots, y_n)$ is

$$\mathbf{x} \cdot \mathbf{y} = x_1y_1 + x_2y_2 + \dots + x_ny_n.$$

Alternative notation: (\mathbf{x}, \mathbf{y}) or $\langle \mathbf{x}, \mathbf{y} \rangle$.

Properties of scalar product:

$$\mathbf{x} \cdot \mathbf{x} \geq 0, \quad \mathbf{x} \cdot \mathbf{x} = 0 \text{ only if } \mathbf{x} = \mathbf{0} \quad (\text{positivity})$$

$$\mathbf{x} \cdot \mathbf{y} = \mathbf{y} \cdot \mathbf{x} \quad (\text{symmetry})$$

$$(\mathbf{x} + \mathbf{y}) \cdot \mathbf{z} = \mathbf{x} \cdot \mathbf{z} + \mathbf{y} \cdot \mathbf{z} \quad (\text{distributive law})$$

$$(r\mathbf{x}) \cdot \mathbf{y} = r(\mathbf{x} \cdot \mathbf{y}) \quad (\text{homogeneity})$$

In particular, $\mathbf{x} \cdot \mathbf{y}$ is a **bilinear** function (i.e., it is both a linear function of \mathbf{x} and a linear function of \mathbf{y}).

Relations between lengths and scalar products:

$$\|\mathbf{x}\| = \sqrt{\mathbf{x} \cdot \mathbf{x}}$$

$$|\mathbf{x} \cdot \mathbf{y}| \leq \|\mathbf{x}\| \|\mathbf{y}\| \quad (\text{Cauchy-Schwarz inequality})$$

$$\|\mathbf{x} - \mathbf{y}\|^2 = \|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 - 2\mathbf{x} \cdot \mathbf{y}$$

By the Cauchy-Schwarz inequality, for any nonzero vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ we have

$$\cos \theta = \frac{\mathbf{x} \cdot \mathbf{y}}{\|\mathbf{x}\| \|\mathbf{y}\|} \quad \text{for some } 0 \leq \theta \leq \pi.$$

θ is called the **angle** between the vectors \mathbf{x} and \mathbf{y} .

The vectors \mathbf{x} and \mathbf{y} are said to be **orthogonal** (denoted $\mathbf{x} \perp \mathbf{y}$) if $\mathbf{x} \cdot \mathbf{y} = 0$ (i.e., if $\theta = 90^\circ$).

Problem. Find the angle θ between vectors $\mathbf{x} = (2, -1)$ and $\mathbf{y} = (3, 1)$.

$$\mathbf{x} \cdot \mathbf{y} = 5, \quad \|\mathbf{x}\| = \sqrt{5}, \quad \|\mathbf{y}\| = \sqrt{10}.$$

$$\cos \theta = \frac{\mathbf{x} \cdot \mathbf{y}}{\|\mathbf{x}\| \|\mathbf{y}\|} = \frac{5}{\sqrt{5} \sqrt{10}} = \frac{1}{\sqrt{2}} \implies \theta = 45^\circ$$

Problem. Find the angle ϕ between vectors $\mathbf{v} = (-2, 1, 3)$ and $\mathbf{w} = (4, 5, 1)$.

$$\mathbf{v} \cdot \mathbf{w} = 0 \implies \mathbf{v} \perp \mathbf{w} \implies \phi = 90^\circ$$