

MATH 311-504

Topics in Applied Mathematics

**Lecture 3-3:**

**Norms and inner products.**

## Norm

The notion of *norm* generalizes the notion of length of a vector in  $\mathbb{R}^n$ .

*Definition.* Let  $V$  be a vector space. A function  $\alpha : V \rightarrow \mathbb{R}$  is called a **norm** on  $V$  if it has the following properties:

- (i)  $\alpha(\mathbf{x}) \geq 0$ ,  $\alpha(\mathbf{x}) = 0$  only for  $\mathbf{x} = \mathbf{0}$  (positivity)
- (ii)  $\alpha(r\mathbf{x}) = |r| \alpha(\mathbf{x})$  for all  $r \in \mathbb{R}$  (homogeneity)
- (iii)  $\alpha(\mathbf{x} + \mathbf{y}) \leq \alpha(\mathbf{x}) + \alpha(\mathbf{y})$  (triangle inequality)

*Notation.* The norm of a vector  $\mathbf{x} \in V$  is usually denoted  $\|\mathbf{x}\|$ . Different norms on  $V$  are distinguished by subscripts, e.g.,  $\|\mathbf{x}\|_1$  and  $\|\mathbf{x}\|_2$ .

*Examples.*  $V = \mathbb{R}^n$ ,  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ .

- $\|\mathbf{x}\|_\infty = \max(|x_1|, |x_2|, \dots, |x_n|)$ .
- $\|\mathbf{x}\|_p = (|x_1|^p + |x_2|^p + \dots + |x_n|^p)^{1/p}$ ,  $p \geq 1$ .

In particular,


- $\|\mathbf{x}\|_1 = |x_1| + |x_2| + \dots + |x_n|$ ,
- $\|\mathbf{x}\|_2 = (|x_1|^2 + |x_2|^2 + \dots + |x_n|^2)^{1/2} = |\mathbf{x}|$ .

*Definition.* A **normed vector space** is a vector space endowed with a norm.

The norm defines a distance function on the normed vector space:  $\text{dist}(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|$ .

Then we say that a sequence  $\mathbf{x}_1, \mathbf{x}_2, \dots$  converges to a vector  $\mathbf{x}$  if  $\text{dist}(\mathbf{x}, \mathbf{x}_n) \rightarrow 0$  as  $n \rightarrow \infty$ .

Unit circle:  $\|\mathbf{x}\| = 1$



$$\|\mathbf{x}\| = (x_1^2 + x_2^2)^{1/2} \quad \text{black}$$

$$\|\mathbf{x}\| = \left(\frac{1}{2}x_1^2 + x_2^2\right)^{1/2} \quad \text{green}$$

$$\|\mathbf{x}\| = |x_1| + |x_2| \quad \text{blue}$$

$$\|\mathbf{x}\| = \max(|x_1|, |x_2|) \quad \text{red}$$

*Examples.*  $V = C[a, b]$ ,  $f : [a, b] \rightarrow \mathbb{R}$ .

- $\|f\|_\infty = \max_{a \leq x \leq b} |f(x)|$  (uniform norm).

- $\|f\|_1 = \int_a^b |f(x)| dx$ .

- $\|f\|_p = \left( \int_a^b |f(x)|^p dx \right)^{1/p}$ ,  $p > 0$ .

**Theorem**  $\|f\|_p$  is a norm on  $C[a, b]$  for any  $p \geq 1$ .

## Inner product

The notion of *inner product* generalizes the notion of dot product of vectors in  $\mathbb{R}^n$ .

*Definition.* Let  $V$  be a vector space. A function  $\beta : V \times V \rightarrow \mathbb{R}$ , usually denoted  $\beta(\mathbf{x}, \mathbf{y}) = \langle \mathbf{x}, \mathbf{y} \rangle$ , is called an **inner product** on  $V$  if it is positive, symmetric, and bilinear. That is, if

- (i)  $\langle \mathbf{x}, \mathbf{y} \rangle \geq 0$ ,  $\langle \mathbf{x}, \mathbf{x} \rangle = 0$  only for  $\mathbf{x} = \mathbf{0}$  (positivity)
- (ii)  $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$  (symmetry)
- (iii)  $\langle r\mathbf{x}, \mathbf{y} \rangle = r\langle \mathbf{x}, \mathbf{y} \rangle$  (homogeneity)
- (iv)  $\langle \mathbf{x} + \mathbf{y}, \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{z} \rangle + \langle \mathbf{y}, \mathbf{z} \rangle$  (distributive law)

An **inner product space** is a vector space endowed with an inner product.

*Examples.*  $V = \mathbb{R}^n$ .

- $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x} \cdot \mathbf{y} = x_1y_1 + x_2y_2 + \cdots + x_ny_n$ .

- $\langle \mathbf{x}, \mathbf{y} \rangle = d_1x_1y_1 + d_2x_2y_2 + \cdots + d_nx_ny_n$ ,

where  $d_1, d_2, \dots, d_n > 0$ .

- $\langle \mathbf{x}, \mathbf{y} \rangle = (D\mathbf{x}) \cdot (D\mathbf{y})$ ,

where  $D$  is an invertible  $n \times n$  matrix.

*Remarks.* (a) Invertibility of  $D$  is necessary to show that  $\langle \mathbf{x}, \mathbf{x} \rangle = 0 \implies \mathbf{x} = \mathbf{0}$ .

(b) The second example is a particular case of the third one when  $D = \text{diag}(d_1^{1/2}, d_2^{1/2}, \dots, d_n^{1/2})$ .



*Examples.*  $V = C[a, b]$ .

- $\langle f, g \rangle = \int_a^b f(x)g(x) dx.$

- $\langle f, g \rangle = \int_a^b f(x)g(x)w(x) dx,$

where  $w$  is bounded, piecewise continuous, and  $w > 0$  everywhere on  $[a, b]$ .

$w$  is called the **weight** function.

*Counterexamples.*  $V = \mathbb{R}^2$ .

- $\langle \mathbf{x}, \mathbf{y} \rangle = x_1y_1 - x_2y_2$ .

Let  $\mathbf{v} = (1, 2)$ , then  $\langle \mathbf{v}, \mathbf{v} \rangle = 1^2 - 2^2 = -3$ .

$\langle \mathbf{x}, \mathbf{y} \rangle$  is symmetric and bilinear, but not positive.

- $\langle \mathbf{x}, \mathbf{y} \rangle = 2x_1y_1 + x_1x_2 + 2x_2y_2 + y_1y_2$ .

$\mathbf{v} = (1, 1)$ ,  $\mathbf{w} = (1, 0) \implies \langle \mathbf{v}, \mathbf{w} \rangle = 3$ ,  $\langle 2\mathbf{v}, \mathbf{w} \rangle = 8$ .

$\langle \mathbf{x}, \mathbf{y} \rangle$  is positive and symmetric, but not bilinear.

- $\langle \mathbf{x}, \mathbf{y} \rangle = x_1y_1 + x_1y_2 - x_2y_1 + x_2y_2$ .

$\mathbf{v} = (1, 1)$ ,  $\mathbf{w} = (1, 0) \implies \langle \mathbf{v}, \mathbf{w} \rangle = 0$ ,  $\langle \mathbf{w}, \mathbf{v} \rangle = 2$ .

$\langle \mathbf{x}, \mathbf{y} \rangle$  is positive and bilinear, but not symmetric.

*Problem.* Find an inner product on  $\mathbb{R}^2$  such that  $\langle \mathbf{e}_1, \mathbf{e}_1 \rangle = 2$ ,  $\langle \mathbf{e}_2, \mathbf{e}_2 \rangle = 3$ , and  $\langle \mathbf{e}_1, \mathbf{e}_2 \rangle = -1$ , where  $\mathbf{e}_1 = (1, 0)$ ,  $\mathbf{e}_2 = (0, 1)$ .

Let  $\mathbf{x} = (x_1, x_2)$ ,  $\mathbf{y} = (y_1, y_2) \in \mathbb{R}^2$ .

Then  $\mathbf{x} = x_1\mathbf{e}_1 + x_2\mathbf{e}_2$ ,  $\mathbf{y} = y_1\mathbf{e}_1 + y_2\mathbf{e}_2$ .

It follows that

$$\begin{aligned}\langle \mathbf{x}, \mathbf{y} \rangle &= \langle x_1\mathbf{e}_1 + x_2\mathbf{e}_2, y_1\mathbf{e}_1 + y_2\mathbf{e}_2 \rangle \\ &= x_1\langle \mathbf{e}_1, y_1\mathbf{e}_1 + y_2\mathbf{e}_2 \rangle + x_2\langle \mathbf{e}_2, y_1\mathbf{e}_1 + y_2\mathbf{e}_2 \rangle \\ &= x_1y_1\langle \mathbf{e}_1, \mathbf{e}_1 \rangle + x_1y_2\langle \mathbf{e}_1, \mathbf{e}_2 \rangle + x_2y_1\langle \mathbf{e}_2, \mathbf{e}_1 \rangle + x_2y_2\langle \mathbf{e}_2, \mathbf{e}_2 \rangle \\ &= 2x_1y_1 - x_1y_2 - x_2y_1 + 3x_2y_2.\end{aligned}$$

**Theorem** Suppose  $\langle \mathbf{x}, \mathbf{y} \rangle$  is an inner product on a vector space  $V$ . Then

$$\langle \mathbf{x}, \mathbf{y} \rangle^2 \leq \langle \mathbf{x}, \mathbf{x} \rangle \langle \mathbf{y}, \mathbf{y} \rangle \quad \text{for all } \mathbf{x}, \mathbf{y} \in V.$$

*Proof:* For any  $t \in \mathbb{R}$  let  $\mathbf{v}_t = \mathbf{x} + t\mathbf{y}$ . Then

$$\langle \mathbf{v}_t, \mathbf{v}_t \rangle = \langle \mathbf{x}, \mathbf{x} \rangle + 2t\langle \mathbf{x}, \mathbf{y} \rangle + t^2\langle \mathbf{y}, \mathbf{y} \rangle.$$

The right-hand side is a quadratic polynomial in  $t$  (provided that  $\mathbf{y} \neq \mathbf{0}$ ). Since  $\langle \mathbf{v}_t, \mathbf{v}_t \rangle \geq 0$  for all  $t$ , the discriminant  $D$  is nonpositive. But

$$D = 4\langle \mathbf{x}, \mathbf{y} \rangle^2 - 4\langle \mathbf{x}, \mathbf{x} \rangle \langle \mathbf{y}, \mathbf{y} \rangle.$$

**Cauchy-Schwarz Inequality:**

$$|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} \sqrt{\langle \mathbf{y}, \mathbf{y} \rangle}.$$

## Cauchy-Schwarz Inequality:

$$|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} \sqrt{\langle \mathbf{y}, \mathbf{y} \rangle}.$$

**Corollary 1**  $|\mathbf{x} \cdot \mathbf{y}| \leq |\mathbf{x}| |\mathbf{y}|$  for all  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ .

Equivalently, for all  $x_i, y_i \in \mathbb{R}$ ,

$$(x_1 y_1 + \cdots + x_n y_n)^2 \leq (x_1^2 + \cdots + x_n^2)(y_1^2 + \cdots + y_n^2).$$

**Corollary 2** For any  $f, g \in C[a, b]$ ,

$$\left( \int_a^b f(x)g(x) dx \right)^2 \leq \int_a^b |f(x)|^2 dx \cdot \int_a^b |g(x)|^2 dx.$$