

MATH 311

Topics in Applied Mathematics I

**Lecture 17:**

**Nullity of a matrix.**

**Basis and coordinates.**

**Change of basis.**

## Rank of a matrix

*Definition.* The **row space** of an  $m \times n$  matrix  $A$  is the subspace of  $\mathbb{R}^n$  spanned by rows of  $A$ . The **column space** of  $A$  is a subspace of  $\mathbb{R}^m$  spanned by columns of  $A$ .

The row space and the column space of  $A$  have the same dimension, which is called the **rank** of  $A$ .

**Theorem 1** Elementary row operations do not change the row space of a matrix.

**Theorem 2** If a matrix  $A$  is in row echelon form, then the nonzero rows of  $A$  form a basis for the row space.

**Theorem 3** The rank of a matrix is equal to the number of nonzero rows in its row echelon form.

## Nullspace of a matrix

Let  $A = (a_{ij})$  be an  $m \times n$  matrix.

*Definition.* The **nullspace** of the matrix  $A$ , denoted  $N(A)$ , is the set of all  $n$ -dimensional column vectors  $\mathbf{x}$  such that

$$\mathbf{Ax} = \mathbf{0}.$$

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

The nullspace  $N(A)$  is the solution set of a system of linear homogeneous equations (with  $A$  as the coefficient matrix).

**Theorem**  $N(A)$  is a subspace of the vector space  $\mathbb{R}^n$ .

*Definition.* The dimension of the nullspace  $N(A)$  is called the **nullity** of the matrix  $A$ .

## rank + nullity

**Theorem** The rank of a matrix  $A$  plus the nullity of  $A$  equals the number of columns in  $A$ .

*Sketch of the proof:* The rank of  $A$  equals the number of nonzero rows in the row echelon form, which equals the number of leading entries.

The nullity of  $A$  equals the number of free variables in the corresponding homogeneous system, which equals the number of columns without leading entries in the row echelon form.

Consequently, rank+nullity is the number of all columns in the matrix  $A$ .

**Problem.** Find the nullity of the matrix

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 2 & 3 & 4 & 5 \end{pmatrix}.$$

Clearly, the rows of  $A$  are linearly independent.

Therefore the rank of  $A$  is 2. Since

$$(\text{rank of } A) + (\text{nullity of } A) = 4,$$

it follows that the nullity of  $A$  is 2.

## Basis and dimension

*Definition.* Let  $V$  be a vector space. A linearly independent spanning set for  $V$  is called a **basis**.

**Theorem** Any vector space  $V$  has a basis. If  $V$  has a finite basis, then all bases for  $V$  are finite and have the same number of elements (called the *dimension* of  $V$ ).

*Example.* Vectors  $\mathbf{e}_1 = (1, 0, 0, \dots, 0, 0)$ ,  $\mathbf{e}_2 = (0, 1, 0, \dots, 0, 0), \dots, \mathbf{e}_n = (0, 0, 0, \dots, 0, 1)$  form a basis for  $\mathbb{R}^n$  (called *standard*) since

$$(x_1, x_2, \dots, x_n) = x_1\mathbf{e}_1 + x_2\mathbf{e}_2 + \cdots + x_n\mathbf{e}_n.$$

## Basis and coordinates

If  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  is a basis for a vector space  $V$ , then any vector  $\mathbf{v} \in V$  has a unique representation

$$\mathbf{v} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \dots + x_n\mathbf{v}_n,$$

where  $x_i \in \mathbb{R}$ . The coefficients  $x_1, x_2, \dots, x_n$  are called the **coordinates** of  $\mathbf{v}$  with respect to the ordered basis  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ .

The mapping

$$\text{vector } \mathbf{v} \mapsto \text{its coordinates } (x_1, x_2, \dots, x_n)$$

is a one-to-one correspondence between  $V$  and  $\mathbb{R}^n$ . This correspondence respects linear operations in  $V$  and in  $\mathbb{R}^n$ .

*Examples.* • Coordinates of a vector

$\mathbf{v} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$  relative to the standard basis  $\mathbf{e}_1 = (1, 0, \dots, 0, 0)$ ,  $\mathbf{e}_2 = (0, 1, \dots, 0, 0), \dots$ ,  $\mathbf{e}_n = (0, 0, \dots, 0, 1)$  are  $(x_1, x_2, \dots, x_n)$ .

• Coordinates of a matrix  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathcal{M}_{2,2}(\mathbb{R})$

relative to the basis  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$  are  $(a, c, b, d)$ .

• Coordinates of a polynomial

$p(x) = a_0 + a_1x + \dots + a_{n-1}x^{n-1} \in \mathcal{P}_n$  relative to the basis  $1, x, x^2, \dots, x^{n-1}$  are  $(a_0, a_1, \dots, a_{n-1})$ .



Vectors  $\mathbf{u}_1=(3, 1)$  and  $\mathbf{u}_2=(2, 1)$  form a basis for  $\mathbb{R}^2$ .

**Problem 1.** Find coordinates of the vector  $\mathbf{v} = (7, 4)$  with respect to the basis  $\mathbf{u}_1, \mathbf{u}_2$ .

The desired coordinates  $x, y$  satisfy

$$\mathbf{v} = x\mathbf{u}_1 + y\mathbf{u}_2 \iff \begin{cases} 3x + 2y = 7 \\ x + y = 4 \end{cases} \iff \begin{cases} x = -1 \\ y = 5 \end{cases}$$

**Problem 2.** Find the vector  $\mathbf{w}$  whose coordinates with respect to the basis  $\mathbf{u}_1, \mathbf{u}_2$  are  $(7, 4)$ .

$$\mathbf{w} = 7\mathbf{u}_1 + 4\mathbf{u}_2 = 7(3, 1) + 4(2, 1) = (29, 11)$$

## Change of coordinates

Given a vector  $\mathbf{v} \in \mathbb{R}^2$ , let  $(x, y)$  be its standard coordinates, i.e., coordinates with respect to the standard basis  $\mathbf{e}_1 = (1, 0)$ ,  $\mathbf{e}_2 = (0, 1)$ , and let  $(x', y')$  be its coordinates with respect to the basis  $\mathbf{u}_1 = (3, 1)$ ,  $\mathbf{u}_2 = (2, 1)$ .

**Problem.** Find a relation between  $(x, y)$  and  $(x', y')$ .

By definition,  $\mathbf{v} = x\mathbf{e}_1 + y\mathbf{e}_2 = x'\mathbf{u}_1 + y'\mathbf{u}_2$ .

In standard coordinates,

$$\begin{aligned} \begin{pmatrix} x \\ y \end{pmatrix} &= x' \begin{pmatrix} 3 \\ 1 \end{pmatrix} + y' \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x' \\ y' \end{pmatrix} \\ \implies \begin{pmatrix} x' \\ y' \end{pmatrix} &= \begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & -2 \\ -1 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \end{aligned}$$

## Change of coordinates in $\mathbb{R}^n$

The usual (standard) coordinates of a vector  $\mathbf{v} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$  are coordinates relative to the standard basis  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ . Let  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$  be another basis for  $\mathbb{R}^n$  and  $(x'_1, x'_2, \dots, x'_n)$  be the coordinates of the same vector  $\mathbf{v}$  with respect to this basis. Then

$$\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ u_{21} & u_{22} & \dots & u_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ u_{n1} & u_{n2} & \dots & u_{nn} \end{pmatrix} \begin{pmatrix} x'_1 \\ x'_2 \\ \vdots \\ x'_n \end{pmatrix},$$

where the matrix  $U = (u_{ij})$  does not depend on the vector  $\mathbf{v}$ . Namely, columns of  $U$  are coordinates of vectors  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$  with respect to the standard basis.  $U$  is called the **transition matrix** from the basis  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$  to the standard basis  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ . The inverse matrix  $U^{-1}$  is called the **transition matrix** from  $\mathbf{e}_1, \dots, \mathbf{e}_n$  to  $\mathbf{u}_1, \dots, \mathbf{u}_n$ .

**Problem.** Find coordinates of the vector  $\mathbf{x} = (1, 2, 3)$  with respect to the basis  $\mathbf{u}_1 = (1, 1, 0)$ ,  $\mathbf{u}_2 = (0, 1, 1)$ ,  $\mathbf{u}_3 = (1, 1, 1)$ .

The nonstandard coordinates  $(x', y', z')$  of  $\mathbf{x}$  satisfy

$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = U \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix},$$

where  $U$  is the transition matrix from the standard basis  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  to the basis  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ .

The transition matrix from  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  to  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  is

$$U_0 = (\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3) = \left( \begin{array}{c|c|c} 1 & 0 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{array} \right).$$

The transition matrix from  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  to  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  is the inverse matrix:  $U = U_0^{-1}$ .

The inverse matrix can be computed using row reduction.

$$(U_0 | I) = \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{array} \right)$$

$$\rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{array} \right) \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & 1 & -1 & 1 \end{array} \right)$$

$$\rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & 1 & -1 & 1 \end{array} \right) = (I | U_0^{-1})$$

Thus


$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 0 & 1 & -1 \\ -1 & 1 & 0 \\ 1 & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}.$$

## Change of coordinates: general case

Let  $V$  be a vector space of dimension  $n$ .

Let  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  be a basis for  $V$  and  $g_1 : V \rightarrow \mathbb{R}^n$  be the coordinate mapping corresponding to this basis.

Let  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$  be another basis for  $V$  and  $g_2 : V \rightarrow \mathbb{R}^n$  be the coordinate mapping corresponding to this basis.



The composition  $g_2 \circ g_1^{-1}$  is a transformation of  $\mathbb{R}^n$ .

It has the form  $\mathbf{x} \mapsto U\mathbf{x}$ , where  $U$  is an  $n \times n$  matrix.

$U$  is called the **transition matrix** from  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  to  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ . Columns of  $U$  are coordinates of the vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  with respect to the basis  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ .

**Problem.** Find the transition matrix from the basis  $p_1(x) = 1$ ,  $p_2(x) = x + 1$ ,  $p_3(x) = (x + 1)^2$  to the basis  $q_1(x) = 1$ ,  $q_2(x) = x$ ,  $q_3(x) = x^2$  for the vector space  $\mathcal{P}_3$ .

We have to find coordinates of the polynomials  $p_1, p_2, p_3$  with respect to the basis  $q_1, q_2, q_3$ :

$$p_1(x) = 1 = q_1(x),$$

$$p_2(x) = x + 1 = q_1(x) + q_2(x),$$

$$p_3(x) = (x+1)^2 = x^2 + 2x + 1 = q_1(x) + 2q_2(x) + q_3(x).$$

Hence the transition matrix is 
$$\begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix}.$$

Thus the polynomial identity

$$a_1 + a_2(x + 1) + a_3(x + 1)^2 = b_1 + b_2x + b_3x^2$$

is equivalent to the relation

$$\begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}.$$