

MATH 311

Topics in Applied Mathematics I

**Lecture 19:**

**Range and kernel.**

**General linear equations.**

**Matrix transformations.**

## Linear transformation

*Definition.* Given vector spaces  $V_1$  and  $V_2$ , a mapping  $L : V_1 \rightarrow V_2$  is **linear** if

$$L(\mathbf{x} + \mathbf{y}) = L(\mathbf{x}) + L(\mathbf{y}),$$

$$L(r\mathbf{x}) = rL(\mathbf{x})$$

for any  $\mathbf{x}, \mathbf{y} \in V_1$  and  $r \in \mathbb{R}$ .

*Basic properties of linear mappings:*

- $L(r_1\mathbf{v}_1 + \cdots + r_k\mathbf{v}_k) = r_1L(\mathbf{v}_1) + \cdots + r_kL(\mathbf{v}_k)$   
for all  $k \geq 1$ ,  $\mathbf{v}_1, \dots, \mathbf{v}_k \in V_1$ , and  $r_1, \dots, r_k \in \mathbb{R}$ .
- $L(\mathbf{0}_1) = \mathbf{0}_2$ , where  $\mathbf{0}_1$  and  $\mathbf{0}_2$  are zero vectors in  $V_1$  and  $V_2$ , respectively.
- $L(-\mathbf{v}) = -L(\mathbf{v})$  for any  $\mathbf{v} \in V_1$ .

## Range and kernel

Let  $V, W$  be vector spaces and  $L : V \rightarrow W$  be a linear mapping.

*Definition.* The **range** (or **image**) of  $L$  is the set of all vectors  $\mathbf{w} \in W$  such that  $\mathbf{w} = L(\mathbf{v})$  for some  $\mathbf{v} \in V$ . The range of  $L$  is denoted  $L(V)$ .

The **kernel** of  $L$ , denoted  $\ker L$ , is the set of all vectors  $\mathbf{v} \in V$  such that  $L(\mathbf{v}) = \mathbf{0}$ .

**Theorem** (i) The range of  $L$  is a subspace of  $W$ .  
(ii) The kernel of  $L$  is a subspace of  $V$ .

*Example.*  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ ,  $L \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 & 0 & -1 \\ 1 & 2 & -1 \\ 1 & 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ .

The kernel  $\ker(L)$  is the nullspace of the matrix.

$$L \begin{pmatrix} x \\ y \\ z \end{pmatrix} = x \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + y \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix} + z \begin{pmatrix} -1 \\ -1 \\ -1 \end{pmatrix}$$

The range  $L(\mathbb{R}^3)$  is the column space of the matrix.

*Example.*  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ ,  $L \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 & 0 & -1 \\ 1 & 2 & -1 \\ 1 & 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ .

The range of  $L$  is spanned by vectors  $(1, 1, 1)$ ,  $(0, 2, 0)$ , and  $(-1, -1, -1)$ . It follows that  $L(\mathbb{R}^3)$  is the plane spanned by  $(1, 1, 1)$  and  $(0, 1, 0)$ .

To find  $\ker(L)$ , we apply row reduction to the matrix:

$$\begin{pmatrix} 1 & 0 & -1 \\ 1 & 2 & -1 \\ 1 & 0 & -1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & -1 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Hence  $(x, y, z) \in \ker(L)$  if  $x - z = y = 0$ .

It follows that  $\ker(L)$  is the line spanned by  $(1, 0, 1)$ .

*Example.*  $L: C^3(\mathbb{R}) \rightarrow C(\mathbb{R})$ ,  $L(u) = u''' - 2u'' + u'$ .

According to the theory of differential equations, the initial value problem

$$\begin{cases} u'''(x) - 2u''(x) + u'(x) = g(x), & x \in \mathbb{R}, \\ u(a) = b_0, \\ u'(a) = b_1, \\ u''(a) = b_2 \end{cases}$$

has a unique solution for any  $g \in C(\mathbb{R})$  and any  $b_0, b_1, b_2 \in \mathbb{R}$ . It follows that  $L(C^3(\mathbb{R})) = C(\mathbb{R})$ .

Also, the initial data evaluation  $I(u) = (u(a), u'(a), u''(a))$ , which is a linear mapping  $I: C^3(\mathbb{R}) \rightarrow \mathbb{R}^3$ , becomes invertible when restricted to  $\ker(L)$ . Hence  $\dim \ker(L) = 3$ .

It is easy to check that  $L(xe^x) = L(e^x) = L(1) = 0$ .

Besides, the functions  $xe^x$ ,  $e^x$ , and 1 are linearly independent (use Wronskian). It follows that  $\ker(L) = \text{Span}(xe^x, e^x, 1)$ .

## General linear equations

*Definition.* A **linear equation** is an equation of the form

$$L(\mathbf{x}) = \mathbf{b},$$

where  $L : V \rightarrow W$  is a linear mapping,  $\mathbf{b}$  is a given vector from  $W$ , and  $\mathbf{x}$  is an unknown vector from  $V$ .

The range of  $L$  is the set of all vectors  $\mathbf{b} \in W$  such that the equation  $L(\mathbf{x}) = \mathbf{b}$  has a solution.

The kernel of  $L$  is the solution set of the **homogeneous** linear equation  $L(\mathbf{x}) = \mathbf{0}$ .

**Theorem** If the linear equation  $L(\mathbf{x}) = \mathbf{b}$  is solvable and  $\dim \ker L < \infty$ , then the general solution is

$$\mathbf{x}_0 + t_1 \mathbf{v}_1 + \cdots + t_k \mathbf{v}_k,$$

where  $\mathbf{x}_0$  is a particular solution,  $\mathbf{v}_1, \dots, \mathbf{v}_k$  is a basis for the kernel of  $L$ , and  $t_1, \dots, t_k$  are arbitrary scalars.

*Example.* 
$$\begin{cases} x + y + z = 4, \\ x + 2y = 3. \end{cases}$$

$$L: \mathbb{R}^3 \rightarrow \mathbb{R}^2, \quad L \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

Linear equation:  $L(\mathbf{x}) = \mathbf{b}$ , where  $\mathbf{b} = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$ .

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & 4 \\ 1 & 2 & 0 & 3 \end{array} \right) \rightarrow \left( \begin{array}{ccc|c} 1 & 1 & 1 & 4 \\ 0 & 1 & -1 & -1 \end{array} \right) \rightarrow \left( \begin{array}{ccc|c} 1 & 0 & 2 & 5 \\ 0 & 1 & -1 & -1 \end{array} \right)$$

$$\begin{cases} x + 2z = 5 \\ y - z = -1 \end{cases} \iff \begin{cases} x = 5 - 2z \\ y = -1 + z \end{cases}$$

$$(x, y, z) = (5 - 2t, -1 + t, t) = (5, -1, 0) + t(-2, 1, 1).$$



*Example.*  $u'''(x) - 2u''(x) + u'(x) = e^{2x}$ .

Linear operator  $L : C^3(\mathbb{R}) \rightarrow C(\mathbb{R})$ ,

$$Lu = u''' - 2u'' + u'.$$

Linear equation:  $Lu = b$ , where  $b(x) = e^{2x}$ .

We already know that functions  $xe^x$ ,  $e^x$  and 1 form a basis for the kernel of  $L$ . It remains to find a particular solution.

$$L(e^{2x}) = 8e^{2x} - 2(4e^{2x}) + 2e^{2x} = 2e^{2x}.$$

Since  $L$  is a linear operator,  $L\left(\frac{1}{2}e^{2x}\right) = e^{2x}$ .

Particular solution:  $u_0(x) = \frac{1}{2}e^{2x}$ .

Thus the general solution is

$$u(x) = \frac{1}{2}e^{2x} + t_1xe^x + t_2e^x + t_3.$$

## Matrix transformations

Any  $m \times n$  matrix  $A$  gives rise to a transformation  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  given by  $L(\mathbf{x}) = A\mathbf{x}$ , where  $\mathbf{x} \in \mathbb{R}^n$  and  $L(\mathbf{x}) \in \mathbb{R}^m$  are regarded as column vectors. This transformation is **linear**.

*Example.* 
$$L \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 & 0 & 2 \\ 3 & 4 & 7 \\ 0 & 5 & 8 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

Let  $\mathbf{e}_1 = (1, 0, 0)$ ,  $\mathbf{e}_2 = (0, 1, 0)$ ,  $\mathbf{e}_3 = (0, 0, 1)$  be the standard basis for  $\mathbb{R}^3$ . We have that  $L(\mathbf{e}_1) = (1, 3, 0)$ ,  $L(\mathbf{e}_2) = (0, 4, 5)$ ,  $L(\mathbf{e}_3) = (2, 7, 8)$ . Thus  $L(\mathbf{e}_1)$ ,  $L(\mathbf{e}_2)$ ,  $L(\mathbf{e}_3)$  are columns of the matrix.

**Problem.** Find a linear mapping  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  such that  $L(\mathbf{e}_1) = (1, 1)$ ,  $L(\mathbf{e}_2) = (0, -2)$ ,  $L(\mathbf{e}_3) = (3, 0)$ , where  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  is the standard basis for  $\mathbb{R}^3$ .

$$\begin{aligned}L(x, y, z) &= L(x\mathbf{e}_1 + y\mathbf{e}_2 + z\mathbf{e}_3) \\ &= xL(\mathbf{e}_1) + yL(\mathbf{e}_2) + zL(\mathbf{e}_3) \\ &= x(1, 1) + y(0, -2) + z(3, 0) = (x + 3z, x - 2y)\end{aligned}$$

$$L(x, y, z) = \begin{pmatrix} x + 3z \\ x - 2y \end{pmatrix} = \begin{pmatrix} 1 & 0 & 3 \\ 1 & -2 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

Columns of the matrix are vectors  $L(\mathbf{e}_1), L(\mathbf{e}_2), L(\mathbf{e}_3)$ .

**Theorem** Suppose  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a linear map. Then there exists an  $m \times n$  matrix  $A$  such that  $L(\mathbf{x}) = A\mathbf{x}$  for all  $\mathbf{x} \in \mathbb{R}^n$ . Columns of  $A$  are vectors  $L(\mathbf{e}_1), L(\mathbf{e}_2), \dots, L(\mathbf{e}_n)$ , where  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$  is the standard basis for  $\mathbb{R}^n$ .

$$\mathbf{y} = A\mathbf{x} \iff \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

$$\iff \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{pmatrix} = x_1 \begin{pmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{pmatrix} + x_2 \begin{pmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{pmatrix} + \dots + x_n \begin{pmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{pmatrix}$$