

MATH 311

Topics in Applied Mathematics I

**Lecture 23:**

**Basis of eigenvectors.**

**Diagonalization.**

## Eigenvalues and eigenvectors of an operator

*Definition.* Let  $V$  be a vector space and  $L : V \rightarrow V$  be a linear operator. A number  $\lambda$  is called an **eigenvalue** of the operator  $L$  if  $L(\mathbf{v}) = \lambda\mathbf{v}$  for a nonzero vector  $\mathbf{v} \in V$ . The vector  $\mathbf{v}$  is called an **eigenvector** of  $L$  associated with the eigenvalue  $\lambda$ .

The set  $V_\lambda$  of all eigenvectors of  $L$  associated with the eigenvalue  $\lambda$  along with the zero vector is a subspace of  $V$ . It is called the **eigenspace** of  $L$  corresponding to the eigenvalue  $\lambda$ .

## Basis of eigenvectors

Let  $V$  be a finite-dimensional vector space and  $L : V \rightarrow V$  be a linear operator. Let  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  be a basis for  $V$  and  $A$  be the matrix of the operator  $L$  with respect to this basis.

**Theorem** The matrix  $A$  is diagonal if and only if vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  are eigenvectors of  $L$ .

If this is the case, then the diagonal entries of the matrix  $A$  are the corresponding eigenvalues of  $L$ .

$$L(\mathbf{v}_i) = \lambda_i \mathbf{v}_i \iff A = \begin{pmatrix} \lambda_1 & & & 0 \\ & \lambda_2 & & \\ & & \ddots & \\ 0 & & & \lambda_n \end{pmatrix}$$

## How to find a basis of eigenvectors

**Theorem** If  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  are eigenvectors of a linear operator  $L$  associated with distinct eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_k$ , then  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  are linearly independent.

**Corollary 1** Suppose  $\lambda_1, \lambda_2, \dots, \lambda_k$  are all eigenvalues of a linear operator  $L : V \rightarrow V$ . For any  $1 \leq i \leq k$ , let  $S_i$  be a basis for the eigenspace associated to the eigenvalue  $\lambda_i$ . Then these bases are disjoint and the union  $S = S_1 \cup S_2 \cup \dots \cup S_k$  is a linearly independent set.

Moreover, if the vector space  $V$  admits a basis consisting of eigenvectors of  $L$ , then  $S$  is such a basis.

**Corollary 2** Let  $A$  be an  $n \times n$  matrix such that the characteristic equation  $\det(A - \lambda I) = 0$  has  $n$  distinct roots. Then (i) there is a basis for  $\mathbb{R}^n$  consisting of eigenvectors of  $A$ ; (ii) all eigenspaces of  $A$  are one-dimensional.

# Diagonalization

**Theorem 1** Let  $L$  be a linear operator on a finite-dimensional vector space  $V$ . Then the following conditions are equivalent:

- the matrix of  $L$  with respect to some basis is diagonal;
- there exists a basis for  $V$  formed by eigenvectors of  $L$ .

The operator  $L$  is **diagonalizable** if it satisfies these conditions.

**Theorem 2** Let  $A$  be an  $n \times n$  matrix. Then the following conditions are equivalent:

- $A$  is the matrix of a diagonalizable operator;
- $A$  is similar to a diagonal matrix, i.e., it is represented as  $A = UBU^{-1}$ , where the matrix  $B$  is diagonal;
- there exists a basis for  $\mathbb{R}^n$  formed by eigenvectors of  $A$ .

The matrix  $A$  is **diagonalizable** if it satisfies these conditions.

*Example.*  $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$ .

- The matrix  $A$  has two eigenvalues: 1 and 3.
- The eigenspace of  $A$  associated with the eigenvalue 1 is the line spanned by  $\mathbf{v}_1 = (-1, 1)$ .
- The eigenspace of  $A$  associated with the eigenvalue 3 is the line spanned by  $\mathbf{v}_2 = (1, 1)$ .
- Eigenvectors  $\mathbf{v}_1$  and  $\mathbf{v}_2$  form a basis for  $\mathbb{R}^2$ .

Thus the matrix  $A$  is diagonalizable. Namely,  $A = UBU^{-1}$ , where

$$B = \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix}, \quad U = \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Notice that  $U$  is the transition matrix from the basis  $\mathbf{v}_1, \mathbf{v}_2$  to the standard basis.

Example.  $A = \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix}$ .

- The matrix  $A$  has two eigenvalues: 0 and 2.
- The eigenspace for 0 is one-dimensional; it has a basis  $S_1 = \{\mathbf{v}_1\}$ , where  $\mathbf{v}_1 = (-1, 1, 0)$ .
- The eigenspace for 2 is two-dimensional; it has a basis  $S_2 = \{\mathbf{v}_2, \mathbf{v}_3\}$ , where  $\mathbf{v}_2 = (1, 1, 0)$ ,  $\mathbf{v}_3 = (-1, 0, 1)$ .
- The union  $S_1 \cup S_2 = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is a linearly independent set, hence it is a basis for  $\mathbb{R}^3$ .

Thus the matrix  $A$  is diagonalizable. Namely,  $A = UBU^{-1}$ , where

$$B = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}, \quad U = \begin{pmatrix} -1 & 1 & -1 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

To *diagonalize* an  $n \times n$  matrix  $A$  is to find a diagonal matrix  $B$  and an invertible matrix  $U$  such that  $A = UBU^{-1}$ .

Suppose there exists a basis  $\mathbf{v}_1, \dots, \mathbf{v}_n$  for  $\mathbb{R}^n$  consisting of eigenvectors of  $A$ . That is,  $A\mathbf{v}_k = \lambda_k\mathbf{v}_k$ , where  $\lambda_k \in \mathbb{R}$ .

Then  $A = UBU^{-1}$ , where  $B = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$  and  $U$  is a transition matrix whose columns are vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ .

*Example.*  $A = \begin{pmatrix} 4 & 3 \\ 0 & 1 \end{pmatrix}$ .  $\det(A - \lambda I) = (4 - \lambda)(1 - \lambda)$ .

Eigenvalues:  $\lambda_1 = 4$ ,  $\lambda_2 = 1$ .

Associated eigenvectors:  $\mathbf{v}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ,  $\mathbf{v}_2 = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$ .

Thus  $A = UBU^{-1}$ , where

$$B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$



Suppose we have a problem that involves a square matrix  $A$  in the context of matrix multiplication.

Also, suppose that the case when  $A$  is a diagonal matrix is simple. Then the diagonalization may help in solving this problem (or may not). Namely, it may reduce the case of a diagonalizable matrix to that of a diagonal one.

An example of such a problem is, given a square matrix  $A$ , to find its power  $A^k$ :

$$A = \begin{pmatrix} s_1 & & & 0 \\ & s_2 & & \\ & & \ddots & \\ 0 & & & s_n \end{pmatrix} \implies A^k = \begin{pmatrix} s_1^k & & & 0 \\ & s_2^k & & \\ & & \ddots & \\ 0 & & & s_n^k \end{pmatrix}$$

**Problem.** Let  $A = \begin{pmatrix} 4 & 3 \\ 0 & 1 \end{pmatrix}$ . Find  $A^5$ .

We know that  $A = UBU^{-1}$ , where

$$B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Then  $A^5 = UBU^{-1}UBU^{-1}UBU^{-1}UBU^{-1}UBU^{-1}$

$$= UB^5U^{-1} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1024 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 1024 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1024 & 1023 \\ 0 & 1 \end{pmatrix}.$$

**Problem.** Let  $A = \begin{pmatrix} 4 & 3 \\ 0 & 1 \end{pmatrix}$ . Find  $A^k$  ( $k \geq 1$ ).

We know that  $A = UBU^{-1}$ , where

$$B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Then

$$\begin{aligned} A^k &= U B^k U^{-1} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 4^k & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 4^k & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 4^k & 4^k - 1 \\ 0 & 1 \end{pmatrix}. \end{aligned}$$

**Problem.** Let  $A = \begin{pmatrix} 4 & 3 \\ 0 & 1 \end{pmatrix}$ . Find a matrix  $C$  such that  $C^2 = A$ .

We know that  $A = UBU^{-1}$ , where

$$B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Suppose that  $D^2 = B$  for some matrix  $D$ . Let  $C = UDU^{-1}$ . Then  $C^2 = UDU^{-1}UDU^{-1} = UD^2U^{-1} = UBU^{-1} = A$ .

We can take  $D = \begin{pmatrix} \sqrt{4} & 0 \\ 0 & \sqrt{1} \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$ .

Then  $C = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$ .

Initial value problem for a system of linear ODEs:

$$\begin{cases} \frac{dx}{dt} = 4x + 3y, \\ \frac{dy}{dt} = y, \end{cases} \quad x(0) = 1, \quad y(0) = 1.$$

The system can be rewritten in vector form:

$$\frac{d\mathbf{v}}{dt} = A\mathbf{v}, \quad \text{where } A = \begin{pmatrix} 4 & 3 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} x \\ y \end{pmatrix}.$$

Matrix  $A$  is diagonalizable:  $A = UBU^{-1}$ , where

$$B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Let  $\mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$  be coordinates of the vector  $\mathbf{v}$  relative to the basis  $\mathbf{v}_1 = (1, 0)$ ,  $\mathbf{v}_2 = (-1, 1)$  of eigenvectors of  $A$ . Then  $\mathbf{v} = U\mathbf{w} \implies \mathbf{w} = U^{-1}\mathbf{v}$ .

It follows that

$$\frac{d\mathbf{w}}{dt} = \frac{d}{dt}(U^{-1}\mathbf{v}) = U^{-1}\frac{d\mathbf{v}}{dt} = U^{-1}A\mathbf{v} = U^{-1}AU\mathbf{w}.$$

$$\text{Hence } \frac{d\mathbf{w}}{dt} = B\mathbf{w} \iff \begin{cases} \frac{dw_1}{dt} = 4w_1, \\ \frac{dw_2}{dt} = w_2. \end{cases}$$

General solution:  $w_1(t) = c_1 e^{4t}$ ,  $w_2(t) = c_2 e^t$ , where  $c_1, c_2 \in \mathbb{R}$ .

Initial condition:

$$\mathbf{w}(0) = U^{-1}\mathbf{v}(0) = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}.$$

Thus  $w_1(t) = 2e^{4t}$ ,  $w_2(t) = e^t$ . Then

$$\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = U\mathbf{w}(t) = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2e^{4t} \\ e^t \end{pmatrix} = \begin{pmatrix} 2e^{4t} - e^t \\ e^t \end{pmatrix}.$$

There are **two obstructions** to existence of a basis consisting of eigenvectors. They are illustrated by the following examples.

*Example 1.*  $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ .

$\det(A - \lambda I) = (\lambda - 1)^2$ . Hence  $\lambda = 1$  is the only eigenvalue. The associated eigenspace is the line  $t(1, 0)$ .

*Example 2.*  $A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ .

$\det(A - \lambda I) = \lambda^2 + 1$ .

$\implies$  no real eigenvalues or eigenvectors

(However there are *complex* eigenvalues/eigenvectors.)