

MATH 311

Topics in Applied Mathematics I

Lecture 29:

Orthogonality in inner product spaces.

Norm

The notion of *norm* generalizes the notion of length of a vector in \mathbb{R}^n .

Definition. Let V be a vector space. A function $\alpha : V \rightarrow \mathbb{R}$, usually denoted $\alpha(\mathbf{x}) = \|\mathbf{x}\|$, is called a **norm** on V if it has the following properties:

- (i) $\|\mathbf{x}\| \geq 0$, $\|\mathbf{x}\| = 0$ only for $\mathbf{x} = \mathbf{0}$ (positivity)
- (ii) $\|r\mathbf{x}\| = |r| \|\mathbf{x}\|$ for all $r \in \mathbb{R}$ (homogeneity)
- (iii) $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$ (triangle inequality)

A **normed vector space** is a vector space endowed with a norm. The norm defines a distance function on the normed vector space: $\text{dist}(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|$.

Examples. $V = \mathbb{R}^n$, $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$.

- $\|\mathbf{x}\|_\infty = \max(|x_1|, |x_2|, \dots, |x_n|)$.
- $\|\mathbf{x}\|_p = (|x_1|^p + |x_2|^p + \dots + |x_n|^p)^{1/p}$, $p \geq 1$.

Examples. $V = C[a, b]$, $f : [a, b] \rightarrow \mathbb{R}$.

- $\|f\|_\infty = \max_{a \leq x \leq b} |f(x)|$.
- $\|f\|_p = \left(\int_a^b |f(x)|^p dx \right)^{1/p}$, $p \geq 1$.

Inner product

The notion of *inner product* generalizes the notion of dot product of vectors in \mathbb{R}^n .

Definition. Let V be a vector space. A function $\beta : V \times V \rightarrow \mathbb{R}$, usually denoted $\beta(\mathbf{x}, \mathbf{y}) = \langle \mathbf{x}, \mathbf{y} \rangle$, is called an **inner product** on V if it is positive, symmetric, and bilinear. That is, if

- (i) $\langle \mathbf{x}, \mathbf{x} \rangle \geq 0$, $\langle \mathbf{x}, \mathbf{x} \rangle = 0$ only for $\mathbf{x} = \mathbf{0}$ (positivity)
- (ii) $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$ (symmetry)
- (iii) $\langle r\mathbf{x}, \mathbf{y} \rangle = r\langle \mathbf{x}, \mathbf{y} \rangle$ (homogeneity)
- (iv) $\langle \mathbf{x} + \mathbf{y}, \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{z} \rangle + \langle \mathbf{y}, \mathbf{z} \rangle$ (distributive law)

An **inner product space** is a vector space endowed with an inner product.

Examples. $V = \mathbb{R}^n$.

- $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x} \cdot \mathbf{y} = x_1y_1 + x_2y_2 + \cdots + x_ny_n$.

- $\langle \mathbf{x}, \mathbf{y} \rangle = d_1x_1y_1 + d_2x_2y_2 + \cdots + d_nx_ny_n$,

where $d_1, d_2, \dots, d_n > 0$.

Examples. $V = C[a, b]$.

- $\langle f, g \rangle = \int_a^b f(x)g(x) dx$.

- $\langle f, g \rangle = \int_a^b f(x)g(x)w(x) dx$,

where w is bounded, piecewise continuous, and $w > 0$ everywhere on $[a, b]$.

Theorem Suppose $\langle \mathbf{x}, \mathbf{y} \rangle$ is an inner product on a vector space V . Then

$$\langle \mathbf{x}, \mathbf{y} \rangle^2 \leq \langle \mathbf{x}, \mathbf{x} \rangle \langle \mathbf{y}, \mathbf{y} \rangle \quad \text{for all } \mathbf{x}, \mathbf{y} \in V.$$

Proof: For any $t \in \mathbb{R}$ let $\mathbf{v}_t = \mathbf{x} + t\mathbf{y}$. Then

$$\begin{aligned} \langle \mathbf{v}_t, \mathbf{v}_t \rangle &= \langle \mathbf{x} + t\mathbf{y}, \mathbf{x} + t\mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{x} + t\mathbf{y} \rangle + t\langle \mathbf{y}, \mathbf{x} + t\mathbf{y} \rangle \\ &= \langle \mathbf{x}, \mathbf{x} \rangle + t\langle \mathbf{x}, \mathbf{y} \rangle + t\langle \mathbf{y}, \mathbf{x} \rangle + t^2\langle \mathbf{y}, \mathbf{y} \rangle. \end{aligned}$$

Assume that $\mathbf{y} \neq \mathbf{0}$ and let $t = -\frac{\langle \mathbf{x}, \mathbf{y} \rangle}{\langle \mathbf{y}, \mathbf{y} \rangle}$. Then

$$\langle \mathbf{v}_t, \mathbf{v}_t \rangle = \langle \mathbf{x}, \mathbf{x} \rangle + t\langle \mathbf{y}, \mathbf{x} \rangle = \langle \mathbf{x}, \mathbf{x} \rangle - \frac{\langle \mathbf{x}, \mathbf{y} \rangle^2}{\langle \mathbf{y}, \mathbf{y} \rangle}.$$

Since $\langle \mathbf{v}_t, \mathbf{v}_t \rangle \geq 0$, the desired inequality follows. In the case $\mathbf{y} = \mathbf{0}$, we have $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{y} \rangle = 0$.

Cauchy-Schwarz Inequality:

$$|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} \sqrt{\langle \mathbf{y}, \mathbf{y} \rangle}.$$

Corollary 1 $|\mathbf{x} \cdot \mathbf{y}| \leq \|\mathbf{x}\| \|\mathbf{y}\|$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$.

Equivalently, for all $x_i, y_i \in \mathbb{R}$,

$$(x_1y_1 + \cdots + x_ny_n)^2 \leq (x_1^2 + \cdots + x_n^2)(y_1^2 + \cdots + y_n^2).$$

Corollary 2 For any $f, g \in C[a, b]$,

$$\left(\int_a^b f(x)g(x) dx \right)^2 \leq \int_a^b |f(x)|^2 dx \cdot \int_a^b |g(x)|^2 dx.$$

Norms induced by inner products

Theorem Suppose $\langle \mathbf{x}, \mathbf{y} \rangle$ is an inner product on a vector space V . Then $\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}$ is a norm.

Proof: Positivity is obvious. Homogeneity:

$$\|r\mathbf{x}\| = \sqrt{\langle r\mathbf{x}, r\mathbf{x} \rangle} = \sqrt{r^2 \langle \mathbf{x}, \mathbf{x} \rangle} = |r| \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}.$$

Triangle inequality (follows from Cauchy-Schwarz's):

$$\begin{aligned} \|\mathbf{x} + \mathbf{y}\|^2 &= \langle \mathbf{x} + \mathbf{y}, \mathbf{x} + \mathbf{y} \rangle \\ &= \langle \mathbf{x}, \mathbf{x} \rangle + \langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{y}, \mathbf{x} \rangle + \langle \mathbf{y}, \mathbf{y} \rangle \\ &\leq \langle \mathbf{x}, \mathbf{x} \rangle + |\langle \mathbf{x}, \mathbf{y} \rangle| + |\langle \mathbf{y}, \mathbf{x} \rangle| + \langle \mathbf{y}, \mathbf{y} \rangle \\ &\leq \|\mathbf{x}\|^2 + 2\|\mathbf{x}\| \|\mathbf{y}\| + \|\mathbf{y}\|^2 = (\|\mathbf{x}\| + \|\mathbf{y}\|)^2. \end{aligned}$$

Examples. • The length of a vector in \mathbb{R}^n ,

$$|\mathbf{x}| = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2},$$

is the norm induced by the dot product

$$\mathbf{x} \cdot \mathbf{y} = x_1y_1 + x_2y_2 + \cdots + x_ny_n.$$

• The norm $\|f\|_2 = \left(\int_a^b |f(x)|^2 dx \right)^{1/2}$ on the vector space $C[a, b]$ is induced by the inner product

$$\langle f, g \rangle = \int_a^b f(x)g(x) dx.$$

Angle

Let V be an inner product space with an inner product $\langle \cdot, \cdot \rangle$ and the induced norm $\| \cdot \|$. Then

$$|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \| \mathbf{x} \| \| \mathbf{y} \|$$

for all $\mathbf{x}, \mathbf{y} \in V$ (the Cauchy-Schwarz inequality).

Therefore we can define the **angle** between nonzero vectors in V by

$$\angle(\mathbf{x}, \mathbf{y}) = \arccos \frac{\langle \mathbf{x}, \mathbf{y} \rangle}{\| \mathbf{x} \| \| \mathbf{y} \|}.$$

Then $\langle \mathbf{x}, \mathbf{y} \rangle = \| \mathbf{x} \| \| \mathbf{y} \| \cos \angle(\mathbf{x}, \mathbf{y})$.

In particular, vectors \mathbf{x} and \mathbf{y} are **orthogonal** (denoted $\mathbf{x} \perp \mathbf{y}$) if $\langle \mathbf{x}, \mathbf{y} \rangle = 0$.

Orthogonal sets

Let V be an inner product space with an inner product $\langle \cdot, \cdot \rangle$ and the induced norm $\| \cdot \|$.

Definition. A nonempty set $S \subset V$ of nonzero vectors is called an **orthogonal set** if all vectors in S are mutually orthogonal. That is, $\mathbf{0} \notin S$ and $\langle \mathbf{x}, \mathbf{y} \rangle = 0$ for any $\mathbf{x}, \mathbf{y} \in S$, $\mathbf{x} \neq \mathbf{y}$.

An orthogonal set $S \subset V$ is called **orthonormal** if $\|\mathbf{x}\| = 1$ for any $\mathbf{x} \in S$.

Remark. Vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in V$ form an orthonormal set if and only if

$$\langle \mathbf{v}_i, \mathbf{v}_j \rangle = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$$

Example

- $V = C[-\pi, \pi]$, $\langle f, g \rangle = \int_{-\pi}^{\pi} f(x)g(x) dx$.

$$f_1(x) = \sin x, \quad f_2(x) = \sin 2x, \quad \dots, \quad f_n(x) = \sin nx, \quad \dots$$

$$\langle f_m, f_n \rangle = \int_{-\pi}^{\pi} \sin(mx) \sin(nx) dx = \begin{cases} \pi & \text{if } m = n, \\ 0 & \text{if } m \neq n. \end{cases}$$

Thus the set $\{f_1, f_2, f_3, \dots\}$ is orthogonal but not orthonormal.

It is orthonormal with respect to a scaled inner product

$$\langle\langle f, g \rangle\rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x)g(x) dx.$$

Orthogonality \implies linear independence

Theorem Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are nonzero vectors that form an orthogonal set. Then $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly independent.

Proof: Suppose $t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k = \mathbf{0}$ for some $t_1, t_2, \dots, t_k \in \mathbb{R}$.

Then for any index $1 \leq i \leq k$ we have

$$\langle t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k, \mathbf{v}_i \rangle = \langle \mathbf{0}, \mathbf{v}_i \rangle = 0.$$

$$\implies t_1\langle \mathbf{v}_1, \mathbf{v}_i \rangle + t_2\langle \mathbf{v}_2, \mathbf{v}_i \rangle + \dots + t_k\langle \mathbf{v}_k, \mathbf{v}_i \rangle = 0$$

By orthogonality, $t_i\langle \mathbf{v}_i, \mathbf{v}_i \rangle = 0 \implies t_i = 0$.

Orthonormal basis

Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is an orthonormal basis for an inner product space V .

Theorem 1 Let $\mathbf{x} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \dots + x_n\mathbf{v}_n$ and $\mathbf{y} = y_1\mathbf{v}_1 + y_2\mathbf{v}_2 + \dots + y_n\mathbf{v}_n$, where $x_i, y_j \in \mathbb{R}$.

Then

$$(i) \quad \langle \mathbf{x}, \mathbf{y} \rangle = x_1y_1 + x_2y_2 + \dots + x_ny_n,$$

$$(ii) \quad \|\mathbf{x}\| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}.$$


Theorem 2 For any vector $\mathbf{x} \in V$,

$$\mathbf{x} = \langle \mathbf{x}, \mathbf{v}_1 \rangle \mathbf{v}_1 + \langle \mathbf{x}, \mathbf{v}_2 \rangle \mathbf{v}_2 + \dots + \langle \mathbf{x}, \mathbf{v}_n \rangle \mathbf{v}_n.$$

Orthogonal projection

Theorem Let V be an inner product space and V_0 be a finite-dimensional subspace of V . Then any vector $\mathbf{x} \in V$ is uniquely represented as $\mathbf{x} = \mathbf{p} + \mathbf{o}$, where $\mathbf{p} \in V_0$ and $\mathbf{o} \perp V_0$.

The component \mathbf{p} is called the **orthogonal projection** of the vector \mathbf{x} onto the subspace V_0 .



The projection \mathbf{p} is closer to \mathbf{x} than any other vector in V_0 . Hence the distance from \mathbf{x} to V_0 is $\|\mathbf{x} - \mathbf{p}\| = \|\mathbf{o}\|$.

Theorem Let V be an inner product space and V_0 be a finite-dimensional subspace of V . Then any vector $\mathbf{x} \in V$ is uniquely represented as $\mathbf{x} = \mathbf{p} + \mathbf{o}$, where $\mathbf{p} \in V_0$ and $\mathbf{o} \perp V_0$.

Theorem Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is an orthogonal basis for the subspace V_0 . Then for any vector $\mathbf{x} \in V$ the orthogonal projection \mathbf{p} onto V_0 is given by

$$\mathbf{p} = \frac{\langle \mathbf{x}, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 + \frac{\langle \mathbf{x}, \mathbf{v}_2 \rangle}{\langle \mathbf{v}_2, \mathbf{v}_2 \rangle} \mathbf{v}_2 + \dots + \frac{\langle \mathbf{x}, \mathbf{v}_n \rangle}{\langle \mathbf{v}_n, \mathbf{v}_n \rangle} \mathbf{v}_n.$$

The Gram-Schmidt orthogonalization process

Let V be a vector space with an inner product. Suppose $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ is a basis for V . Let

$$\mathbf{v}_1 = \mathbf{x}_1,$$

$$\mathbf{v}_2 = \mathbf{x}_2 - \frac{\langle \mathbf{x}_2, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1,$$

$$\mathbf{v}_3 = \mathbf{x}_3 - \frac{\langle \mathbf{x}_3, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 - \frac{\langle \mathbf{x}_3, \mathbf{v}_2 \rangle}{\langle \mathbf{v}_2, \mathbf{v}_2 \rangle} \mathbf{v}_2,$$

.....

$$\mathbf{v}_n = \mathbf{x}_n - \frac{\langle \mathbf{x}_n, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 - \dots - \frac{\langle \mathbf{x}_n, \mathbf{v}_{n-1} \rangle}{\langle \mathbf{v}_{n-1}, \mathbf{v}_{n-1} \rangle} \mathbf{v}_{n-1}.$$

Then $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is an orthogonal basis for V .

Normalization

Let V be a vector space with an inner product.

Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is an orthogonal basis for V .

Let $\mathbf{w}_1 = \frac{\mathbf{v}_1}{\|\mathbf{v}_1\|}$, $\mathbf{w}_2 = \frac{\mathbf{v}_2}{\|\mathbf{v}_2\|}$, \dots , $\mathbf{w}_n = \frac{\mathbf{v}_n}{\|\mathbf{v}_n\|}$.

Then $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n$ is an orthonormal basis for V .

Theorem Any finite-dimensional vector space with an inner product has an orthonormal basis.

Remark. An infinite-dimensional vector space with an inner product may or may not have an orthonormal basis.

Problem. Approximate the function $f(x) = e^x$ on the interval $[-1, 1]$ by a quadratic polynomial.

The best approximation would be a polynomial $p(x)$ that minimizes the distance relative to the uniform norm:

$$\|f - p\|_{\infty} = \max_{|x| \leq 1} |f(x) - p(x)|.$$

However there is no analytic way to find such a polynomial. Instead, one can find a **“least squares”** approximation that minimizes the integral norm

$$\|f - p\|_2 = \left(\int_{-1}^1 |f(x) - p(x)|^2 dx \right)^{1/2}.$$

The norm $\| \cdot \|_2$ is induced by the inner product

$$\langle g, h \rangle = \int_{-1}^1 g(x)h(x) dx.$$

Therefore $\|f - p\|_2$ is minimal if p is the orthogonal projection of the function f on the subspace \mathcal{P}_3 of quadratic polynomials.

We should apply the Gram-Schmidt process to the polynomials $1, x, x^2$, which form a basis for \mathcal{P}_3 .

This would yield an orthogonal basis p_0, p_1, p_2 .

Then

$$p(x) = \frac{\langle f, p_0 \rangle}{\langle p_0, p_0 \rangle} p_0(x) + \frac{\langle f, p_1 \rangle}{\langle p_1, p_1 \rangle} p_1(x) + \frac{\langle f, p_2 \rangle}{\langle p_2, p_2 \rangle} p_2(x).$$

Fourier series: view from linear algebra

Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n, \dots$ are nonzero vectors in an inner product space V that form an orthogonal set S . Given $\mathbf{x} \in V$, the **Fourier series** of the vector \mathbf{x} relative to the orthogonal set S is a series

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_n\mathbf{v}_n + \cdots, \quad \text{where } c_i = \frac{\langle \mathbf{x}, \mathbf{v}_i \rangle}{\langle \mathbf{v}_i, \mathbf{v}_i \rangle}.$$

The numbers c_1, c_2, \dots are called the **Fourier coefficients** of \mathbf{x} relative to S .

By construction, a partial sum $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_n\mathbf{v}_n$ of the Fourier series is the orthogonal projection of the vector \mathbf{x} onto the subspace $\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n)$.

Classical Fourier series

Consider a functional vector space $V = C[-\pi, \pi]$ with the standard inner product $\langle f, g \rangle = \int_{-\pi}^{\pi} f(x)g(x) dx$.

Then the functions $1, \sin x, \cos x, \sin 2x, \cos 2x, \dots$ form an orthogonal set in the inner product space V . This gives rise to the classical Fourier series of a function $F \in C[-\pi, \pi]$:

$$a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx,$$

where

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} F(x) dx$$

and for $n \geq 1$,

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} F(x) \cos nx dx, \quad b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} F(x) \sin nx dx.$$

Convergence of Fourier series

Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n, \dots$ are vectors in an inner product space V that form an orthogonal set S . The set S is called a **Hilbert basis** for V if any vector $\mathbf{x} \in V$ can be expanded into a series $\mathbf{x} = \sum_{n=1}^{\infty} \alpha_n \mathbf{v}_n$, where α_n are some scalars.

Theorem 1 If S is a Hilbert basis for V , then the above expansion is unique for any vector $\mathbf{x} \in V$. Namely, it coincides with the Fourier series of \mathbf{x} relative to S .

Theorem 2 The functions $1, \sin x, \cos x, \sin 2x, \cos 2x, \dots$ form a Hilbert basis for the space $C[-\pi, \pi]$.

As a consequence, Fourier series of a continuous function on $[-\pi, \pi]$ converges to this function with respect to the distance

$$\text{dist}(f, g) = \|f - g\| = \left(\int_{-\pi}^{\pi} |f(x) - g(x)|^2 dx \right)^{1/2}.$$

Note that this need not imply pointwise convergence.