

MATH 311

Topics in Applied Mathematics I

**Lecture 7:**

**Inverse matrix (continued).**

## Inverse matrix

*Definition.* Let  $A$  be an  $n \times n$  matrix. The **inverse** of  $A$  is an  $n \times n$  matrix, denoted  $A^{-1}$ , such that

$$\boxed{AA^{-1} = A^{-1}A = I.}$$

If  $A^{-1}$  exists then the matrix  $A$  is called **invertible**. Otherwise  $A$  is called **singular**.

## Inverting diagonal matrices

**Theorem** A diagonal matrix  $D = \text{diag}(d_1, \dots, d_n)$  is invertible if and only if all diagonal entries are nonzero:  $d_i \neq 0$  for  $1 \leq i \leq n$ .

If  $D$  is invertible then  $D^{-1} = \text{diag}(d_1^{-1}, \dots, d_n^{-1})$ .

$$\begin{pmatrix} d_1 & 0 & \dots & 0 \\ 0 & d_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & d_n \end{pmatrix}^{-1} = \begin{pmatrix} d_1^{-1} & 0 & \dots & 0 \\ 0 & d_2^{-1} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & d_n^{-1} \end{pmatrix}$$

## Inverting $2 \times 2$ matrices

*Definition.* The **determinant** of a  $2 \times 2$  matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \text{ is } \det A = ad - bc.$$

**Theorem** A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  is invertible if and only if  $\det A \neq 0$ .

If  $\det A \neq 0$  then

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

**Problem.** Solve a system  $\begin{cases} 4x + 3y = 5, \\ 3x + 2y = -1. \end{cases}$

This system is equivalent to a matrix equation  $A\mathbf{x} = \mathbf{b}$ ,  
where  $A = \begin{pmatrix} 4 & 3 \\ 3 & 2 \end{pmatrix}$ ,  $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$ ,  $\mathbf{b} = \begin{pmatrix} 5 \\ -1 \end{pmatrix}$ .

We have  $\det A = -1 \neq 0$ . Hence  $A$  is invertible.

$$\begin{aligned} A\mathbf{x} = \mathbf{b} &\implies A^{-1}(A\mathbf{x}) = A^{-1}\mathbf{b} \implies (A^{-1}A)\mathbf{x} = A^{-1}\mathbf{b} \\ &\implies \mathbf{x} = A^{-1}\mathbf{b}. \end{aligned}$$

Conversely,  $\mathbf{x} = A^{-1}\mathbf{b} \implies A\mathbf{x} = A(A^{-1}\mathbf{b}) = (AA^{-1})\mathbf{b} = \mathbf{b}$ .

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4 & 3 \\ 3 & 2 \end{pmatrix}^{-1} \begin{pmatrix} 5 \\ -1 \end{pmatrix} = \frac{1}{-1} \begin{pmatrix} 2 & -3 \\ -3 & 4 \end{pmatrix} \begin{pmatrix} 5 \\ -1 \end{pmatrix} = \begin{pmatrix} -13 \\ 19 \end{pmatrix}$$

System of  $n$  linear equations in  $n$  variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \dots\dots\dots \\ a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n = b_n \end{cases} \iff \mathbf{Ax} = \mathbf{b},$$

where

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix}, \quad \mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}.$$

**Theorem** If the matrix  $A$  is invertible then the system has a unique solution, which is  $\mathbf{x} = A^{-1}\mathbf{b}$ .

## General results on inverse matrices

**Theorem 1** Given an  $n \times n$  matrix  $A$ , the following conditions are equivalent:

- (i)  $A$  is invertible;
- (ii)  $\mathbf{x} = \mathbf{0}$  is the only solution of the matrix equation  $A\mathbf{x} = \mathbf{0}$ ;
- (iii) the matrix equation  $A\mathbf{x} = \mathbf{b}$  has a unique solution for any  $n$ -dimensional column vector  $\mathbf{b}$ ;
- (iv) the row echelon form of  $A$  has no zero rows;
- (v) the reduced row echelon form of  $A$  is the identity matrix.

**Theorem 2** Suppose that a sequence of elementary row operations converts a matrix  $A$  into the identity matrix.

Then the same sequence of operations converts the identity matrix into the inverse matrix  $A^{-1}$ .

*Row echelon form of a square matrix:*

A 7x7 matrix in row echelon form. The leading entries are represented by squares in the main diagonal. A blue staircase path starts from the top-left square and moves down and right, ending at the bottom-right square. All other entries are represented by asterisks.

invertible case

A 7x7 matrix in row echelon form, similar to the first one. However, the asterisks in the third row, fourth column and the sixth row, seventh column are circled in red. A blue staircase path is also shown, ending at the bottom-right square.

noninvertible case

For any matrix in row echelon form, the number of columns with leading entries equals the number of rows with leading entries. For a square matrix, also the number of columns *without* leading entries (i.e., the number of free variables in a related system of linear equations) equals the number of rows *without* leading entries (i.e., zero rows).



*Row echelon form of a square matrix:*

A 7x7 matrix in row echelon form. The main diagonal elements are represented by empty boxes. All elements below the diagonal are zero. All elements above the diagonal are represented by asterisks. A blue line traces the path of the leading ones from the top-left to the bottom-right.

invertible case

A 7x7 matrix in row echelon form. The main diagonal elements are represented by empty boxes. All elements below the diagonal are zero. All elements above the diagonal are represented by asterisks. Two asterisks are circled in red: one in the third row, fourth column, and one in the seventh row, seventh column. A blue line traces the path of the leading ones from the top-left to the bottom-right.

noninvertible case

Hence the row echelon form of a square matrix  $A$  is either strict triangular or else it has a zero row. In the former case, the equation  $A\mathbf{x} = \mathbf{b}$  *always* has a unique solution. In the latter case,  $A\mathbf{x} = \mathbf{b}$  *never* has a unique solution. Also, in the former case the reduced row echelon form of  $A$  is  $I$ .

*Example.*  $A = \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}.$

*To check whether  $A$  is invertible, we convert it to row echelon form.*

Interchange the 1st row with the 2nd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 3 & -2 & 0 \\ -2 & 3 & 0 \end{pmatrix}$$

Add  $-3$  times the 1st row to the 2nd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & -2 & -3 \\ -2 & 3 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & -2 & -3 \\ -2 & 3 & 0 \end{pmatrix}$$

Add 2 times the 1st row to the 3rd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & -2 & -3 \\ 0 & 3 & 2 \end{pmatrix}$$

Multiply the 2nd row by  $-0.5$ :

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 3 & 2 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 3 & 2 \end{pmatrix}$$

Add  $-3$  times the 2nd row to the 3rd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 0 & -2.5 \end{pmatrix}$$

Multiply the 3rd row by  $-0.4$ :

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} \boxed{1} & 0 & 1 \\ 0 & \boxed{1} & 1.5 \\ 0 & 0 & \boxed{1} \end{pmatrix}$$

*We already know that the matrix  $A$  is invertible.*

*Let's proceed towards reduced row echelon form.*

Add  $-1.5$  times the 3rd row to the 2nd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Add  $-1$  times the 3rd row to the 1st row:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

To obtain  $A^{-1}$ , we need to apply the following sequence of elementary row operations to the identity matrix:

- interchange the 1st row with the 2nd row,
- add  $-3$  times the 1st row to the 2nd row,
- add 2 times the 1st row to the 3rd row,
- multiply the 2nd row by  $-0.5$ ,
- add  $-3$  times the 2nd row to the 3rd row,
- multiply the 3rd row by  $-0.4$ ,
- add  $-1.5$  times the 3rd row to the 2nd row,
- add  $-1$  times the 3rd row to the 1st row.

A convenient way to compute the inverse matrix  $A^{-1}$  is to merge the matrices  $A$  and  $I$  into one  $3 \times 6$  matrix  $(A | I)$ , and apply elementary row operations to this new matrix.

$$A = \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$(A | I) = \left( \begin{array}{ccc|ccc} 3 & -2 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

$$\left( \begin{array}{ccc|ccc} 3 & -2 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

Interchange the 1st row with the 2nd row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 3 & -2 & 0 & 1 & 0 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

Add  $-3$  times the 1st row to the 2nd row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & -3 & 1 & -3 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$



$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & -3 & 1 & -3 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

Add 2 times the 1st row to the 3rd row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & -3 & 1 & -3 & 0 \\ 0 & 3 & 2 & 0 & 2 & 1 \end{array} \right)$$

Multiply the 2nd row by  $-0.5$ :

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 3 & 2 & 0 & 2 & 1 \end{array} \right)$$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 3 & 2 & 0 & 2 & 1 \end{array} \right)$$

Add  $-3$  times the 2nd row to the 3rd row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 0 & -2.5 & 1.5 & -2.5 & 1 \end{array} \right)$$

Multiply the 3rd row by  $-0.4$ :

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right)$$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right)$$

Add  $-1.5$  times the 3rd row to the 2nd row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0.4 & 0 & 0.6 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right)$$

Add  $-1$  times the 3rd row to the 1st row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 0 & 0.6 & 0 & 0.4 \\ 0 & 1 & 0 & 0.4 & 0 & 0.6 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right) = (I | A^{-1})$$

Thus 
$$\begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}^{-1} = \begin{pmatrix} \frac{3}{5} & 0 & \frac{2}{5} \\ \frac{2}{5} & 0 & \frac{3}{5} \\ -\frac{3}{5} & 1 & -\frac{2}{5} \end{pmatrix}.$$

That is,

$$\begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix} \begin{pmatrix} \frac{3}{5} & 0 & \frac{2}{5} \\ \frac{2}{5} & 0 & \frac{3}{5} \\ -\frac{3}{5} & 1 & -\frac{2}{5} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$$\begin{pmatrix} \frac{3}{5} & 0 & \frac{2}{5} \\ \frac{2}{5} & 0 & \frac{3}{5} \\ -\frac{3}{5} & 1 & -\frac{2}{5} \end{pmatrix} \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

## Why does it work?

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} = \begin{pmatrix} a_1 & a_2 & a_3 \\ 2b_1 & 2b_2 & 2b_3 \\ c_1 & c_2 & c_3 \end{pmatrix},$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1+3a_1 & b_2+3a_2 & b_3+3a_3 \\ c_1 & c_2 & c_3 \end{pmatrix},$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} = \begin{pmatrix} a_1 & a_2 & a_3 \\ c_1 & c_2 & c_3 \\ b_1 & b_2 & b_3 \end{pmatrix}.$$

**Proposition** Any elementary row operation can be simulated as left multiplication by a certain matrix.



## Elementary matrices

$$E = \begin{pmatrix} 1 & & & & & & & \\ \vdots & \ddots & & & & & & \\ 0 & \cdots & 1 & & & & & \\ \vdots & & \vdots & \ddots & & & & \\ 0 & \cdots & r & \cdots & 1 & & & \\ \vdots & & \vdots & & \vdots & \ddots & & \\ 0 & \cdots & 0 & \cdots & 0 & \cdots & 1 & \end{pmatrix} \begin{array}{l} \text{row \#}i \\ \\ \text{row \#}j \end{array}$$

To obtain the matrix  $EA$  from  $A$ , add  $r$  times the  $i$ th row to the  $j$ th row. To obtain the matrix  $AE$  from  $A$ , add  $r$  times the  $j$ th column to the  $i$ th column.





## Why does it work? (continued)

Assume that a square matrix  $A$  can be converted to the identity matrix by a sequence of elementary row operations. Then  $E_k E_{k-1} \dots E_2 E_1 A = I$ , where  $E_1, E_2, \dots, E_k$  are elementary matrices simulating those operations.

Applying the same sequence of operations to the identity matrix, we obtain the matrix

$$B = E_k E_{k-1} \dots E_2 E_1 I = E_k E_{k-1} \dots E_2 E_1.$$

Thus  $BA = I$ . Besides,  $B$  is invertible since elementary matrices are invertible (why?). It follows that  $A = B^{-1}$ , then  $B = A^{-1}$ .