

MATH 311

Topics in Applied Mathematics I

Lecture 22:
Eigenvalues and eigenvectors
of a linear operator.

Eigenvalues and eigenvectors of a matrix

Definition. Let A be an $n \times n$ matrix. A number $\lambda \in \mathbb{R}$ is called an **eigenvalue** of the matrix A if $A\mathbf{v} = \lambda\mathbf{v}$ for a nonzero column vector $\mathbf{v} \in \mathbb{R}^n$.

The vector \mathbf{v} is called an **eigenvector** of A belonging to (or associated with) the eigenvalue λ .

If λ is an eigenvalue of A then the nullspace $N(A - \lambda I)$, which is nontrivial, is called the **eigenspace** of A corresponding to λ . The eigenspace consists of all eigenvectors belonging to the eigenvalue λ plus the zero vector.

Characteristic equation

Definition. Given a square matrix A , the equation $\det(A - \lambda I) = 0$ is called the **characteristic equation** of A .

Eigenvalues λ of A are roots of the characteristic equation.

If A is an $n \times n$ matrix then $p(\lambda) = \det(A - \lambda I)$ is a polynomial of degree n . It is called the **characteristic polynomial** of A .

Theorem Any $n \times n$ matrix has at most n eigenvalues.

Example. $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$.

Characteristic equation: $\begin{vmatrix} 2 - \lambda & 1 \\ 1 & 2 - \lambda \end{vmatrix} = 0$.

$$(2 - \lambda)^2 - 1 = 0 \implies \lambda_1 = 1, \lambda_2 = 3.$$

$$(A - I)\mathbf{x} = \mathbf{0} \iff \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\iff \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \iff x + y = 0.$$

The general solution is $(-t, t) = t(-1, 1)$, $t \in \mathbb{R}$.

Thus $\mathbf{v}_1 = (-1, 1)$ is an eigenvector associated with the eigenvalue 1. The corresponding eigenspace is the line spanned by \mathbf{v}_1 .

$$(A - 3I)\mathbf{x} = \mathbf{0} \iff \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$
$$\iff \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \iff x - y = 0.$$

The general solution is $(t, t) = t(1, 1)$, $t \in \mathbb{R}$.

Thus $\mathbf{v}_2 = (1, 1)$ is an eigenvector associated with the eigenvalue 3. The corresponding eigenspace is the line spanned by \mathbf{v}_2 .

Summary. $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$.

- The matrix A has two eigenvalues: 1 and 3.
- The eigenspace of A associated with the eigenvalue 1 is the line $t(-1, 1)$.
- The eigenspace of A associated with the eigenvalue 3 is the line $t(1, 1)$.
- Eigenvectors $\mathbf{v}_1 = (-1, 1)$ and $\mathbf{v}_2 = (1, 1)$ of the matrix A form a basis for \mathbb{R}^2 .
- Geometrically, the mapping $\mathbf{x} \mapsto A\mathbf{x}$ is a stretch by a factor of 3 away from the line $x + y = 0$ in the orthogonal direction.

Example. $A = \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix}.$

Characteristic equation:

$$\begin{vmatrix} 1 - \lambda & 1 & -1 \\ 1 & 1 - \lambda & 1 \\ 0 & 0 & 2 - \lambda \end{vmatrix} = 0.$$

Expand the determinant by the 3rd row:

$$(2 - \lambda) \begin{vmatrix} 1 - \lambda & 1 \\ 1 & 1 - \lambda \end{vmatrix} = 0.$$

$$\begin{aligned} ((1 - \lambda)^2 - 1)(2 - \lambda) &= 0 \iff -\lambda(2 - \lambda)^2 = 0 \\ \implies \lambda_1 &= 0, \lambda_2 = 2. \end{aligned}$$

$$A\mathbf{x} = \mathbf{0} \iff \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

Convert the matrix to reduced row echelon form:

$$\begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & -1 \\ 0 & 0 & 2 \\ 0 & 0 & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$A\mathbf{x} = \mathbf{0} \iff \begin{cases} x + y = 0, \\ z = 0. \end{cases}$$

The general solution is $(-t, t, 0) = t(-1, 1, 0)$, $t \in \mathbb{R}$. Thus $\mathbf{v}_1 = (-1, 1, 0)$ is an eigenvector associated with the eigenvalue 0. The corresponding eigenspace is the line spanned by \mathbf{v}_1 .

$$(A - 2I)\mathbf{x} = \mathbf{0} \iff \begin{pmatrix} -1 & 1 & -1 \\ 1 & -1 & 1 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\iff \begin{pmatrix} 1 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \iff x - y + z = 0.$$

The general solution is $x = t - s$, $y = t$, $z = s$, where $t, s \in \mathbb{R}$. Equivalently,

$$\mathbf{x} = (t - s, t, s) = t(1, 1, 0) + s(-1, 0, 1).$$

Thus $\mathbf{v}_2 = (1, 1, 0)$ and $\mathbf{v}_3 = (-1, 0, 1)$ are eigenvectors associated with the eigenvalue 2.

The corresponding eigenspace is the plane spanned by \mathbf{v}_2 and \mathbf{v}_3 .

Summary. $A = \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix}.$

- The matrix A has two eigenvalues: 0 and 2.
- The eigenvalue 0 is *simple*: the corresponding eigenspace is a line.
- The eigenvalue 2 is of *multiplicity* 2: the corresponding eigenspace is a plane.
- Eigenvectors $\mathbf{v}_1 = (-1, 1, 0)$, $\mathbf{v}_2 = (1, 1, 0)$, and $\mathbf{v}_3 = (-1, 0, 1)$ of the matrix A form a basis for \mathbb{R}^3 .
- Geometrically, the map $\mathbf{x} \mapsto A\mathbf{x}$ is the projection on the plane $\text{Span}(\mathbf{v}_2, \mathbf{v}_3)$ along the lines parallel to \mathbf{v}_1 with the subsequent scaling by a factor of 2.

Eigenvalues and eigenvectors of an operator

Definition. Let V be a vector space and $L : V \rightarrow V$ be a linear operator. A number λ is called an **eigenvalue** of the operator L if $L(\mathbf{v}) = \lambda\mathbf{v}$ for a nonzero vector $\mathbf{v} \in V$. The vector \mathbf{v} is called an **eigenvector** of L associated with the eigenvalue λ . (If V is a functional space then eigenvectors are also called **eigenfunctions**.)

If $V = \mathbb{R}^n$ then the linear operator L is given by $L(\mathbf{x}) = A\mathbf{x}$, where A is an $n \times n$ matrix.

In this case, eigenvalues and eigenvectors of the operator L are precisely eigenvalues and eigenvectors of the matrix A .

Suppose $L : V \rightarrow V$ is a linear operator on a **finite-dimensional** vector space V .

Let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ be a basis for V and $g : V \rightarrow \mathbb{R}^n$ be the corresponding coordinate mapping. Let A be the matrix of L with respect to this basis. Then

$$L(\mathbf{v}) = \lambda \mathbf{v} \iff A g(\mathbf{v}) = \lambda g(\mathbf{v}).$$

Hence the eigenvalues of L coincide with those of the matrix A . Moreover, the associated eigenvectors of A are coordinates of the eigenvectors of L .

Definition. The characteristic polynomial $p(\lambda) = \det(A - \lambda I)$ of the matrix A is called the **characteristic polynomial** of the operator L .

Then eigenvalues of L are roots of its characteristic polynomial.

Theorem. The characteristic polynomial of the operator L is well defined. That is, it does not depend on the choice of a basis.

Proof: Let B be the matrix of L with respect to a different basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$. Then $A = UBU^{-1}$, where U is the transition matrix from the basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ to $\mathbf{u}_1, \dots, \mathbf{u}_n$. We have to show that $\det(A - \lambda I) = \det(B - \lambda I)$ for all $\lambda \in \mathbb{R}$. We obtain

$$\begin{aligned}\det(A - \lambda I) &= \det(UBU^{-1} - \lambda I) \\ &= \det(UBU^{-1} - U(\lambda I)U^{-1}) = \det(U(B - \lambda I)U^{-1}) \\ &= \det(U) \det(B - \lambda I) \det(U^{-1}) = \det(B - \lambda I).\end{aligned}$$

Eigenspaces

Let $L : V \rightarrow V$ be a linear operator.

For any $\lambda \in \mathbb{R}$, let V_λ denotes the set of all solutions of the equation $L(\mathbf{x}) = \lambda\mathbf{x}$.

Then V_λ is a *subspace* of V since V_λ is the *kernel* of a linear operator given by $\mathbf{x} \mapsto L(\mathbf{x}) - \lambda\mathbf{x}$.

V_λ minus the zero vector is the set of all eigenvectors of L associated with the eigenvalue λ . In particular, $\lambda \in \mathbb{R}$ is an eigenvalue of L if and only if $V_\lambda \neq \{\mathbf{0}\}$.

If $V_\lambda \neq \{\mathbf{0}\}$ then it is called the **eigenspace** of L corresponding to the eigenvalue λ .

Example. $V = C^\infty(\mathbb{R})$, $D : V \rightarrow V$, $Df = f'$.

A function $f \in C^\infty(\mathbb{R})$ is an eigenfunction of the operator D belonging to an eigenvalue λ if $f'(x) = \lambda f(x)$ for all $x \in \mathbb{R}$.

It follows that $f(x) = ce^{\lambda x}$, where c is a nonzero constant.

Thus each $\lambda \in \mathbb{R}$ is an eigenvalue of D .

The corresponding eigenspace is spanned by $e^{\lambda x}$.

Example. $V = C^\infty(\mathbb{R})$, $L : V \rightarrow V$, $Lf = f''$.

$$Lf = \lambda f \iff f''(x) - \lambda f(x) = 0 \text{ for all } x \in \mathbb{R}.$$

It follows that each $\lambda \in \mathbb{R}$ is an eigenvalue of L and the corresponding eigenspace V_λ is two-dimensional. Note that $L = D^2$, hence $Df = \mu f \implies Lf = \mu^2 f$.

If $\lambda > 0$ then $V_\lambda = \text{Span}(e^{\mu x}, e^{-\mu x})$, where $\mu = \sqrt{\lambda}$.

If $\lambda < 0$ then $V_\lambda = \text{Span}(\sin(\mu x), \cos(\mu x))$, where $\mu = \sqrt{-\lambda}$.

If $\lambda = 0$ then $V_\lambda = \text{Span}(1, x)$.