

MATH 311

Topics in Applied Mathematics I

Lecture 39:

Review for the final exam (continued).

Topics for the final exam: Part I

Elementary linear algebra (L/C 1.1–1.5, 2.1–2.2)

- Systems of linear equations: elementary operations, Gaussian elimination, back substitution.
- Matrix of coefficients and augmented matrix. Elementary row operations, row echelon form and reduced row echelon form.
- Matrix algebra. Inverse matrix.
- Determinants: explicit formulas for 2×2 and 3×3 matrices, row and column expansions, elementary row and column operations.

Topics for the final exam: Part II

Abstract linear algebra (L/C 3.1–3.6, 4.1–4.3)

- Vector spaces (vectors, matrices, polynomials, functional spaces).
- Subspaces. Nullspace, column space, and row space of a matrix.
- Span, spanning set. Linear independence.
- Bases and dimension.
- Rank and nullity of a matrix.
- Coordinates relative to a basis.
- Change of basis, transition matrix.
- Linear transformations.
- Matrix transformations.
- Matrix of a linear mapping.
- Change of basis for a linear operator.
- Similarity of matrices.

Topics for the final exam: Part III

Advanced linear algebra (L/C 5.1–5.6, 6.1, 6.3)

- Eigenvalues, eigenvectors, eigenspaces
- Characteristic polynomial
- Bases of eigenvectors, diagonalization
- Euclidean structure in \mathbb{R}^n (length, angle, dot product)
- Inner products and norms
- Orthogonal complement, orthogonal projection
- Least squares problems
- The Gram-Schmidt orthogonalization process

Topics for the final exam: Part IV

Vector analysis (L/C 8.1–8.4, 9.1–9.5, 10.1–10.3, 11.1–11.3)

- Gradient, divergence, and curl
- Fubini's Theorem
- Change of coordinates in a multiple integral
- Geometric meaning of the determinant
- Length of a curve
- Line integrals
- Green's Theorem
- Conservative vector fields
- Area of a surface
- Surface integrals
- Gauss' Theorem
- Stokes' Theorem

Problem. Consider a linear operator $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $L(\mathbf{v}) = \mathbf{v}_0 \times \mathbf{v}$, where $\mathbf{v}_0 = (3/5, 0, -4/5)$.

- (a) Find the matrix B of the operator L .
- (b) Find the range and kernel of L .
- (c) Find the eigenvalues of L .
- (d) Find the matrix of the operator L^{2018} (L applied 2018 times).

$$L(\mathbf{v}) = \mathbf{v}_0 \times \mathbf{v}, \quad \mathbf{v}_0 = (3/5, 0, -4/5).$$

Let $\mathbf{v} = (x, y, z) = x\mathbf{e}_1 + y\mathbf{e}_2 + z\mathbf{e}_3$. Then

$$L(\mathbf{v}) = \mathbf{v}_0 \times \mathbf{v} = \begin{vmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \\ 3/5 & 0 & -4/5 \\ x & y & z \end{vmatrix}$$

$$= \begin{vmatrix} 0 & -4/5 \\ y & z \end{vmatrix} \mathbf{e}_1 - \begin{vmatrix} 3/5 & -4/5 \\ x & z \end{vmatrix} \mathbf{e}_2 + \begin{vmatrix} 3/5 & 0 \\ x & y \end{vmatrix} \mathbf{e}_3$$

$$= \frac{4}{5}y\mathbf{e}_1 - \left(\frac{4}{5}x + \frac{3}{5}z\right)\mathbf{e}_2 + \frac{3}{5}y\mathbf{e}_3 = \left(\frac{4}{5}y, -\frac{4}{5}x - \frac{3}{5}z, \frac{3}{5}y\right).$$

In particular, $L(\mathbf{e}_1) = (0, -\frac{4}{5}, 0)$, $L(\mathbf{e}_2) = (\frac{4}{5}, 0, \frac{3}{5})$,
 $L(\mathbf{e}_3) = (0, -\frac{3}{5}, 0)$.

Therefore $B = \begin{pmatrix} 0 & 4/5 & 0 \\ -4/5 & 0 & -3/5 \\ 0 & 3/5 & 0 \end{pmatrix}$.

The range of the operator L is spanned by columns of the matrix B . It follows that $\text{Range}(L)$ is the plane spanned by $\mathbf{v}_1 = (0, 1, 0)$ and $\mathbf{v}_2 = (4, 0, 3)$.

The kernel of L is the nullspace of the matrix B , i.e., the solution set for the equation $B\mathbf{x} = \mathbf{0}$.

$$\begin{pmatrix} 0 & 4/5 & 0 \\ -4/5 & 0 & -3/5 \\ 0 & 3/5 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 3/4 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\implies x + \frac{3}{4}z = y = 0 \implies \mathbf{x} = t(-3/4, 0, 1).$$

Alternatively, the kernel of L is the set of vectors $\mathbf{v} \in \mathbb{R}^3$ such that $L(\mathbf{v}) = \mathbf{v}_0 \times \mathbf{v} = \mathbf{0}$.

It follows that this is the line spanned by $\mathbf{v}_0 = (3/5, 0, -4/5)$.

Characteristic polynomial of the matrix B :

$$\begin{aligned} \det(B - \lambda I) &= \begin{vmatrix} -\lambda & 4/5 & 0 \\ -4/5 & -\lambda & -3/5 \\ 0 & 3/5 & -\lambda \end{vmatrix} \\ &= -\lambda^3 - (3/5)^2\lambda - (4/5)^2\lambda = -\lambda^3 - \lambda = -\lambda(\lambda^2 + 1). \end{aligned}$$

The eigenvalues are 0 , i , and $-i$.

The matrix of the operator L^{2018} is B^{2018} .

Since the matrix B has eigenvalues 0 , i , and $-i$, it is diagonalizable in \mathbb{C}^3 . Namely, $B = UDU^{-1}$, where U is an invertible matrix with complex entries and

$$D = \begin{pmatrix} 0 & 0 & 0 \\ 0 & i & 0 \\ 0 & 0 & -i \end{pmatrix}.$$

Then $B^{2018} = UD^{2018}U^{-1}$. We have that $D^{2018} = \text{diag}(0, i^{2018}, (-i)^{2018}) = \text{diag}(0, -1, -1) = D^2$.

Hence

$$B^{2018} = UD^2U^{-1} = B^2 = \begin{pmatrix} -0.64 & 0 & -0.48 \\ 0 & -1 & 0 \\ -0.48 & 0 & -0.36 \end{pmatrix}.$$

Problem. Find the distance from the point $\mathbf{y} = (0, 0, 0, 1)$ to the subspace $V \subset \mathbb{R}^4$ spanned by vectors $\mathbf{x}_1 = (1, -1, 1, -1)$, $\mathbf{x}_2 = (1, 1, 3, -1)$, and $\mathbf{x}_3 = (-3, 7, 1, 3)$.

First we apply the Gram-Schmidt process to vectors $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ and obtain an orthogonal basis $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ for the subspace V . Next we compute the orthogonal projection \mathbf{p} of the vector \mathbf{y} onto V :

$$\mathbf{p} = \frac{\langle \mathbf{y}, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 + \frac{\langle \mathbf{y}, \mathbf{v}_2 \rangle}{\langle \mathbf{v}_2, \mathbf{v}_2 \rangle} \mathbf{v}_2 + \frac{\langle \mathbf{y}, \mathbf{v}_3 \rangle}{\langle \mathbf{v}_3, \mathbf{v}_3 \rangle} \mathbf{v}_3.$$

Then the distance from \mathbf{y} to V equals $\|\mathbf{y} - \mathbf{p}\|$.

Alternatively, we can apply the Gram-Schmidt process to vectors $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{y}$. We should obtain an orthogonal system $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4$. Then the desired distance will be $\|\mathbf{v}_4\|$.

$$\mathbf{x}_1 = (1, -1, 1, -1), \quad \mathbf{x}_2 = (1, 1, 3, -1), \\ \mathbf{x}_3 = (-3, 7, 1, 3), \quad \mathbf{y} = (0, 0, 0, 1).$$

$$\mathbf{v}_1 = \mathbf{x}_1 = (1, -1, 1, -1),$$

$$\mathbf{v}_2 = \mathbf{x}_2 - \frac{\langle \mathbf{x}_2, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 = (1, 1, 3, -1) - \frac{4}{4}(1, -1, 1, -1) \\ = (0, 2, 2, 0),$$

$$\mathbf{v}_3 = \mathbf{x}_3 - \frac{\langle \mathbf{x}_3, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 - \frac{\langle \mathbf{x}_3, \mathbf{v}_2 \rangle}{\langle \mathbf{v}_2, \mathbf{v}_2 \rangle} \mathbf{v}_2 \\ = (-3, 7, 1, 3) - \frac{-12}{4}(1, -1, 1, -1) - \frac{16}{8}(0, 2, 2, 0) \\ = (0, 0, 0, 0).$$

The Gram-Schmidt process can be used to check linear independence of vectors! It failed because the vector \mathbf{x}_3 is a linear combination of \mathbf{x}_1 and \mathbf{x}_2 . V is a plane, not a 3-dimensional subspace. To fix things, it is enough to drop \mathbf{x}_3 , i.e., we should orthogonalize vectors $\mathbf{x}_1, \mathbf{x}_2, \mathbf{y}$.

$$\begin{aligned}\tilde{\mathbf{v}}_3 &= \mathbf{y} - \frac{\langle \mathbf{y}, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 - \frac{\langle \mathbf{y}, \mathbf{v}_2 \rangle}{\langle \mathbf{v}_2, \mathbf{v}_2 \rangle} \mathbf{v}_2 \\ &= (0, 0, 0, 1) - \frac{-1}{4}(1, -1, 1, -1) - \frac{0}{8}(0, 2, 2, 0) \\ &= (1/4, -1/4, 1/4, 3/4).\end{aligned}$$

$$\|\tilde{\mathbf{v}}_3\| = \left| \left(\frac{1}{4}, -\frac{1}{4}, \frac{1}{4}, \frac{3}{4} \right) \right| = \frac{1}{4} |(1, -1, 1, 3)| = \frac{\sqrt{12}}{4} = \frac{\sqrt{3}}{2}.$$

Area, volume, and determinants

- 2×2 determinants and plane geometry


Let P be a parallelogram in the plane \mathbb{R}^2 . Suppose that vectors $\mathbf{v}_1, \mathbf{v}_2 \in \mathbb{R}^2$ are represented by adjacent sides of P . Then $\text{area}(P) = |\det A|$, where $A = (\mathbf{v}_1, \mathbf{v}_2)$, a matrix whose columns are \mathbf{v}_1 and \mathbf{v}_2 .

Consider a linear operator $L_A : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $L_A(\mathbf{v}) = A\mathbf{v}$ for any column vector \mathbf{v} . Then $\text{area}(L_A(D)) = |\det A| \text{area}(D)$ for any bounded domain D .

- 3×3 determinants and space geometry

Let Π be a parallelepiped in space \mathbb{R}^3 . Suppose that vectors $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3 \in \mathbb{R}^3$ are represented by adjacent edges of Π . Then $\text{volume}(\Pi) = |\det B|$, where $B = (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$, a matrix whose columns are $\mathbf{v}_1, \mathbf{v}_2$, and \mathbf{v}_3 .

Similarly, $\text{volume}(L_B(D)) = |\det B| \text{volume}(D)$ for any bounded domain $D \subset \mathbb{R}^3$.




Parallelepiped is a prism.

(Volume) = (area of the base) \times (height)

Area of the base = $\|\mathbf{y} \times \mathbf{z}\|$

Volume = $|\mathbf{x} \cdot (\mathbf{y} \times \mathbf{z})|$



Tetrahedron is a pyramid.

$$(\text{Volume}) = \frac{1}{3} (\text{area of the base}) \times (\text{height})$$

$$\text{Area of the base} = \frac{1}{2} \|\mathbf{y} \times \mathbf{z}\|$$

$$\implies \text{Volume} = \frac{1}{6} |\mathbf{x} \cdot (\mathbf{y} \times \mathbf{z})|$$