

MATH 323
Linear Algebra

Lecture 23:
Review for Test 2.

Topics for Test 2

Vector spaces (Leon/de Pillis 3.4–3.6)

- Basis and dimension
- Rank and nullity of a matrix
- Coordinates relative to a basis
- Change of basis, transition matrix

Linear transformations (Leon/de Pillis 4.1–4.3)

- Linear transformations
- Range and kernel
- Matrix transformations
- Matrix of a linear transformation
- Change of basis for a linear operator
- Similar matrices

Topics for Test 2

Eigenvalues and eigenvectors (Leon/de Pillis 6.1, 6.3)

- Eigenvalues, eigenvectors, eigenspaces
- Characteristic polynomial
- Diagonalization

Orthogonality (Leon/de Pillis 5.1–5.3, 5.5–5.6)

- Euclidean structure in \mathbb{R}^n
- Orthogonal complement
- Orthogonal projection
- Least squares problems
- Orthogonal sets
- The Gram-Schmidt process

Sample problems for Test 2

Problem 1. Let $A = \begin{pmatrix} 0 & -1 & 4 & 1 \\ 1 & 1 & 2 & -1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix}$.

- (i) Find the rank and the nullity of the matrix A .
- (ii) Find a basis for the row space of A , then extend this basis to a basis for \mathbb{R}^4 .
- (iii) Find a basis for the nullspace of A .

Problem 2. Let A and B be two matrices such that the product AB is well defined.

- (i) Prove that $\text{rank}(AB) \leq \text{rank}(B)$.
- (ii) Prove that $\text{rank}(AB) \leq \text{rank}(A)$.

Sample problems for Test 2

Problem 3. Complex numbers \mathbb{C} form a vector space of (real) dimension 2. Consider a function $f : \mathbb{C} \rightarrow \mathbb{C}$ given by $f(z) = (3 + 2i)z$ for all $z \in \mathbb{C}$.

- (i) Prove that f is a linear operator on the vector space \mathbb{C} .
- (ii) Find the matrix of f relative to the basis $1, i$.

Problem 4. Let V be a subspace of $F(\mathbb{R})$ spanned by functions e^x and e^{-x} . Let L be a linear operator on V such that $\begin{pmatrix} 2 & -1 \\ -3 & 2 \end{pmatrix}$ is the matrix of L relative to the basis e^x, e^{-x} . Find the matrix of L relative to the basis $\cosh x = \frac{1}{2}(e^x + e^{-x}), \sinh x = \frac{1}{2}(e^x - e^{-x})$.

Sample problems for Test 2

Problem 5. Let $A = \begin{pmatrix} 1 & 2 & 0 \\ 1 & 1 & 1 \\ 0 & 2 & 1 \end{pmatrix}$.

- (i) Find all eigenvalues of the matrix A .
- (ii) For each eigenvalue of A , find an associated eigenvector.
- (iii) Is the matrix A diagonalizable? Explain.
- (iv) Find all eigenvalues of the matrix A^2 .

Sample problems for Test 2

Problem 6. Find a linear polynomial which is the best least squares fit to the following data:

x	-2	-1	0	1	2
$f(x)$	-3	-2	1	2	5

Problem 7. Let V be a subspace of \mathbb{R}^4 spanned by the vectors $\mathbf{x}_1 = (1, 1, 1, 1)$ and $\mathbf{x}_2 = (1, 0, 3, 0)$.

- (i) Find an orthonormal basis for V .
- (ii) Find an orthonormal basis for the orthogonal complement V^\perp .
- (iii) Find the distance from the vector $\mathbf{y} = (1, 0, 0, 0)$ to the subspaces V and V^\perp .

Problem 1. Let $A = \begin{pmatrix} 0 & -1 & 4 & 1 \\ 1 & 1 & 2 & -1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix}$.

(i) Find the rank and the nullity of the matrix A .

The rank (= dimension of the row space) and the nullity (= dimension of the nullspace) of a matrix are preserved under elementary row operations. We apply such operations to convert the matrix A into row echelon form.

Interchange the 1st row with the 2nd row:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & -1 & 4 & 1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix}$$

Add 3 times the 1st row to the 3rd row, then subtract 2 times the 1st row from the 4th row:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & -1 & 4 & 1 \\ 0 & 3 & 5 & -3 \\ 2 & -1 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & -1 & 4 & 1 \\ 0 & 3 & 5 & -3 \\ 0 & -3 & -4 & 3 \end{pmatrix}$$

Multiply the 2nd row by -1 :

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 3 & 5 & -3 \\ 0 & -3 & -4 & 3 \end{pmatrix}$$

Add the 4th row to the 3rd row:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -3 & -4 & 3 \end{pmatrix}$$

Add 3 times the 2nd row to the 4th row:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -16 & 0 \end{pmatrix}$$

Add 16 times the 3rd row to the 4th row:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Now that the matrix is in row echelon form, its rank equals the number of nonzero rows, which is 3. Since $(\text{rank of } A) + (\text{nullity of } A) = (\text{the number of columns of } A) = 4$, it follows that the nullity of A equals 1.

Problem 1. Let $A = \begin{pmatrix} 0 & -1 & 4 & 1 \\ 1 & 1 & 2 & -1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix}$.

(ii) Find a basis for the row space of A , then extend this basis to a basis for \mathbb{R}^4 .

The row space of a matrix is invariant under elementary row operations. Therefore the row space of the matrix A is the same as the row space of its row echelon form:

$$\begin{pmatrix} 0 & -1 & 4 & 1 \\ 1 & 1 & 2 & -1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

The nonzero rows of the latter matrix are linearly independent so that they form a basis for its row space:

$$\mathbf{v}_1 = (1, 1, 2, -1), \quad \mathbf{v}_2 = (0, 1, -4, -1), \quad \mathbf{v}_3 = (0, 0, 1, 0).$$

To extend the basis $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ to a basis for \mathbb{R}^4 , we need a vector $\mathbf{v}_4 \in \mathbb{R}^4$ that is not a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$.

It is known that at least one of the vectors $\mathbf{e}_1 = (1, 0, 0, 0)$, $\mathbf{e}_2 = (0, 1, 0, 0)$, $\mathbf{e}_3 = (0, 0, 1, 0)$, and $\mathbf{e}_4 = (0, 0, 0, 1)$ can be chosen as \mathbf{v}_4 .

In particular, the vectors $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{e}_4$ form a basis for \mathbb{R}^4 .

This follows from the fact that the 4×4 matrix whose rows are these vectors is not singular:

$$\begin{vmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{vmatrix} = 1 \neq 0.$$

Problem 1. Let $A = \begin{pmatrix} 0 & -1 & 4 & 1 \\ 1 & 1 & 2 & -1 \\ -3 & 0 & -1 & 0 \\ 2 & -1 & 0 & 1 \end{pmatrix}$.

(iii) Find a basis for the nullspace of A .

The nullspace of A is the solution set of the system of linear homogeneous equations with A as the coefficient matrix. To solve the system, we convert A to reduced row echelon form:

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & -1 \\ 0 & 1 & -4 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\implies x_1 = x_2 - x_4 = x_3 = 0$$

General solution: $(x_1, x_2, x_3, x_4) = (0, t, 0, t) = t(0, 1, 0, 1)$.

Thus the vector $(0, 1, 0, 1)$ forms a basis for the nullspace of A .

Problem 2. Let A and B be two matrices such that the product AB is well defined.

(i) Prove that $\text{rank}(AB) \leq \text{rank}(B)$.

Suppose that $B\mathbf{x} = \mathbf{0}$ for some column vector \mathbf{x} . Then $(AB)\mathbf{x} = A(B\mathbf{x}) = A\mathbf{0} = \mathbf{0}$. It follows that the nullspace of B is contained in the nullspace of AB . Consequently, $\text{nullity}(B) \leq \text{nullity}(AB)$. Since matrices AB and B have the same number of columns, we obtain $\text{rank}(AB) \leq \text{rank}(B)$.

(ii) Prove that $\text{rank}(AB) \leq \text{rank}(A)$.

Note that $\text{rank}(M) = \text{rank}(M^T)$ for any matrix M . In particular, $\text{rank}(AB) = \text{rank}((AB)^T) = \text{rank}(B^T A^T)$. By the above, $\text{rank}(B^T A^T) \leq \text{rank}(A^T) = \text{rank}(A)$.

Remark. Alternatively, one can show that the row space of AB is contained in the row space of B while the column space of AB is contained in the column space of A .

Problem 3. Complex numbers \mathbb{C} form a vector space of (real) dimension 2. Consider a function $f : \mathbb{C} \rightarrow \mathbb{C}$ given by $f(z) = (3 + 2i)z$ for all $z \in \mathbb{C}$.

(i) Prove that f is a linear operator on the vector space \mathbb{C} .

We need to show that $f(z + w) = f(z) + f(w)$ for $z, w \in \mathbb{C}$ and $f(rz) = rf(z)$ for all $r \in \mathbb{R}$ and $z \in \mathbb{C}$. The first condition means that $(3 + 2i)(z + w) = (3 + 2i)z + (3 + 2i)w$; it follows from the distributive law for complex numbers. The second condition means that $(3 + 2i)(rz) = r((3 + 2i)z)$; it follows from commutativity and associativity of multiplication of complex numbers.

(ii) Find the matrix of f relative to the basis $1, i$.

Columns of the matrix are coordinates of the images $f(1)$ and $f(i)$ relative to the basis $1, i$. Observe that the coordinates of a complex number $x + yi$ are (x, y) . We obtain that

$f(1) = 3 + 2i$ and $f(i) = (3 + 2i)i = -2 + 3i$. Hence the

matrix of f is $\begin{pmatrix} 3 & -2 \\ 2 & 3 \end{pmatrix}$.

Problem 4. Let V be a subspace of $F(\mathbb{R})$ spanned by functions e^x and e^{-x} . Let L be a linear operator on V such that $\begin{pmatrix} 2 & -1 \\ -3 & 2 \end{pmatrix}$ is the matrix of L relative to the basis e^x, e^{-x} . Find the matrix of L relative to the basis $\cosh x = \frac{1}{2}(e^x + e^{-x}), \sinh x = \frac{1}{2}(e^x - e^{-x})$.

Let A denote the matrix of the operator L relative to the basis e^x, e^{-x} (which is given) and B denote the matrix of L relative to the basis $\cosh x, \sinh x$ (which is to be found). By definition of the functions $\cosh x$ and $\sinh x$, the transition matrix from $\cosh x, \sinh x$ to e^x, e^{-x} is $U = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$.

It follows that $B = U^{-1}AU$. We obtain that

$$B = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -3 & 2 \end{pmatrix} \cdot \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 4 \end{pmatrix}.$$

Problem 5. Let $A = \begin{pmatrix} 1 & 2 & 0 \\ 1 & 1 & 1 \\ 0 & 2 & 1 \end{pmatrix}$.

(i) Find all eigenvalues of the matrix A .

The eigenvalues of A are roots of the characteristic equation $\det(A - \lambda I) = 0$. We obtain that

$$\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 2 & 0 \\ 1 & 1 - \lambda & 1 \\ 0 & 2 & 1 - \lambda \end{vmatrix}$$

$$\begin{aligned} &= (1 - \lambda)^3 - 2(1 - \lambda) - 2(1 - \lambda) = (1 - \lambda)((1 - \lambda)^2 - 4) \\ &= (1 - \lambda)((1 - \lambda) - 2)((1 - \lambda) + 2) = -(\lambda - 1)(\lambda + 1)(\lambda - 3). \end{aligned}$$

Hence the matrix A has three eigenvalues: -1 , 1 , and 3 .

Problem 5. Let $A = \begin{pmatrix} 1 & 2 & 0 \\ 1 & 1 & 1 \\ 0 & 2 & 1 \end{pmatrix}$.

(ii) For each eigenvalue of A , find an associated eigenvector.

An eigenvector $\mathbf{v} = (x, y, z)$ of the matrix A associated with an eigenvalue λ is a nonzero solution of the vector equation

$$(A - \lambda I)\mathbf{v} = \mathbf{0} \iff \begin{pmatrix} 1 - \lambda & 2 & 0 \\ 1 & 1 - \lambda & 1 \\ 0 & 2 & 1 - \lambda \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

To solve the equation, we convert the matrix $A - \lambda I$ to reduced row echelon form.

First consider the case $\lambda = -1$. The row reduction yields

$$A + I = \begin{pmatrix} 2 & 2 & 0 \\ 1 & 2 & 1 \\ 0 & 2 & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 2 & 2 \end{pmatrix}$$

$$\rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 2 & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}.$$

Hence

$$(A + I)\mathbf{v} = \mathbf{0} \iff \begin{cases} x - z = 0, \\ y + z = 0. \end{cases}$$

The general solution is $x = t$, $y = -t$, $z = t$, where $t \in \mathbb{R}$. In particular, $\mathbf{v}_1 = (1, -1, 1)$ is an eigenvector of A associated with the eigenvalue -1 .

Secondly, consider the case $\lambda = 1$. The row reduction yields

$$A - I = \begin{pmatrix} 0 & 2 & 0 \\ 1 & 0 & 1 \\ 0 & 2 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 0 & 2 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 2 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Hence

$$(A - I)\mathbf{v} = \mathbf{0} \iff \begin{cases} x + z = 0, \\ y = 0. \end{cases}$$

The general solution is $x = -t$, $y = 0$, $z = t$, where $t \in \mathbb{R}$. In particular, $\mathbf{v}_2 = (-1, 0, 1)$ is an eigenvector of A associated with the eigenvalue 1.

Finally, consider the case $\lambda = 3$. The row reduction yields

$$\begin{aligned} A-3I &= \begin{pmatrix} -2 & 2 & 0 \\ 1 & -2 & 1 \\ 0 & 2 & -2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & -1 & 0 \\ 1 & -2 & 1 \\ 0 & 2 & -2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & -1 & 0 \\ 0 & -1 & 1 \\ 0 & 2 & -2 \end{pmatrix} \\ &\rightarrow \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 2 & -2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{pmatrix}. \end{aligned}$$

Hence

$$(A - 3I)\mathbf{v} = \mathbf{0} \iff \begin{cases} x - z = 0, \\ y - z = 0. \end{cases}$$

The general solution is $x = t$, $y = t$, $z = t$, where $t \in \mathbb{R}$. In particular, $\mathbf{v}_3 = (1, 1, 1)$ is an eigenvector of A associated with the eigenvalue 3.

Problem 5. Let $A = \begin{pmatrix} 1 & 2 & 0 \\ 1 & 1 & 1 \\ 0 & 2 & 1 \end{pmatrix}$.

(iii) Is the matrix A diagonalizable? Explain.

The matrix A is diagonalizable, i.e., there exists a basis for \mathbb{R}^3 formed by its eigenvectors.

Namely, the vectors $\mathbf{v}_1 = (1, -1, 1)$, $\mathbf{v}_2 = (-1, 0, 1)$, and $\mathbf{v}_3 = (1, 1, 1)$ are eigenvectors of the matrix A belonging to distinct eigenvalues. Therefore these vectors are linearly independent. It follows that $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ is a basis for \mathbb{R}^3 .

Alternatively, the existence of a basis for \mathbb{R}^3 consisting of eigenvectors of A already follows from the fact that the matrix A has three distinct eigenvalues.

Problem 5. Let $A = \begin{pmatrix} 1 & 2 & 0 \\ 1 & 1 & 1 \\ 0 & 2 & 1 \end{pmatrix}$.

(iv) Find all eigenvalues of the matrix A^2 .

Suppose that \mathbf{v} is an eigenvector of the matrix A associated with an eigenvalue λ , that is, $\mathbf{v} \neq \mathbf{0}$ and $A\mathbf{v} = \lambda\mathbf{v}$. Then

$$A^2\mathbf{v} = A(A\mathbf{v}) = A(\lambda\mathbf{v}) = \lambda(A\mathbf{v}) = \lambda(\lambda\mathbf{v}) = \lambda^2\mathbf{v}.$$

Therefore \mathbf{v} is also an eigenvector of the matrix A^2 and the associated eigenvalue is λ^2 . We already know that the matrix A has eigenvalues -1 , 1 , and 3 . It follows that A^2 has eigenvalues 1 and 9 .

Since a 3×3 matrix can have up to 3 eigenvalues, we need an additional argument to show that 1 and 9 are the only eigenvalues of A^2 . One reason is that the eigenvalue 1 has multiplicity 2.

Problem 6. Find a linear polynomial which is the best least squares fit to the following data:

x	-2	-1	0	1	2
$f(x)$	-3	-2	1	2	5

We are looking for a function $f(x) = c_1 + c_2x$, where c_1, c_2 are unknown coefficients. The data of the problem give rise to an overdetermined system of linear equations in variables c_1 and c_2 :

$$\begin{cases} c_1 - 2c_2 = -3, \\ c_1 - c_2 = -2, \\ c_1 = 1, \\ c_1 + c_2 = 2, \\ c_1 + 2c_2 = 5. \end{cases}$$

This system is inconsistent.


We can represent the system as a matrix equation $A\mathbf{c} = \mathbf{y}$, where

$$A = \begin{pmatrix} 1 & -2 \\ 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{pmatrix}, \quad \mathbf{c} = \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}, \quad \mathbf{y} = \begin{pmatrix} -3 \\ -2 \\ 1 \\ 2 \\ 5 \end{pmatrix}.$$

The least squares solution \mathbf{c} of the above system is a solution of the normal system $A^T A \mathbf{c} = A^T \mathbf{y}$:

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ -2 & -1 & 0 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & -2 \\ 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ -2 & -1 & 0 & 1 & 2 \end{pmatrix} \begin{pmatrix} -3 \\ -2 \\ 1 \\ 2 \\ 5 \end{pmatrix}$$
$$\iff \begin{pmatrix} 5 & 0 \\ 0 & 10 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} 3 \\ 20 \end{pmatrix} \iff \begin{cases} c_1 = 3/5 \\ c_2 = 2 \end{cases}$$

Thus the function $f(x) = \frac{3}{5} + 2x$ is the best least squares fit to the above data among linear polynomials.



Problem 7. Let V be a subspace of \mathbb{R}^4 spanned by the vectors $\mathbf{x}_1 = (1, 1, 1, 1)$ and $\mathbf{x}_2 = (1, 0, 3, 0)$.

(i) Find an orthonormal basis for V .

First we apply the Gram-Schmidt orthogonalization process to vectors $\mathbf{x}_1, \mathbf{x}_2$ and obtain an orthogonal basis $\mathbf{v}_1, \mathbf{v}_2$ for the subspace V :


$$\mathbf{v}_1 = \mathbf{x}_1 = (1, 1, 1, 1),$$

$$\mathbf{v}_2 = \mathbf{x}_2 - \frac{\mathbf{x}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 = (1, 0, 3, 0) - \frac{4}{4}(1, 1, 1, 1) = (0, -1, 2, -1).$$

Then we normalize vectors $\mathbf{v}_1, \mathbf{v}_2$ to obtain an orthonormal basis $\mathbf{w}_1, \mathbf{w}_2$ for V :

$$\|\mathbf{v}_1\| = 2 \implies \mathbf{w}_1 = \frac{\mathbf{v}_1}{\|\mathbf{v}_1\|} = \frac{1}{2}(1, 1, 1, 1)$$

$$\|\mathbf{v}_2\| = \sqrt{6} \implies \mathbf{w}_2 = \frac{\mathbf{v}_2}{\|\mathbf{v}_2\|} = \frac{1}{\sqrt{6}}(0, -1, 2, -1)$$



Problem 7. Let V be a subspace of \mathbb{R}^4 spanned by the vectors $\mathbf{x}_1 = (1, 1, 1, 1)$ and $\mathbf{x}_2 = (1, 0, 3, 0)$.

(iii) Find the distance from the vector $\mathbf{y} = (1, 0, 0, 0)$ to the subspaces V and V^\perp .

The vector \mathbf{y} is uniquely decomposed as $\mathbf{y} = \mathbf{p} + \mathbf{o}$, where $\mathbf{p} \in V$ and $\mathbf{o} \in V^\perp$. Then \mathbf{p} is the orthogonal projection of \mathbf{y} onto the subspace V while \mathbf{o} is the orthogonal projection of \mathbf{y} onto the orthogonal complement V^\perp . Hence the distance from \mathbf{y} to V equals $\|\mathbf{y} - \mathbf{p}\| = \|\mathbf{o}\|$ and the distance from \mathbf{y} to V^\perp equals $\|\mathbf{y} - \mathbf{o}\| = \|\mathbf{p}\|$.

Since vectors $\mathbf{v}_1 = (1, 1, 1, 1)$ and $\mathbf{v}_2 = (0, -1, 2, -1)$ form an orthogonal basis for V , it follows that

$$\mathbf{p} = \frac{\mathbf{y} \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 + \frac{\mathbf{y} \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2 = \frac{1}{4} \mathbf{v}_1 + \frac{0}{6} \mathbf{v}_2 = \frac{1}{4} (1, 1, 1, 1).$$

Then $\mathbf{o} = \mathbf{y} - \mathbf{p} = \frac{1}{4} (3, -1, -1, -1)$, $\|\mathbf{o}\| = \frac{\sqrt{3}}{2}$, $\|\mathbf{p}\| = \frac{1}{2}$.

Problem 7. Let V be a subspace of \mathbb{R}^4 spanned by the vectors $\mathbf{x}_1 = (1, 1, 1, 1)$ and $\mathbf{x}_2 = (1, 0, 3, 0)$.

(ii) Find an orthonormal basis for the orthogonal complement V^\perp .

Since the subspace V is spanned by vectors $(1, 1, 1, 1)$ and $(1, 0, 3, 0)$, it is the row space of the matrix

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 3 & 0 \end{pmatrix}.$$

Then the orthogonal complement V^\perp is the nullspace of A . To find the nullspace, we convert the matrix A to reduced row echelon form:

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 3 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 3 & 0 \\ 1 & 1 & 1 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 3 & 0 \\ 0 & 1 & -2 & 1 \end{pmatrix}.$$

Hence a vector $(x_1, x_2, x_3, x_4) \in \mathbb{R}^4$ belongs to V^\perp if and only if

$$\begin{pmatrix} 1 & 0 & 3 & 0 \\ 0 & 1 & -2 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\iff \begin{cases} x_1 + 3x_3 = 0 \\ x_2 - 2x_3 + x_4 = 0 \end{cases} \iff \begin{cases} x_1 = -3x_3 \\ x_2 = 2x_3 - x_4 \end{cases}$$

The general solution of the system is $(x_1, x_2, x_3, x_4) = (-3t, 2t - s, t, s) = t(-3, 2, 1, 0) + s(0, -1, 0, 1)$, where $t, s \in \mathbb{R}$.

It follows that V^\perp is spanned by vectors $\mathbf{x}_3 = (0, -1, 0, 1)$ and $\mathbf{x}_4 = (-3, 2, 1, 0)$.

Since the vectors $\mathbf{x}_3 = (0, -1, 0, 1)$ and $\mathbf{x}_4 = (-3, 2, 1, 0)$ are linearly independent, they form a basis for the subspace V^\perp .

It remains to orthogonalize and normalize this basis:

$$\mathbf{v}_3 = \mathbf{x}_3 = (0, -1, 0, 1),$$

$$\begin{aligned}\mathbf{v}_4 &= \mathbf{x}_4 - \frac{\mathbf{x}_4 \cdot \mathbf{v}_3}{\mathbf{v}_3 \cdot \mathbf{v}_3} \mathbf{v}_3 = (-3, 2, 1, 0) - \frac{-2}{2}(0, -1, 0, 1) \\ &= (-3, 1, 1, 1),\end{aligned}$$

$$\|\mathbf{v}_3\| = \sqrt{2} \implies \mathbf{w}_3 = \frac{\mathbf{v}_3}{\|\mathbf{v}_3\|} = \frac{1}{\sqrt{2}}(0, -1, 0, 1),$$

$$\|\mathbf{v}_4\| = \sqrt{12} = 2\sqrt{3} \implies \mathbf{w}_4 = \frac{\mathbf{v}_4}{\|\mathbf{v}_4\|} = \frac{1}{2\sqrt{3}}(-3, 1, 1, 1).$$

Thus the vectors $\mathbf{w}_3 = \frac{1}{\sqrt{2}}(0, -1, 0, 1)$ and $\mathbf{w}_4 = \frac{1}{2\sqrt{3}}(-3, 1, 1, 1)$ form an orthonormal basis for V^\perp .