

MATH 433
Applied Algebra

Lecture 36:
Review for Exam 3.

Topics for Exam 3

- Order of an element in a group
- Subgroups
- Cyclic groups
- Cosets
- Lagrange's Theorem
- Isomorphism of groups

- The ISBN code
- Binary codes, error detection and error correction
- Linear codes, generator matrix
- Coset leaders, coset decoding table
- Parity-check matrix, syndromes

- Division of polynomials
- Greatest common divisor of polynomials
- Factorisation of polynomials

Sample problems

Problem 1. Suppose $\pi, \sigma \in S(5)$ are permutations of order 3. What are possible values for the order of the permutation $\pi\sigma$.

Problem 2. Suppose H and K are subgroups of a group G . Is the union $H \cup K$ necessarily a subgroup of G ? Is the intersection $H \cap K$ necessarily a subgroup of G ?

Problem 3. Prove that the group $(\mathbb{Q} \setminus \{0\}, \times)$ is not cyclic.

Sample problems

Problem 4. Suppose G is a group of order 125. Show that G contains an element of order 5.

Problem 5. The group (G_{15}, \times) has subgroups of what orders?

Problem 6. Determine which of the following groups of order 6 are isomorphic and which are not: \mathbb{Z}_6 , $\mathbb{Z}_3 \times \mathbb{Z}_2$, $S(3)$, and $D(3)$.

Sample problems

Problem 7. Let $f : \mathbf{B}^3 \rightarrow \mathbf{B}^7$ be the coding function that sends each three-character word abc in the alphabet $\mathbf{B} = \{0, 1\}$ to the codeword $abcabcy$, where y is the inverted parity bit of the word abc (i.e., $y = 0$ if abc contains an odd number of 1's and $y = 1$ otherwise). How many errors will this code detect? correct? Is this code linear?

Problem 8. Let $f : \mathbf{B}^3 \rightarrow \mathbf{B}^6$ be a linear coding function defined by the generator matrix

$$\begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}.$$

Suppose that a message encoded by this function is received with errors as 101101 010101 011111. Correct errors and decode the received message.

Sample problems

Problem 9. Find a greatest common divisor of polynomials $p(x) = x^4 - 2x^3 + 5x^2 - 4x + 4$ and $q(x) = 2x^3 - 3x^2 + 5x - 2$ over \mathbb{R} .

Problem 10. Factorise a polynomial $p(x) = x^3 - 3x^2 + 3x - 2$ into irreducible factors over the field \mathbb{Z}_7 .

Problem 1. Suppose $\pi, \sigma \in S(5)$ are permutations of order 3. What are possible values for the order of the permutation $\pi\sigma$.

The order of a permutation equals the least common multiple of the cycle lengths in its cycle decomposition. Hence it equals 3 only if the cycles are of length 1 or 3 (at least one cycle of length 3 is required). For permutations $\pi, \sigma \in S(5)$, this implies that both are cycles of length 3. Note that π and σ are not disjoint.

Up to relabeling of the set $\{1, 2, 3, 4, 5\}$, we can assume that $\pi = (1\ 2\ 3)$. As for σ , there are several possible choices: $\sigma_1 = (1\ 4\ 5)$, $\sigma_2 = (1\ 2\ 4)$, $\sigma_3 = (1\ 2\ 3)$, $\sigma_4 = (1\ 3\ 2)$. Namely, $\sigma = \sigma_1$ if there is only one element that both π and σ move, $\sigma = \sigma_2$ if there are two such elements, and $\sigma = \sigma_3$ or $\sigma = \sigma_4$ if π and σ move the same three elements.

We have $\pi\sigma_1 = (1\ 4\ 5\ 2\ 3)$, $\pi\sigma_2 = (1\ 3)(2\ 4)$, $\pi\sigma_3 = (1\ 3\ 2)$, and $\pi\sigma_4 = \text{id}$. Thus $o(\pi\sigma)$ can be 1, 2, 3 or 5.

Problem 2. Suppose H and K are subgroups of a group G . Is the union $H \cup K$ necessarily a subgroup of G ? Is the intersection $H \cap K$ necessarily a subgroup of G ?

The union $H \cup K$ is a subgroup of G only if $H \subset K$ or $K \subset H$ (so that $H \cup K$ coincides with one of the subgroups H and K).

Otherwise $H \cup K$ is not closed under the group operation. Indeed, if neither of the subgroups contains the other, we can find an element $h \in H \setminus K$ and an element $k \in K \setminus H$. Let $g = hk$. Then $g \notin H$ as otherwise $k = h^{-1}g \in H$, a contradiction. Similarly, $g \notin K$ as otherwise $h = gk^{-1} \in K$, another contradiction. Thus $h, k \in H \cup K$ while $hk \notin H \cup K$.

The intersection $H \cap K$ of two subgroups is always a subgroup (see lecture notes and the textbook).

Problem 3. Prove that the group $(\mathbb{Q} \setminus \{0\}, \times)$ is not cyclic.

Take any non-zero rational number r . It can be represented as a reduced fraction: $r = \frac{m}{n}$, where m and n are non-zero integers and $\gcd(m, n) = 1$.

The cyclic group $\langle r \rangle$ consists of fractions $\frac{m}{n}, \frac{m^2}{n^2}, \frac{m^3}{n^3}, \dots$, fractions $\frac{n}{m}, \frac{n^2}{m^2}, \frac{n^3}{m^3}, \dots$, and 1. Note that all fractions are reduced.

The numbers m and n can have only finitely many prime divisors. Since there are infinitely many prime numbers, we can find a prime number p that divides neither m nor n . It is easy to see that $p \notin \langle r \rangle$. Thus $\langle r \rangle \neq \mathbb{Q} \setminus \{0\}$.

Problem 4. Suppose G is a group of order 125. Show that G contains an element of order 5.

It follows from Lagrange's Theorem that the order of any element of the group G divides 125. Hence the only orders we can expect are 1, 5, 25, and 125.

Let g be any element of G different from the identity element. Then the order of g is 5, 25 or 125.

If $o(g) = 5$ then we are done.

If $o(g) = 25$ then the element g^5 has order 5.

If $o(g) = 125$ then the element g^{25} has order 5.

Problem 5. The group (G_{15}, \times) has subgroups of what orders?

G_{15} is the multiplicative group of invertible congruence classes modulo 15. It has 8 elements:

$$[1], [2], [4], [7], [8], [11], [13], [14].$$

By Lagrange's Theorem, a subgroup of G_{15} can be of order 1, 2, 4 or 8. First we find the cyclic subgroups of G_{15} . These are $\{[1]\}$, $\{[1], [4]\}$, $\{[1], [11]\}$, $\{[1], [14]\}$, $\{[1], [2], [4], [8]\}$, and $\{[1], [4], [7], [13]\}$.

Hence we have cyclic subgroups of orders 1, 2 and 4. Also, the entire group G_{15} is a subgroup of order 8.

Remark. The only other subgroup of G_{15} is a non-cyclic group $\{[1], [4], [11], [14]\}$.

Problem 6. Determine which of the following groups of order 6 are isomorphic and which are not: \mathbb{Z}_6 , $\mathbb{Z}_3 \times \mathbb{Z}_2$, $S(3)$, and $D(3)$.

$\mathbb{Z}_3 \times \mathbb{Z}_2$ is an additive group, where the addition is defined by $(g, h) + (g', h') = (g + g', h + h')$. It is easy to check that the element $([1]_3, [1]_2)$ has order 6. Therefore it generates the entire group so that $\mathbb{Z}_3 \times \mathbb{Z}_2$ is cyclic. Hence it is isomorphic to \mathbb{Z}_6 as another cyclic group of order 6.

$D(3)$ is a dihedral group, the group of symmetries of an equilateral triangle. Any symmetry permutes vertices of the triangle. Once we label the vertices as 1, 2, and 3, each symmetry from $D(3)$ is assigned a permutation from the symmetric group $S(3)$. This correspondence is actually an isomorphism.

Neither of the groups \mathbb{Z}_6 and $\mathbb{Z}_3 \times \mathbb{Z}_2$ is isomorphic to $S(3)$ or $D(3)$ since the first two groups are commutative while the other two are not.

Problem 7. Let $f : \mathbf{B}^3 \rightarrow \mathbf{B}^7$ be the coding function that sends each three-character word abc in the alphabet $\mathbf{B} = \{0, 1\}$ to the codeword $abcabcy$, where y is the inverted parity bit of the word abc (i.e., $y = 0$ if abc contains an odd number of 1's and $y = 1$ otherwise). How many errors will this code detect? correct? Is this code linear?

First we list all 8 codewords for the given code:

0000001, 0010010, 0100100, 0110111,
1001000, 1011011, 1101101, 1111110.

Then we determine the minimum distance between distinct codewords. By inspection, it is 3. Therefore the code allows to detect 2 errors and to correct 1 error.

For any linear coding function $f_0 : \mathbf{B}^3 \rightarrow \mathbf{B}^7$, the set of codewords is a subspace of \mathbf{B}^7 . As a consequence, it contains the zero word. Since the zero word is not a codeword for the function f , this code is not linear.

Problem 8. Let $f : \mathbf{B}^3 \rightarrow \mathbf{B}^6$ be a linear coding function defined by the generator matrix

$$\begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}.$$

Suppose that a message encoded by this function is received with errors as 101101 010101 011111. Correct errors and decode the received message.

The coding function is given by $f(w) = wG$, where G is the generator matrix and w is regarded as a row vector. The 8 codewords are linear combinations of rows of the generator matrix:

$$\begin{aligned} &000000, 001011, 010110, 011101, \\ &100101, 101110, 110011, 111000. \end{aligned}$$

Every received word is corrected to the closest codeword. The corrected message is 100101 011101 011101. Since the code is systematic, decoding consists of truncating the codewords to 3 digits: 100 011 011.

Problem 9. Find a greatest common divisor of polynomials

$$p(x) = x^4 - 2x^3 + 5x^2 - 4x + 4 \quad \text{and}$$

$$q(x) = 2x^3 - 3x^2 + 5x - 2 \quad \text{over } \mathbb{R}.$$

$\gcd(p, q)$ can be found using the Euclidean algorithm. First

$$\begin{aligned} \text{we divide } p \text{ by } q: \quad & x^4 - 2x^3 + 5x^2 - 4x + 4 = \\ & = (2x^3 - 3x^2 + 5x - 2)\left(\frac{1}{2}x - \frac{1}{4}\right) + \frac{7}{4}x^2 - \frac{7}{4}x + \frac{7}{2}. \end{aligned}$$

Hence $\gcd(p, q) = \gcd(q, r)$, where $r(x) = \frac{7}{4}x^2 - \frac{7}{4}x + \frac{7}{2}$ is the remainder of p by q . It is convenient to replace the polynomial r by its scalar multiple $\tilde{r}(x) = \frac{4}{7}r(x) = x^2 - x + 2$.

Clearly, $\gcd(q, r) = \gcd(q, \tilde{r})$.

Next we divide q by \tilde{r} :

$$2x^3 - 3x^2 + 5x - 2 = (x^2 - x + 2)(2x - 1).$$

Since \tilde{r} divides q , it follows that $\gcd(q, \tilde{r}) = \tilde{r}$.

Finally, $\gcd(p, q) = x^2 - x + 2$.

Problem 10. Factorise a polynomial $p(x) = x^3 - 3x^2 + 3x - 2$ into irreducible factors over the field \mathbb{Z}_7 .

A quadratic or cubic polynomial is irreducible if and only if it has no roots. Indeed, if such a polynomial splits into a product of two non-constant polynomials, then at least one of the factors is linear. This implies that the original polynomial has a root.

Let us look for the roots of $p(x)$: $p(0) = -2$, $p(1) = -1$, $p(2) = 0$. Hence $p(x)$ is divisible by $x - 2$:

$$x^3 - 3x^2 + 3x - 2 = (x - 2)(x^2 - x + 1).$$

Now let us look for the roots of the polynomial $q(x) = x^2 - x + 1$. Note that values 0 and 1 can be skipped this time. We obtain $q(2) = 3$, $q(3) = 0$. Hence $q(x)$ is divisible by $x - 3$: $x^2 - x + 1 = (x - 3)(x + 2)$.

Thus $x^3 - 3x^2 + 3x - 2 = (x - 2)(x - 3)(x + 2)$ over the field \mathbb{Z}_7 .