

MATH 433

Applied Algebra

**Lecture 30:**

**Direct product of groups.**

**Quotient group.**

## Direct product of groups

Given nonempty sets  $G$  and  $H$ , the Cartesian product  $G \times H$  is the set of all ordered pairs  $(g, h)$  such that  $g \in G$  and  $h \in H$ . Suppose  $*$  is a binary operation on  $G$  and  $\star$  is a binary operation on  $H$ . Then we can define a binary operation  $\bullet$  on  $G \times H$  by

$$(g_1, h_1) \bullet (g_2, h_2) = (g_1 * g_2, h_1 \star h_2).$$

**Proposition 1** The closure axiom holds for the operation  $\bullet$  if and only if it holds for both  $*$  and  $\star$ .

**Proposition 2** The operation  $\bullet$  is associative if and only if both  $*$  and  $\star$  are associative.

**Proposition 3** A pair  $(e_G, e_H)$  is the identity element in  $G \times H$  if and only if  $e_G$  is the identity element in  $G$  and  $e_H$  is the identity element in  $H$ .

**Proposition 4**  $(g', h') = (g, h)^{-1}$  in  $G \times H$  if and only if  $g' = g^{-1}$  in  $G$  and  $h' = h^{-1}$  in  $H$ .

## Direct product of groups

Given nonempty sets  $G$  and  $H$ , the Cartesian product  $G \times H$  is the set of all ordered pairs  $(g, h)$  such that  $g \in G$  and  $h \in H$ . Suppose  $*$  is a binary operation on  $G$  and  $\star$  is a binary operation on  $H$ . Then we can define a binary operation  $\bullet$  on  $G \times H$  by

$$(g_1, h_1) \bullet (g_2, h_2) = (g_1 * g_2, h_1 \star h_2).$$

**Theorem** The set  $G \times H$  with the operation  $\bullet$  is a group if and only if both  $(G, *)$  and  $(H, \star)$  are groups.

The group  $G \times H$  is called the **direct product** of the groups  $G$  and  $H$ . Usually the same notation (multiplicative or additive) is used for all three groups:

$$\begin{aligned}(g_1, h_1)(g_2, h_2) &= (g_1 g_2, h_1 h_2) \text{ or} \\ (g_1, h_1) + (g_2, h_2) &= (g_1 + g_2, h_1 + h_2).\end{aligned}$$

Similarly, we can define the direct product  $G_1 \times G_2 \times \cdots \times G_n$  of any finite collection of groups  $G_1, G_2, \dots, G_n$ .

*Example.*  $\mathbb{Z}_2 \times \mathbb{Z}_3$  (with addition in  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$ ).

The group consists of 6 elements. It is Abelian since  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$  are both Abelian. The identity element is  $([0]_2, [0]_3)$ .

Let  $g = ([1]_2, [1]_3)$ . Then  $2g = g + g = ([0]_2, [2]_3)$ ,  
 $3g = ([1]_2, [0]_3)$ ,  $4g = ([0]_2, [1]_3)$ ,  $5g = ([1]_2, [2]_3)$ , and  
 $6g = ([0]_2, [0]_3)$ . It follows that  $\mathbb{Z}_2 \times \mathbb{Z}_3$  is a cyclic group,  
 $\mathbb{Z}_2 \times \mathbb{Z}_3 = \langle g \rangle$ .

**Theorem** If  $g$  has finite order in a group  $G$  and  $h$  has finite order in a group  $H$ , then  $(g, h)$  has finite order in  $G \times H$  equal to  $\text{lcm}(o(g), o(h))$ .

**Theorem** The direct product of nontrivial cyclic groups is cyclic if and only if they are all finite and their orders are pairwise coprime.

For example, groups  $\mathbb{Z}_3 \times \mathbb{Z}_5$ ,  $\mathbb{Z}_4 \times \mathbb{Z}_{15}$ , and  $\mathbb{Z}_2 \times \mathbb{Z}_5 \times \mathbb{Z}_7$  are cyclic while groups  $\mathbb{Z}_4 \times \mathbb{Z}_6$ ,  $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3$ ,  $\mathbb{Z}_3 \times \mathbb{Z}$ , and  $\mathbb{Z} \times \mathbb{Z}$  are not.

## Quotient space

Let  $X$  be a nonempty set and  $\sim$  be an equivalence relation on  $X$ . Given an element  $x \in X$ , the **equivalence class** of  $x$ , denoted  $[x]_{\sim}$  or simply  $[x]$ , is the set of all elements of  $X$  that are **equivalent** (i.e., related by  $\sim$ ) to  $x$ :

$$[x]_{\sim} = \{y \in X \mid y \sim x\}.$$

**Theorem** Equivalence classes of the relation  $\sim$  form a partition of the set  $X$ .

The set of all equivalence classes of  $\sim$  is denoted  $X/\sim$  and called the **quotient space** (or **factor space**) of  $X$  by the relation  $\sim$ .

In the case when the set  $X$  carries some structure (algebraic, geometric, analytic, etc.), this structure may (or may not) induce an analogous structure on the quotient space  $X/\sim$ .

## Examples of quotient spaces

- $X = \mathbb{Z}$ ,  $x \sim y$  if and only if  $x \equiv y \pmod{n}$ .

Equivalence class of an integer  $m$  is the congruence class modulo  $n$ ,  $[m]_{\sim} = [m]_n = m + n\mathbb{Z}$ . The quotient space  $\mathbb{Z}/\sim$  is  $\mathbb{Z}_n$ .

- $X = G$ , a group;  $x \sim y$  if and only if  $x \in yH$ , where  $H$  is a subgroup.

Equivalence class of an element  $g \in G$  is the coset of the subgroup  $H$ ,  $[g]_{\sim} = gH$ . The quotient space  $G/\sim$  is the set of all cosets of  $H$  in  $G$ . In this example, the quotient space is usually denoted  $G/H$ .

*Remark.* The first example is a particular case of the second, when  $G = \mathbb{Z}$  and  $H = n\mathbb{Z}$ . Hence  $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ .

## Quotient group

Let  $G$  be a nonempty set with a binary operation  $*$ , which is well defined (i.e., the closure axiom holds). Given an equivalence relation  $\sim$  on  $G$ , we say that the relation  $\sim$  is **compatible** with the operation  $*$  if for any  $g_1, g_2, h_1, h_2 \in G$ ,

$$g_1 \sim g_2 \text{ and } h_1 \sim h_2 \implies g_1 * h_1 \sim g_2 * h_2.$$

If this is the case, we can define an operation on the quotient space  $G/\sim$  by  $[g] \star [h] = [g * h]$  for all  $g, h \in G$ . Note that the operation  $\star$  is well defined: if  $[g'] = [g]$  and  $[h'] = [h]$  then  $[g' * h'] = [g * h]$ .

If the operation  $*$  is associative (commutative, resp.), then so is  $\star$ . If  $e$  is the identity element for  $*$ , then its equivalence class  $[e]$  is the identity element for  $\star$ . If  $h = g^{-1}$  in  $(G, *)$ , then  $[h] = [g]^{-1}$  in  $(G/\sim, \star)$ .

Thus, if  $(G, *)$  is a group then  $(G/\sim, \star)$  is also a group called the **quotient group**.

## Quotient group

**Question.** When is an equivalence relation  $\sim$  on a group  $G$  compatible with the operation?

**Theorem** Assume that the quotient space  $G/\sim$  is also a quotient group. Then

- (i)  $H = [e]_{\sim}$ , the equivalence class of the identity element, is a subgroup of  $G$ ,
- (ii)  $[g]_{\sim} = gH$  for all  $g \in G$ ,
- (iii)  $G/\sim = G/H$ ,
- (iv) the subgroup  $H$  is **normal**, which means that  $ghg^{-1} \in H$  for all  $h \in H$  and  $g \in G$ .

**Theorem** If  $H$  is a normal subgroup of a group  $G$ , then  $G/H$  is a quotient group.