

MATH 433
Applied Algebra

Lecture 1:
Division of integers.
Greatest common divisor.

Integer numbers

Positive integers: $\mathbb{P} = \{1, 2, 3, \dots\}$

Natural numbers: $\mathbb{N} = \{0, 1, 2, 3, \dots\}$

Integers: $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Arithmetic operations: addition, subtraction, multiplication, and division.

Addition and multiplication are well defined for the natural numbers \mathbb{N} . Subtraction is well defined for the integers \mathbb{Z} (only partially defined on \mathbb{N}).

Division by a nonzero number is well defined on the set of *rational numbers* \mathbb{Q} (only partially defined on \mathbb{Z} and \mathbb{N}).

Division of integer numbers

Let a and b be integers and $a \neq 0$. We say that a **divides** b or that b **is divisible by** a if $b = aq$ for some integer q . The integer q is called the **quotient** of b by a .

Notation: $a \mid b$ (a divides b)

$a \nmid b$ (a does not divide b)

Let a and b be integers and $a > 0$. Suppose that $b = aq + r$ for some integers q and r such that $0 \leq r < a$. Then r is the **remainder** and q is the (partial) **quotient** of b by a .

Note that $a \mid b$ means that the remainder is 0.

Ordering of integers

Integer numbers are ordered: for any $a, b \in \mathbb{Z}$ we have either $a < b$ or $b < a$ or $a = b$.

One says that an integer c lies between integers a and b if $a < c < b$ or $b < c < a$.

Well-ordering principle: any nonempty set of natural numbers has the smallest element.

As a consequence, any decreasing sequence of natural numbers is finite.

Remark. The well-ordering principle does not hold for all integers (there is no smallest integer).

Division theorem

Theorem Let a and b be integers and $a > 0$. Then the remainder and the quotient of b by a are well-defined. That is, $b = aq + r$ for some integers q and r such that $0 \leq r < a$.

Proof: First consider the case $b \geq 0$. Let

$$R = \{x \in \mathbb{N} \mid x = b - ay \text{ for some } y \in \mathbb{Z}\}.$$

The set R is not empty as $b = b - a \cdot 0 \in R$. Hence it has the smallest element r . We have $r = b - aq$ for some $q \in \mathbb{Z}$.

Consider the number $r - a$. Since $r - a < r$, it is not contained in R . But $r - a = (b - aq) - a = b - a(q + 1)$. It follows that $r - a$ is not natural, i.e., $r - a < 0$.

Thus $b = aq + r$, where q and r are integers and $0 \leq r < a$.

Now consider the case $b < 0$. In this case $-b > 0$.

By the above $-b = aq + r$ for some integers q and r such that $0 \leq r < a$. If $r = 0$ then $b = -aq = a(-q) + 0$.

If $0 < r < a$ then $b = -aq - r = a(-q - 1) + (a - r)$.

Greatest common divisor

Given two positive integers a and b , the **greatest common divisor** of a and b is the largest natural number that divides both a and b .

Notation: $\gcd(a, b)$ or simply (a, b) .

Example 1. $a = 12$, $b = 18$.

Natural divisors of 12 are 1, 2, 3, 4, 6, and 12.

Natural divisors of 18 are 1, 2, 3, 6, 9, and 18.

Common divisors are 1, 2, 3, and 6.

Thus $\gcd(12, 18) = 6$.

Notice that $\gcd(12, 18)$ is divisible by any other common divisor of 12 and 18.

Example 2. $a = 1356$, $b = 744$. $\gcd(a, b) = ?$

Euclidean algorithm

Lemma 1 If a divides b then $\gcd(a, b) = a$.

Lemma 2 If $a \nmid b$ and r is the remainder of b by a , then $\gcd(a, b) = \gcd(r, a)$.

Example 2. $a = 1356$, $b = 744$. $\gcd(a, b) = ?$

First we divide 1356 by 744: $1356 = 744 \cdot 1 + 612$.

Then divide 744 by 612: $744 = 612 \cdot 1 + 132$.

Then divide 612 by 132: $612 = 132 \cdot 4 + 84$.

Then divide 132 by 84: $132 = 84 \cdot 1 + 48$.

Then divide 84 by 48: $84 = 48 \cdot 1 + 36$.

Then divide 48 by 36: $48 = 36 \cdot 1 + 12$.

Then divide 36 by 12: $36 = 12 \cdot 3$.

Thus $\gcd(1356, 744) = \gcd(744, 612)$
 $= \gcd(612, 132) = \gcd(132, 84) = \gcd(84, 48)$
 $= \gcd(48, 36) = \gcd(36, 12) = 12$.