

MATH 433  
Applied Algebra

**Lecture 14:**  
**Functions.**  
**Relations.**

## Set theory

The primary notions of **set theory** are an **element** (an object that we can work with), a **set** (a collection of objects that we can work with), and **membership**. Namely, given an element  $x$  and a set  $S$ , we have either  $x \in S$  ( $x$  is a member of  $S$ ) or  $x \notin S$  ( $x$  is not a member of  $S$ ).

Any set is determined uniquely by its members (**axiom of extensionality**). Given sets  $S_1$  and  $S_2$ , we say that  $S_1$  is a **subset** of  $S_2$  (and write  $S_1 \subset S_2$ ) if every member of  $S_1$  is also a member of  $S_2$ . The axiom of extensionality can be rephrased as follows: for any sets  $S_1$  and  $S_2$ ,

$$S_1 = S_2 \iff S_1 \subset S_2 \text{ and } S_2 \subset S_1.$$

## Set theory

Set theory can provide the foundation for all of mathematics (though there are other ways as well).

The general idea is that every mathematical object is modeled as a set so that objects of the same kind are the same if and only if the corresponding sets are the same (but the same set can serve as a model for many objects of different kinds).

For example, one way to model nonnegative integers is as follows: 0 is the empty set  $\emptyset$ , 1 is  $\{\emptyset\}$ , 2 is  $\{\emptyset, \{\emptyset\}\}$ , 3 is  $\{\emptyset, \{\emptyset, \{\emptyset, \{\emptyset\}\}\}$ , and so on...

## Cartesian product

*Definition.* The **Cartesian product**  $X \times Y$  of two sets  $X$  and  $Y$  is the set consisting of all ordered pairs  $(x, y)$  such that  $x \in X$  and  $y \in Y$ .

The Cartesian square  $X \times X$  is also denoted  $X^2$ .

If the sets  $X$  and  $Y$  are finite, then

$|X \times Y| = |X| \cdot |Y|$ , where  $|S|$  denotes the number of elements in a set  $S$ .

*Remark.* An ordered pair  $(x, y)$  can be modeled as a set  $S_{x,y}$ , where  $S_{x,y} = \{x, \{x, y\}\}$  if  $x \neq y$  and  $S_{x,y} = \{x, \{x\}\}$  if  $x = y$ .

## Functions

A **function** (or **map**)  $f : X \rightarrow Y$  is an assignment: to each  $x \in X$  we assign an element  $f(x) \in Y$ .


*Definition.* A function  $f : X \rightarrow Y$  is **injective** (or **one-to-one**) if  $f(x') = f(x) \implies x' = x$ .

The function  $f$  is **surjective** (or **onto**) if for each  $y \in Y$  there exists at least one  $x \in X$  such that  $f(x) = y$ .

Finally,  $f$  is **bijective** if it is both surjective and injective. Equivalently, if for each  $y \in Y$  there is exactly one  $x \in X$  such that  $f(x) = y$ .

Suppose we have two functions  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$ . We say that  $g$  is the **inverse function** of  $f$  (denoted  $f^{-1}$ ) if  $y = f(x) \iff g(y) = x$  for all  $x \in X$  and  $y \in Y$ .

**Theorem** The inverse function  $f^{-1}$  exists if and only if  $f$  is bijective.



*Definition.* The **composition** of functions  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  is a function from  $X$  to  $Z$ , denoted  $g \circ f$ , that is defined by  $(g \circ f)(x) = g(f(x))$ ,  $x \in X$ .

$$X \xrightarrow{f} Y \xrightarrow{g} Z$$

*Properties of compositions:*

- If  $f$  and  $g$  are one-to-one, then  $g \circ f$  is also one-to-one.
- If  $g \circ f$  is one-to-one, then  $f$  is also one-to-one.
- If  $f$  and  $g$  are onto, then  $g \circ f$  is also onto.
- If  $g \circ f$  is onto, then  $g$  is also onto.
- If  $f$  and  $g$  are bijective, then  $g \circ f$  is also bijective.
- If  $f$  and  $g$  are invertible, then  $g \circ f$  is also invertible and  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$ .
- If  $\text{id}_Z$  denotes the identity function on a set  $Z$ , then  $f \circ \text{id}_X = f = \text{id}_Y \circ f$  for any function  $f : X \rightarrow Y$ .
- For any functions  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$ , we have  $g = f^{-1}$  if and only if  $g \circ f = \text{id}_X$  and  $f \circ g = \text{id}_Y$ .

## Relations

*Definition.* Let  $X$  and  $Y$  be sets. A **relation**  $R$  from  $X$  to  $Y$  is given by specifying a subset of the Cartesian product:  $S_R \subset X \times Y$ .

If  $(x, y) \in S_R$ , then we say that  $x$  **is related to**  $y$  (in the sense of  $R$  or by  $R$ ) and write  $xRy$ .

*Remarks.* • Usually the relation  $R$  is identified with the set  $S_R$ .

• In the case  $X = Y$ , the relation  $R$  is called a **relation on**  $X$ .



**Examples.** • “is equal to”

$$xRy \iff x = y$$

Equivalently,  $R = \{(x, x) \mid x \in X \cap Y\}$ .

• “is not equal to”

$$xRy \iff x \neq y$$

• “is mapped by  $f$  to”

$xRy \iff y = f(x)$ , where  $f : X \rightarrow Y$  is a function.

Equivalently,  $R$  is the graph of the function  $f$ .

• “is the image under  $f$  of”

(from  $Y$  to  $X$ )  $yRx \iff y = f(x)$ , where  $f : X \rightarrow Y$  is a function. If  $f$  is invertible, then  $R$  is the graph of  $f^{-1}$ .

• reversed  $R'$

$xRy \iff yR'x$ , where  $R'$  is a relation from  $Y$  to  $X$ .

• not  $R'$

$xRy \iff \text{not } xR'y$ , where  $R'$  is a relation from  $X$  to  $Y$ .

Equivalently,  $R = (X \times Y) \setminus R'$  (set difference).

## Relations on a set

- “is equal to”

$$xRy \iff x = y$$

- “is not equal to”

$$xRy \iff x \neq y$$

- “is less than”

$$X = \mathbb{R}, \quad xRy \iff x < y$$

- “is less than or equal to”

$$X = \mathbb{R}, \quad xRy \iff x \leq y$$

- “is contained in”

$X$  = the set of all subsets of some set  $Y$ ,

$$xRy \iff x \subset y$$

- “is congruent modulo  $n$  to”

$$X = \mathbb{Z}, \quad xRy \iff x \equiv y \pmod{n}$$

- “divides”

$$X = \mathbb{P}, \quad xRy \iff x|y$$