


MATH 614

Dynamical Systems and Chaos

**Lecture 17a:**  
**Morse-Smale diffeomorphisms.**



## Maps of the circle



$$T : S^1 \rightarrow S^1,$$

$T$  an orientation-preserving homeomorphism.

## Structurally stable maps of the circle



*Definition.* An orientation-preserving diffeomorphism  $f : S^1 \rightarrow S^1$  is **Morse-Smale** if it has rational rotation number and all of its periodic points are hyperbolic.

If  $\rho(f) = m/n$ , a reduced fraction, then all periodic points of  $f$  have period  $n$ . Hence the only periodic points of  $f^n$  are fixed points, alternately sinks and sources around the circle.

**Theorem** A Morse-Smale diffeomorphism of the circle is  $C^1$ -structurally stable.

**Theorem (The Closing Lemma)** Suppose  $f$  is a  $C^r$ -diffeomorphism of  $S^1$  with an irrational rotation number. Then for any  $\varepsilon > 0$  there exists a diffeomorphism  $g : S^1 \rightarrow S^1$  with a rational rotation number such that  $f$  and  $g$  are  $C^r$ - $\varepsilon$  close.

**Theorem (Kupka-Smale)** For any orientation-preserving  $C^r$ -diffeomorphism  $f$  of  $S^1$  and any  $\varepsilon > 0$  there exists a Morse-Smale diffeomorphism that is  $C^r$ - $\varepsilon$  close to  $f$ .