

MATH 614

Dynamical Systems and Chaos

Lecture 32:

The Julia and Fatou sets.

The Julia set

Suppose $P : U \rightarrow U$ is a holomorphic map, where U is a domain in \mathbb{C} , the entire plane \mathbb{C} , or the Riemann sphere $\overline{\mathbb{C}}$.

Definition. The **Julia set** $J(P)$ of P is the closure of the set of repelling periodic points of P .

Examples. • $L_2(z) = 2z$.

$$J(L_2) = \{0\}.$$

• $L_{1/2}(z) = z/2$.

$J(L_{1/2}) = \{\infty\}$ since $L_{1/2} = H \circ L_2 \circ H^{-1}$, where $H(z) = 1/z$.

• $L_{1,1}(z) = z + 1$.

$$J(L_{1,1}) = \emptyset.$$

• $Q_0(z) = z^2$.

$$J(Q_0) = \{z \in \mathbb{C} : |z| = 1\}.$$

• $Q_{-2}(z) = z^2 - 2$.

$$J(Q_{-2}) = [-2, 2]. \quad \text{Note that } Q_{-2}(2 \cos \alpha) = 2 \cos(2\alpha).$$

Invariance

Proposition 1 The Julia set of a holomorphic map $P : U \rightarrow U$ is invariant: $P(J(P)) \subset J(P)$.

Proof: Let $z \in J(P)$. There are repelling periodic points z_1, z_2, \dots of P such that $z_n \rightarrow z$ as $n \rightarrow \infty$. By continuity, $P(z_n) \rightarrow P(z)$ as $n \rightarrow \infty$. Clearly, $P(z_1), P(z_2), \dots$ are also repelling periodic points of P .

Proposition 2 $P(J(P)) = J(P)$.

Proof: Let $z \in J(P)$ and $z_n \rightarrow z$ as $n \rightarrow \infty$, where z_1, z_2, \dots are repelling periodic points of P . Then there are repelling periodic points w_1, w_2, \dots such that $P(w_n) = z_n$. The sequence w_1, w_2, \dots is bounded, hence there is a converging subsequence: $w_{n_k} \rightarrow w$ as $k \rightarrow \infty$. Then $z_{n_k} = P(w_{n_k}) \rightarrow P(w)$. We have $w \in J(P)$ and $P(w) = z$.

Remark. This argument is incomplete.

Normal family

Let \mathcal{F} be a collection of holomorphic functions $F : U \rightarrow \mathbb{C}$ defined in a domain $U \subset \mathbb{C}$.

Definition. The collection \mathcal{F} is a **normal family** in U if every sequence F_1, F_2, \dots of functions from \mathcal{F} has a subsequence F_{n_1}, F_{n_2}, \dots ($1 \leq n_1 < n_2 < \dots$) which either

- (i) converges uniformly on compact subsets of U , or
- (ii) converges uniformly to ∞ on U .

The condition (i) means that there exists a function $f : U \rightarrow \mathbb{C}$ such that for any compact set $D \subset U$ we have

$$\sup_{z \in D} |F_{n_k}(z) - f(z)| \rightarrow 0 \text{ as } k \rightarrow \infty.$$

The function f is going to be continuous.

The condition (ii) means that for any $R > 0$ there exists an integer $K > 0$ such that $|F_{n_k}(z)| > R$ for all $k \geq K$ and $z \in U$.

The Fatou set

Let \mathcal{F} be a collection of holomorphic functions defined in a domain $U \subset \overline{\mathbb{C}}$.

We say that the collection \mathcal{F} is **normal at** a finite point $z \in U$ if it is a normal family in some neighborhood of z . In the case $\infty \in U$, we say \mathcal{F} is **normal at infinity** if the collection of functions $G(z) = F(1/z)$, $F \in \mathcal{F}$ is normal at 0.

Definition. The **Fatou set** $S(P)$ of a holomorphic map $P : U \rightarrow U$ is the set of all points $z \in U$ such that the family of iterates $\{P^n\}_{n \geq 1}$ is normal at z .

By definition, the Fatou set is open.

Let \mathcal{F} be a collection of holomorphic functions defined in a domain $U \subset \overline{\mathbb{C}}$.

Theorem (Arzelà-Ascoli) Suppose the functions in \mathcal{F} are uniformly bounded and their derivatives are uniformly bounded.

Then any sequence F_1, F_2, \dots of functions from \mathcal{F} has a subsequence F_{n_1}, F_{n_2}, \dots which converges uniformly on compact subsets of U .

Corollary If the iterates of a holomorphic transformation P and their derivatives are uniformly bounded in a neighborhood of a point z , then $z \in S(P)$.

Theorem (Weierstrass) Let F_1, F_2, \dots be holomorphic functions in a domain U . Assume that the sequence F_1, F_2, \dots converges uniformly on compact subsets of U .

Then the limit function F is holomorphic in U and, moreover, the sequence of derivatives F'_1, F'_2, \dots converges to F' uniformly on compact subsets of U .

Corollary Let $z \in \mathbb{C}$. Assume that there exists a sequence of iterates P^{n_1}, P^{n_2}, \dots such that $P^{n_k}(z) \not\rightarrow \infty$ while $(P^{n_k})'(z) \rightarrow \infty$ as $k \rightarrow \infty$. Then $z \notin S(P)$.

The Fatou set and periodic points

Proposition 1 Attracting periodic points of P belong to $S(P)$.

Proof: Suppose z_0 is an attracting periodic point of period n . Then $P^n(z_0) = z_0$ and $|P^n(z) - z_0| \leq \mu|z - z_0|$ for some $0 < \mu < 1$ and all z close enough to z_0 . Hence there is a neighborhood D of z_0 such that the functions $P^n, P^{2n}, P^{3n}, \dots$ converge to the constant z_0 uniformly on D . Then for any integer $k \geq 1$ the functions $P^{n+k}, P^{2n+k}, P^{3n+k}, \dots$ converge to the constant $P^k(z_0)$ uniformly on D .

It follows that the sequence P, P^2, P^3, \dots is normal at z_0 .

The Fatou set and periodic points

Proposition 2 Repelling periodic points of P do not belong to $S(P)$.

Proof: Suppose z is a repelling periodic point of period n . Then $P^n(z) = z$ and $|(P^n)'(z)| > 1$. For any integer $k \geq 1$ we have $P^{nk}(z) = z$ and $(P^{nk})'(z) = ((P^n)'(z))^k$. As a consequence, $P^{nk}(z) \not\rightarrow \infty$ while $(P^{nk})'(z) \rightarrow \infty$ as $k \rightarrow \infty$. By the above, $z \notin S(P)$.

Corollary The Julia set and the Fatou set of P are disjoint.

The Fatou set and periodic points

Neutral periodic points of P may or may not belong to $S(P)$.

Examples. • $P(z) = e^{i\alpha}z$, where $\alpha \in \mathbb{R}$.

The neutral fixed point 0 belongs to the Fatou set $S(P)$. Indeed, $P^n(z) = e^{in\alpha}z$ and $(P^n)'(z) = e^{in\alpha}$ for all $z \in \mathbb{C}$ and $n = 1, 2, \dots$. Therefore all iterates of P and their derivatives are uniformly bounded in a neighborhood of 0.

• $P(z) = z + z^2$.

The neutral fixed point 0 does not belong to the Fatou set $S(P)$. Indeed, for any ε , $0 < \varepsilon < 1$, we have $P^n(-\varepsilon) \rightarrow 0$ as $n \rightarrow \infty$ while $P^n(\varepsilon) \rightarrow \infty$ as $n \rightarrow \infty$. It follows that the iterates of P are not normal at 0.

Invariance

Proposition The Fatou set of P is completely invariant under this map: $P(S(P)) \subset S(P)$ and $P^{-1}(S(P)) \subset S(P)$.

Proof: Suppose $P(w) = z$. We have to show that the family P, P^2, P^3, \dots is normal at z if and only if it is normal at w .

Indeed, let f be a nonconstant holomorphic function such that $f(w) = z$. Then P, P^2, P^3, \dots is normal at z if and only if the family $P \circ f, P^2 \circ f, P^3 \circ f, \dots$ is normal at w .