

MATH 614

Dynamical Systems and Chaos

**Lecture 36:**  
**Invariant measure.**

## Ergodic theory

Topological dynamics is the study of continuous transformations.

Smooth dynamics is the study of smooth transformations.

Holomorphic dynamics is the study of holomorphic transformations.

**Ergodic theory** (a.k.a. metric theory of dynamical systems) is the study of **measure-preserving** transformations.

The **measure** is an abstract concept that generalizes the notions of length, area, and volume.

## Examples


- Bijective self-map  $F : X \rightarrow X$ .

Any set  $E \subset X$  is mapped onto a set with the same number of elements.

- Translation of the real line.

$F : \mathbb{R} \rightarrow \mathbb{R}$ ,  $F(x) = x + x_0$ . Any interval is mapped onto an interval of the same length.


- Rotation of the circle.



Any arc is mapped onto an arc of the same length.

## Non-continuous example

- Interval exchange transformation.



An **interval exchange transformation**  $F : I \rightarrow I$  of an interval  $I$  is defined by cutting the interval into several subintervals and then rearranging them by translation. The image of any subinterval  $I_0 \subset I$  consists of one or several intervals whose total length equals the length of  $I_0$ .

Note that the transformation  $F$  is not well defined at the cutting points. Consequently, the orbit under  $F$  is not defined for a finite or countable set of points which may be dense in  $I$ . However this is not a concern as in ergodic theory sets of zero measure can be neglected.

- Motion of the Euclidean plane.

Any domain is mapped onto a domain of the same area.

- Linear transformation  $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ .

$L(\mathbf{x}) = A\mathbf{x}$ , where  $A$  is a  $2 \times 2$  matrix. The image of any domain of area  $\alpha$  has area  $\alpha |\det A|$ . In the case  $\det A = \pm 1$ , the map  $L$  is area-preserving.

- Translation of the torus.

$F : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ ,  $F(\mathbf{x}) = \mathbf{x} + \mathbf{x}_0$ . This is the quotient of a translation of the Euclidean plane under the natural projection  $\pi : \mathbb{R}^2 \rightarrow \mathbb{T}^2$ .

- Toral automorphism.

$F : \mathbb{T}^2 \rightarrow \mathbb{T}^2$  ( $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ ),  $F(\mathbf{x}) = A\mathbf{x}$ , where  $A$  is a  $2 \times 2$  matrix with integer entries and  $\det A = \pm 1$ . This is the quotient of an area-preserving linear map under the natural projection  $\pi : \mathbb{R}^2 \rightarrow \mathbb{T}^2$ .

## Example with continuous time

- Area-preserving flow.

Consider an autonomous system of two ordinary differential equations of the first order


$$\begin{cases} \dot{x} = g_1(x, y), \\ \dot{y} = g_2(x, y), \end{cases}$$

where  $g_1, g_2$  are differentiable functions defined in a domain  $D \subset \mathbb{R}^2$ . In vector form,  $\dot{\mathbf{v}} = G(\mathbf{v})$ , where  $G : D \rightarrow \mathbb{R}^2$  is a vector field. Assume that for any  $\mathbf{x} \in D$  the initial value problem  $\dot{\mathbf{v}} = G(\mathbf{v})$ ,  $\mathbf{v}(0) = \mathbf{x}$  has a unique solution  $\mathbf{v}_{\mathbf{x}}(t)$ ,  $t \in \mathbb{R}$ . Then the system of ODEs gives rise to a dynamical system with continuous time  $F^t : D \rightarrow D$ ,  $t \in \mathbb{R}$  defined by  $F^t(\mathbf{x}) = \mathbf{v}_{\mathbf{x}}(t)$  for all  $\mathbf{x} \in D$  and  $t \in \mathbb{R}$ .

The flow  $\{F^t\}$  is area-preserving if and only if  $\nabla \cdot G = \partial g_1 / \partial x + \partial g_2 / \partial y = 0$  in  $D$ .

## Non-invertible example

- Doubling map  $F : S^1 \rightarrow S^1$ .



If  $S^1 = \mathbb{R}/\mathbb{Z}$ , then  $F(x) = 2x$  for all  $x \in S^1$ . For any arc  $\gamma = (\omega_1, \omega_2)$ ,  $0 \leq \omega_1 < \omega_2 \leq 1$ , of length  $\alpha = \omega_2 - \omega_1$  the image  $F(\gamma)$  is an arc of length  $2\alpha$  or the entire circle.

However the preimage  $F^{-1}(\gamma)$  consists of two disjoint arcs  $(\frac{1}{2}\omega_1, \frac{1}{2}\omega_2)$  and  $(\frac{1}{2}\omega_1 + \frac{1}{2}, \frac{1}{2}\omega_2 + \frac{1}{2})$  of length  $\alpha/2$  so that  $F^{-1}(\gamma)$  has the same length measure as  $\gamma$ .

## Measure-preserving transformation

*Definition.* A **measured space** is a triple  $(X, \mathcal{B}, \mu)$ , where  $X$  is a set,  $\mathcal{B}$  is a collection of subsets of  $X$ , and  $\mu$  is a function  $\mu : \mathcal{B} \rightarrow [0, \infty]$ . Elements of  $\mathcal{B}$  are referred to as **measurable sets**. The function  $\mu$  is called the **measure** on  $X$ .

A mapping  $T : X \rightarrow X$  is called **measurable** if preimage of any measurable set under  $T$  is also measurable:  $E \in \mathcal{B} \implies T^{-1}(E) \in \mathcal{B}$ .

A measurable mapping  $T : X \rightarrow X$  is called **measure-preserving** if for any  $E \in \mathcal{B}$  one has  $\mu(T^{-1}(E)) = \mu(E)$ .



## Algebra of sets

*Definition.* A collection  $\mathcal{B}$  of subsets of a set  $X$  is called an **algebra** of sets if  $\mathcal{B}$  is closed under taking unions  $B_1 \cup B_2$ , intersections  $B_1 \cap B_2$ , complements  $X \setminus B$ , and if  $\mathcal{B}$  contains the empty set and the entire set  $X$ .

The algebra  $\mathcal{B}$  is also closed under taking finite unions  $B_1 \cup B_2 \cup \dots \cup B_n$ , finite intersections  $B_1 \cap B_2 \cap \dots \cap B_n$ , set differences  $B_1 \setminus B_2 = B_1 \cap (X \setminus B_2)$ , and symmetric differences  $B_1 \Delta B_2 = (B_1 \setminus B_2) \cup (B_2 \setminus B_1)$ .

For any subset  $B \subset X$  let  $\chi_B : X \rightarrow \{0, 1\}$  denote the characteristic function of  $B$ :  $\chi_B(x) = 1$  if  $x \in B$  and  $\chi_B(x) = 0$  otherwise. Then  $\chi_X = 1$ ,  $\chi_\emptyset = 0$ ,  
 $\chi_{B_1 \cap B_2} = \chi_{B_1} \chi_{B_2}$ ,  $\chi_{B_1 \cup B_2} = \chi_{B_1} + \chi_{B_2}$  if  $B_1 \cap B_2 = \emptyset$ ,  
 $\chi_{B_1 \setminus B_2} = \chi_{B_1} - \chi_{B_2}$  if  $B_2 \subset B_1$ , and  
 $\chi_{B_1 \Delta B_2} = \chi_{B_1} + \chi_{B_2} \pmod{2}$ .

## $\sigma$ -algebra

A standard requirement for a measured space  $(X, \mathcal{B}, \mu)$  is that  $\mathcal{B}$  be a  $\sigma$ -algebra.

*Definition.* An algebra of sets is called a  $\sigma$ -**algebra** if it is closed under taking countable unions.

*Examples of  $\sigma$ -algebras:* •  $\{\emptyset, X\}$ ;

- all subsets of  $X$  ( $2^X$ );
- all finite and countable subsets of  $X$  and their complements.

**Proposition** Given a collection  $S$  of subsets of  $X$ , there exists a minimal  $\sigma$ -algebra of subsets of  $X$  that contains  $S$ .

Suppose  $X$  is a topological space. The **Borel**  $\sigma$ -algebra  $\mathcal{B}(X)$  is the minimal  $\sigma$ -algebra that contains all open subsets of  $X$ . Elements of  $\mathcal{B}(X)$  are called **Borel sets**. A mapping  $F : X \rightarrow X$  is measurable relative to  $\mathcal{B}(X)$  if and only if the preimage of any open set is Borel. In particular, each continuous map is measurable.

## $\sigma$ -additive measure

*Definition.* Suppose  $\mathcal{B}$  is an algebra of subsets of a set  $X$ . A function  $\mu : \mathcal{B} \rightarrow [0, \infty]$  is an **additive measure** if  $\mu(\emptyset) = 0$  and, for any disjoint sets  $A_1, A_2, \dots, A_n \in \mathcal{B}$ ,

$$\mu\left(\bigcup_{k=1}^n A_k\right) = \sum_{k=1}^n \mu(A_k).$$

In the case  $\mathcal{B}$  is a  $\sigma$ -algebra, the additive measure  $\mu$  is  **$\sigma$ -additive** if for any disjoint sets  $A_1, A_2, \dots$  from  $\mathcal{B}$ ,

$$\mu\left(\bigcup_{k=1}^{\infty} A_k\right) = \sum_{k=1}^{\infty} \mu(A_k).$$

The measure  $\mu$  is **finite** if  $\mu(X) < \infty$ .  $\mu$  is  **$\sigma$ -finite** if  $X = \bigcup_{k=1}^{\infty} X_k$ , where  $\mu(X_k) < \infty$  for all  $k$ .

Another standard requirement for a measured space  $(X, \mathcal{B}, \mu)$  is that  $\mu$  be a  $\sigma$ -additive measure and also be finite or  $\sigma$ -finite.

*Definition.* A normalized invariant **mean** on  $\mathbb{Z}$  is a function  $m : 2^{\mathbb{Z}} \rightarrow [0, \infty)$  such that

- $m(\emptyset) = 0$ ,  $m(\mathbb{Z}) = 1$ ;
- if  $A_1, A_2, \dots, A_k$  are disjoint subsets of  $\mathbb{Z}$  then  $m(A_1 \cup \dots \cup A_k) = m(A_1) + \dots + m(A_k)$ ;
- $m(n + S) = m(S)$  for all  $n \in \mathbb{Z}$  and  $S \subset \mathbb{Z}$ .

The mean  $m$  is a finite, additive measure on  $\mathbb{Z}$ .

Note that  $m(\{n\})$  is the same for all  $n \in \mathbb{Z}$ .

Since  $m(\mathbb{Z}) < \infty$ , it follows that  $m(\{n\}) = 0$ .

Besides, it follows that  $m$  is not  $\sigma$ -additive.

**Theorem (Banach)** There exists a normalized invariant mean on  $\mathbb{Z}$ .

That is, the group  $\mathbb{Z}$  is **amenable**.