

Power Series, Taylor Series and Analytic Functions (section 5.1)

DEFINITION 1. A power series about $x = x_0$ (or centered at $x = x_0$), or just **power series**, is any series that can be written in the form

$$\sum_{n=0}^{\infty} a_n(x - x_0)^n,$$

where x_0 and a_n are numbers.

DEFINITION 2. A power series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ is said to converge at a point x if the limit $\lim_{m \rightarrow \infty} \sum_{n=0}^m a_n(x - x_0)^n$ exists and finite.

REMARK 3. A power series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ always converges at $x = x_0$.

EXAMPLE 4. For what x the power series $\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \dots$ converges ?

$$\sum_{n=0}^m x^n =$$

$$\text{If } |x| < 1$$

$$\text{If } |x| > 1$$

Absolute Convergence: The series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ is said to converge absolutely at x if

$$\sum_{n=0}^{\infty} |a_n| |x - x_0|^n \text{ converges.}$$

If a series converges absolutely then it converges (but in general not vice versa).

EXAMPLE 5. The series $\sum_{n=1}^{\infty} \frac{x^n}{n}$ converges at $x = -1$, but it doesn't converges absolutely there:

$$1 - \frac{1}{2} + \frac{1}{3} - \dots = \ln 2$$

but the series of absolute values is the so-called harmonic series

$$1 + \frac{1}{2} + \frac{1}{3} + \dots$$

and it is divergent.

Fact: If the series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ converges absolutely at $x = x_1$ then it converges absolutely for all x such that $|x - x_0| < |x_1 - x_0|$

This immediately implies the following:

THEOREM 6. For a given power series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ there are only 3 possibilities:

1. The series converges only for $x = x_0$.
2. The series converges for all x .
3. There is $R > 0$ such that the series converges if $|x - x_0| < R$ and diverges if $|x - x_0| > R$. We call such R the **radius of convergence**.

REMARK 7. In case 1 of the theorem we say that $R = 0$ and in case 2 we say that $R = \infty$

EXAMPLE 8. What is the radius of convergence of the geometric power series $\sum_{n=0}^{\infty} x^n$?

How to find Radius of convergence: If $a_n \neq 0$ for any n and $\lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$ exists, then

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$$

More generally,

$$R = \frac{1}{\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|}},$$

the Cauchy-Hadamard formula.

EXAMPLE 9. Find the radius of convergence of the power series $\sum_{n=0}^{\infty} \frac{n^2}{3^n}(x + 1)^n$.

The Taylor series for $f(x)$ about $x = x_0$

Assume that f has derivatives of any order at $x = x_0$. Then for any m

$$f(x) = \sum_{n=0}^m \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n + \frac{f^{(m+1)}(c)}{(m+1)!} (x - x_0)^{m+1}$$

where c is a number between x and x_0 . The remainder converges to zero at least as fast as $(x - x_0)^{m+1}$ when $x \rightarrow x_0$.

Formally we can consider the following power series:

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n,$$

the *Taylor series of the function f about x_0* .

The Taylor series may converge and may not converge in a neighborhood of x_0 and even if it converges for any x close to x_0 it may not converge to $f(x)$.

DEFINITION 10. *The function f is called **analytic** at the point x_0 if there exist a power series*

$$\sum_{n=0}^{\infty} a_n (x - x_0)^n$$

such that

$$f(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^n$$

for all x sufficiently close to x_0 .

In this case, one can show that a_n must be equal to $\frac{f^{(n)}(x_0)}{n!}$. This implies:

The function f is analytic at the point x_0 if its Taylor series

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

converges to $f(x)$ for all x sufficiently closed to x_0 .

EXAMPLE 11. *From Example 3 it follows that $f(x) = \frac{1}{1-x}$ is analytic at 0. In general, it is analytic at any $x_0 \neq 1$*

EXAMPLE 12. *More generally, any rational function $f(x) = \frac{Q(x)}{P(x)}$, where $P(x)$ and $Q(x)$ are polynomials without common linear factors (the latter can be always assumed, because the common factors can be canceled) is analytic at all points except zeros of the denominator $P(x)$*

EXAMPLE 13. *The functions e^x , $\sin x$ and $\cos x$ are analytic at any x . Here are their Taylor series at 0:*

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$$

REMARK 14. *Not any function having derivatives of any order at any point is analytic. For example, take*

$$f(x) = \begin{cases} e^{-\frac{1}{x}}, & x > 0 \\ 0, & x \leq 0 \end{cases}.$$

Term by term differentiation If f is analytic at a point x_0 , i.e. $f(x) = \sum_{n=0}^{\infty} a_n(x - x_0)^n$ for x sufficiently close to x_0 , then $f'(x)$ is also analytic at x_0 and

$$f'(x) = \sum_{n=1}^{\infty} n a_n (x - x_0)^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} (x - x_0)^n$$

In other words, the derivative of a (convergent) power series is obtained by term by term differentiation of the series.

EXAMPLE 15. *What is the Taylor expansion of $f''(x)$?*

$$\begin{aligned} f''(x) &= \left(\sum_{n=0}^{\infty} (n+1) a_{n+1} (x - x_0)^n \right)' = \sum_{n=1}^{\infty} (n+1) n a_{n+1} (x - x_0)^{n-1} = \\ &\quad \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} (x - x_0)^n. \end{aligned}$$