

Math 151 Fall 2010 Exam I Solutions-Form B

1. C: A force $\vec{F} = 2\vec{i} + 6\vec{j}$ moves an object from the point $P(1, 3)$ to the point $Q(3, 7)$. How much work is done if the force is measured in pounds and the distance is measured in feet? Here, $\vec{F} = \langle 2, 6 \rangle$ and $\vec{D} = \overrightarrow{PQ} = \langle 2, 4 \rangle$. Hence the work done is

$$W = \vec{F} \cdot \vec{D} = \langle 2, 6 \rangle \cdot \langle 2, 4 \rangle = 28 \text{ foot pounds.}$$

2. D: If $\triangle ABC$ is an equilateral triangle with sides of length 2, compute $\overrightarrow{BA} \cdot \overrightarrow{BC}$.

$\overrightarrow{BA} \cdot \overrightarrow{BC} = |\overrightarrow{BA}| |\overrightarrow{BC}| \cos \theta$. We are given $|\overrightarrow{BA}| = 2$, $|\overrightarrow{BC}| = 2$, and since the triangle is equilateral, $\theta = 60^\circ$. Hence $\overrightarrow{BA} \cdot \overrightarrow{BC} = |\overrightarrow{BA}| |\overrightarrow{BC}| \cos \theta = (2)(2) \cos 60^\circ = 2$.

3. A: The parametric equations $x = 3 + \sin t$, $y = -2 + \cos t$ describe a circle since $x - 3 = \sin t$ and $y + 2 = \cos t$ and therefore $(x - 3)^2 + (y + 2)^2 = 1$.

4. E: What is the slope of the line that passes through the point $(4, 7)$ and is perpendicular to the vector $\langle -5, 6 \rangle$? The direction vector is $\langle -5, 6 \rangle^\perp = \langle -6, -5 \rangle$. Thus the slope of the line with direction vector $\langle -6, -5 \rangle$ is $\frac{5}{6}$.

5. B: Find the vector \vec{a} that has magnitude $|\vec{a}| = 12$ and makes an angle of 300° with the positive x -axis. Since $300^\circ = 360^\circ - 60^\circ$, our reference angle is 60° in quadrant IV. Thus

$$\vec{a} = \langle 12 \cos 60^\circ, -12 \sin 60^\circ \rangle = \langle 6, -6\sqrt{3} \rangle.$$

6. A: Which of the following intervals contains a solution to the equation $x^3 + 2x + 2 = 7$? First, note $f(x) = x^3 + 2x + 2$ is continuous, so we will apply the Intermediate Value Theorem. Since $f(1) = 5 < 7$ and $f(2) = 14 > 7$, there is a solution to $f(c) = 7$ on the interval $[1, 2]$.

7. D: Find all vertical asymptotes for $f(x) = \frac{x^2 - 1}{x^3 + x^2}$. First by simplifying $f(x)$, we find that

$$f(x) = \frac{(x+1)(x-1)}{x^2(x+1)} = \frac{x-1}{x^2}, \text{ provided } x \neq -1$$

and $x \neq 0$. Thus the only vertical asymptote occurs at $x = 0$.

8. E: Find the equation of the tangent line to the graph of $f(x) = \frac{x}{1+2x}$ at $x = 1$. The slope of the tangent line at $x = 1$ is $m = f'(1)$. Using the quotient rule, we find that $f'(x) = \frac{1}{(1+2x)^2}$. Thus $m = \frac{1}{9}$. The line is tangent to the curve at the point $(1, \frac{1}{3})$. Thus the equation of the tangent line is $y - \frac{1}{3} = \frac{1}{9}(x - 1)$.

$$\begin{aligned} 9. \text{ A: } \lim_{t \rightarrow \infty} \frac{6t^2 + 3t}{(3-t)(2t+4)} &= \lim_{t \rightarrow \infty} \frac{6t^2 + 3t}{-2t^2 + 2t + 12} \\ &= \lim_{t \rightarrow \infty} \frac{6 + 3/t}{-2 + 2/t + 12/t^2} = -3 \end{aligned}$$

10. A: Find the value of a and b that makes

$$f(x) = \begin{cases} -x + a & \text{if } x \leq 1 \\ bx^2 + 3 & \text{if } x > 1 \end{cases} \text{ continuous and differentiable at } x = 1.$$

For f to be continuous at $x = 1$, we need for $\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^-} f(x)$. Thus $b+3 = -1+a$. For f to be differentiable at $x = 1$, we need for $\lim_{x \rightarrow 1^+} f'(x) = \lim_{x \rightarrow 1^-} f'(x)$. Thus $2b = -1$. This yields $b = -\frac{1}{2}$ and therefore $a = \frac{7}{2}$.

11. Find the distance from the point $P(1, 7)$ to the line $y = 2x + 3$. First we will choose any two points on the line.

Let $A(0, 3)$ and $B(1, 5)$. Define $\vec{a} = \overrightarrow{AB} = \langle 1, 2 \rangle$ and $\vec{b} = \overrightarrow{AP} = \langle 1, 4 \rangle$. Then the distance from the point $P(1, 7)$ to the line $y = 2x + 3$ is

$$d = \text{comp}_{\vec{a}^\perp} \vec{b} = \frac{\vec{a}^\perp \cdot \vec{b}}{|\vec{a}^\perp|} = \frac{\langle -2, 1 \rangle \cdot \langle 1, 4 \rangle}{|\langle -2, 1 \rangle|} = \frac{2}{\sqrt{5}}$$

12. Find the coordinates of the intersection point of the lines L1 and L2 given below.

$$\text{L1: } x = -1 - 2t, y = 2 + t$$

$$\text{L2: } x = 15 - 3w, y = 3 + 6w$$

We find the intersection of the lines by solving the equations

$$(i) -1 - 2t = 15 - 3w \text{ and } (ii) 2 + t = 3 + 6w.$$

Solving (ii) for t yields $t = 1 + 6w$. Substituting this into (i) for t yields $-1 - 2(1 + 6w) = 15 - 3w$. Solving this for w gives $w = -2$, and therefore $t = -11$. Now to find the intersection point, we can either substitute $t = -11$ into L1 or substitute $w = -2$ into L2. Both yield the point $(21, -9)$.

13. For each of the following limits, either calculate the limit, if it exists, or else explain why the limit does not exist.

$$(i) \lim_{x \rightarrow 0^+} \frac{x^2 + 2x}{|x|} = \lim_{x \rightarrow 0^+} \frac{x(x+2)}{x} = 2$$


$$(ii) \lim_{x \rightarrow 0^-} \frac{x^2 + 2x}{|x|} = \lim_{x \rightarrow 0^-} \frac{x(x+2)}{-x} = -2$$

(iii) $\lim_{x \rightarrow 0} \frac{x^2 + 2x}{|x|}$ does not exist since the left and right hand limits at 0 are not equal.

14. Using the *definition of the derivative*, find $f'(x)$ for $f(x) = \frac{4}{x+1}$.

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{4}{x+h+1} - \frac{4}{x+1}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{4}{x+h+1} - \frac{4}{x+1}}{h} \cdot \frac{(x+h+1)(x+1)}{(x+h+1)(x+1)} \\ &= \lim_{h \rightarrow 0} \frac{4(x+1) - 4(x+h+1)}{h(x+h+1)(x+1)} \\ &= \lim_{h \rightarrow 0} \frac{-4h}{h(x+h+1)(x+1)} \\ &= \frac{-4}{(x+1)^2} \end{aligned}$$

15. The horizontal line through the point $(3, 1)$ is tangent to the parabola $y = x^2 + 1$. We see from the figure below that there is a second line tangent to the parabola at $(a, a^2 + 1)$ that also passes through the point $(3, 1)$. Find the value of a .



The slope of the tangent line to the curve $y = x^2 + 1$ at $x = a$ is $m_1 = 2a$. This tangent line can also be viewed as a line passing through the points $(3, 1)$ and $(a, a^2 + 1)$, and therefore the slope is also

$$m_2 = \frac{a^2 + 1 - 1}{a - 3}. \text{ Thus to find } a \text{ we will solve } m_1 = m_2:$$

$$2a = \frac{a^2}{a-3} \Rightarrow 2a(a-3) = a^2 \Rightarrow 2a^2 - 6a = a^2, \text{ thus } a = 0 \text{ or } a = 6. \text{ Clearly from the picture provided, } a > 0, \text{ hence } a = 6.$$

$$\begin{aligned} 16. (a) \text{ Find } \lim_{x \rightarrow 2} \frac{x - \sqrt{3x-2}}{x^2 - 4} \\ &= \lim_{x \rightarrow 2} \frac{(x - \sqrt{3x-2})(x + \sqrt{3x-2})}{(x^2 - 4)(x + \sqrt{3x-2})} \\ &= \lim_{x \rightarrow 2} \frac{x^2 - (3x-2)}{(x+2)(x-2)(x + \sqrt{3x-2})} \\ &= \lim_{x \rightarrow 2} \frac{(x-2)(x-1)}{(x+2)(x-2)(x + \sqrt{3x-2})} \\ &= \lim_{x \rightarrow 2} \frac{x-1}{(x+2)(x + \sqrt{3x-2})} = \frac{1}{16} \end{aligned}$$

- (b) Is there a value of a for which $\lim_{x \rightarrow -3} \frac{x^2 + ax + a + 5}{x^2 - 2x - 15}$ exists? If so, find the value of a . If not, explain why.

Note that $\lim_{x \rightarrow -3} \frac{x^2 + ax + a + 5}{x^2 - 2x - 15} = \lim_{x \rightarrow -3} \frac{x^2 + ax + a + 5}{(x-5)(x+3)}$. Hence in order for this limit to exist, the numerator must have a factor of $(x+3)$ (thereby eliminating the division by zero). Thus the numerator evaluated at $x = -3$ must be zero:

$$(-3)^2 + a(-3) + a + 5 = 0 \Rightarrow 14 - 2a = 0, \text{ thus } a = 7.$$