

Fall 2010 Math 152

Exam III Version B Solutions

1. **D** We need $\mathbf{a} \cdot \mathbf{b} = 0$.

$$(t+2)(t-2) + (t)(t+1) + t^2 = 3t^2 + t - 4 = 0,$$

$$(3t+4)(t-1) = 0, \text{ so } t = -\frac{4}{3} \text{ or } t = 1.$$

2. **B** Apply the Ratio Test to (I): $\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+3)3^{n+1}}{2^{2(n+1)+1}} \cdot \frac{2^{2n+1}}{(n+2)3^n} = \frac{(n+3)3^{n+1}}{2^{2n+3}} \cdot \frac{2^{2n+1}}{(n+2)3^n} = \frac{3(n+3)}{4(n+2)}$, which approaches $\frac{3}{4} < 1$, so (I) is **absolutely convergent**. (II) is an Alternating Series, but $|a_n| = \frac{n^3+4}{2n^2} \rightarrow \infty$, which means a_n is divergent (hence not approaching 0), so (II) is **divergent by the Test for Divergence**.

3. **E** $f(x)$ is a geometric sum with $a = 1, r = -4x^2$, so $f(x) = \sum_{n=0}^{\infty} (-4x^2)^n = 1 - 4x^2 + 16x^4 - 64x^6 + \dots$

4. **A** The power series has a radius of convergence of at least 4, so it is certain that the series is convergent for all $x \in (-4, 4]$ (convergence/divergence is unknown for all other values of x). Only $x = -2$ is in this interval, therefore, we know for certain that $\sum_{n=0}^{\infty} c_n(-2)^n$ is convergent.

5. **D** $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$, so $e^{-x^2} = \sum_{n=0}^{\infty} \frac{(-x^2)^n}{n!} = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{n!}$

6. **E** $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} = \frac{8}{\sqrt{3}\sqrt{26}} = \frac{8}{\sqrt{78}}$

7. **B** Apply the Ratio Test: $\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{4^{n+1}(x-2)^{n+1}}{(n+1)!} \cdot \frac{n!}{4^n(x-2)^n} \right| = \frac{4|x-2|}{n+1}$, which approaches 0 (hence < 1) for all values of x . Therefore, the radius of convergence is ∞ .

8. **C** The series is an Alternating Series, so $|s - s_3| < |a_4| = 48e^{-64}$

9. **A** $f'(x) = \frac{1}{1+x}$, which is a geometric sum with $a = 1, r = -x$, so $f'(x) = \sum_{n=0}^{\infty} (-x)^n = \sum_{n=0}^{\infty} (-1)^n x^n$. Integration with respect to x yields $f(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{n+1}}{n+1}$ (since $f(0) = 0$, the constant $C = 0$).

10. **B** Completing the square yields $(x+1)^2 + (y-2)^2 + z^2 = 5$, so the center is $(-1, 2, 0)$ and the radius is $\sqrt{5}$.

11. From the chart below, $\cos x = \frac{-(x-\frac{\pi}{2})}{1!} + \frac{(x-\frac{\pi}{2})^3}{3!} - \frac{(x-\frac{\pi}{2})^5}{5!} + \dots$
- $$= \sum_{n=0}^{\infty} \frac{(-1)^{n+1} (x-\frac{\pi}{2})^{2n+1}}{(2n+1)!}$$
- | n | $f^{(n)}(x)$ | $f^{(n)}(\frac{\pi}{2})$ | $\frac{f^{(n)}(\frac{\pi}{2})}{n!} (x-\frac{\pi}{2})^n$ |
|-----|--------------|--------------------------|---|
| 0 | $\cos x$ | 0 | 0 |
| 1 | $-\sin x$ | -1 | $\frac{-1}{1!} (x-\frac{\pi}{2})^1$ |
| 2 | $-\cos x$ | 0 | 0 |
| 3 | $\sin x$ | 1 | $\frac{1}{3!} (x-\frac{\pi}{2})^3$ |
| 4 | $\cos x$ | 0 | 0 |

12. .

(a) From the chart below, $\sqrt{x} \approx T_3(x) =$

$$1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3$$

n	$f^{(n)}(x)$	$f^{(n)}(1)$	$\frac{f^{(n)}(1)}{n!}(x-1)^n$
0	\sqrt{x}	1	$\frac{1}{0!}(x-1)^0 = 1$
1	$\frac{1}{2}x^{-1/2}$	$\frac{1}{2}$	$\frac{\frac{1}{2}}{1!}(x-1)^1 = \frac{1}{2}(x-1)$
2	$-\frac{1}{4}x^{-3/2}$	$-\frac{1}{4}$	$\frac{-\frac{1}{4}}{2!}(x-1)^2 = -\frac{1}{8}(x-1)^2$
3	$\frac{3}{8}x^{-5/2}$	$\frac{3}{8}$	$\frac{\frac{3}{8}}{3!}(x-1)^3 = \frac{1}{16}(x-1)^3$

(b) $a = 1, n = 3$. To find M , note that $|f^{(4)}(x)| = \left| -\frac{15}{16}x^{-7/2} \right| = \frac{15}{16x^{7/2}}$, which is largest on the interval $x \in [1, 2]$ at $x = 1$. Therefore, $M = f^{(4)}(1) = \frac{15}{16}$.

Therefore, $|R_3(x)| \leq \frac{15}{4!}|x-1|^3$. This error is largest on the interval $x \in [1, 2]$ at $x = 2$, so on the interval, $|R_3(x)| \leq \frac{15}{4!}|2-1|^3 = \frac{15}{16 \cdot 24}$.


13.

(a) The series is an Alternating Series. $|a_n| = \frac{n}{n^2+4}$, which is decreasing and approaching 0. Therefore, the series is convergent by the Alternating Series Test.

(b) $\sum_{n=1}^{\infty} |a_n| = \sum_{n=1}^{\infty} \frac{n}{n^2+4}$. Let $b_n = \frac{1}{n}$. Both $|a_n|$ and b_n are greater than 0, and $\frac{a_n}{b_n} = \frac{n}{n^2+4} \cdot \frac{n}{1} = \frac{n^2}{n^2+4}$, which approaches 1. Since $\sum_{n=1}^{\infty} \frac{1}{n}$ is divergent by the P-Test, $\sum_{n=1}^{\infty} |a_n|$ is divergent by the Limit Comparison Test with $\sum_{n=1}^{\infty} \frac{1}{n}$, hence the Alternating Series is not absolutely convergent. (NOTE: Integral Test could be used to show divergence as well).

14.

- (a)
- $\mathbf{a} = \langle 3, 5, 3 \rangle - \langle 2, 4, 5 \rangle = \langle 1, 1, -2 \rangle$
 - $\mathbf{b} = \langle 2, 8, -3 \rangle - \langle 2, 4, 5 \rangle = \langle 0, 4, -8 \rangle$
 - $\mathbf{c} = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{b}|^2} \mathbf{b} = \frac{20}{16+64} \langle 0, 4, -8 \rangle = \langle 0, 1, -2 \rangle$
 - $\mathbf{a} - \mathbf{c} = \langle 1, 0, 0 \rangle$



(b)

15. Apply the Ratio Test: $\left| \frac{a_{n+1}}{a_n} \right| =$

$$\left| \frac{(x+3)^{n+1}}{(n+3)4^{n+1}} \cdot \frac{(n+2)4^n}{(x+3)^n} \right| = \frac{(n+2)}{4(n+3)} |x+3|.$$

We want the limit, $\frac{|x+3|}{4} < 1$, which is true when $|x+3| < 4$, $-4 < x+3 < 4$, or $-7 < x < 1$. Therefore, the radius of convergence is 4. The series diverges (limit > 1) when $|x+3| > 4$, i.e., $x > 1$ or $x < -7$. Since the Ratio Test is inconclusive (limit = 1) at the endpoints, we must test them separately: When $x = -7$, $\sum_{n=1}^{\infty} \frac{(-7+3)^n}{(n+2)4^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n+2}$, which is convergent by the Alternating Series Test. When $x = 1$, $\sum_{n=1}^{\infty} \frac{(1+3)^n}{(n+2)4^n} = \sum_{n=1}^{\infty} \frac{1}{n+2}$, which is divergent by the Integral Test or by Limit Comparison Test with $\sum_{n=1}^{\infty} \frac{1}{n}$. Therefore, the interval of convergence is $[-7, 1)$.