## Section 11.3 The Integral Test and Estimation of Sums

The Integral Test: If f(x) is a positive, continuous, decreasing function on  $[m, \infty]$ , and  $a_k = f(k)$ , then  $\sum_{k=m}^{\infty} a_k$  and  $\int_m^{\infty} f(x) dx$  either both converge or both diverge.

## We can only use the integral test on series whose terms are (eventually) positive and decreasing!



- 1. Determine whether the following series converge or diverge.
  - a.)  $\sum_{n=1}^{\infty} \frac{n}{n+1}$

b.) 
$$\sum_{n=1}^{\infty} \frac{1}{(7n+8)^3}$$

©Amy Austin, February 28, 2019

c.) 
$$\sum_{n=1}^{\infty} \frac{n}{n^2 + 1}$$

d.) 
$$\sum_{n=0}^{\infty} n e^{-n^2}$$

e.) 
$$\sum_{n=2}^{\infty} \frac{1}{n\sqrt{\ln n}}$$

We learned in section 7.8 that  $\int_{1}^{\infty} \frac{1}{x^{p}} dx$  will converge if p > 1 and will diverge if  $p \le 1$ . This gives us the following result:

**P-series Test**: The **p-series**  $\sum_{n=1}^{\infty} \frac{1}{n^p}$ , where p > 0, is convergent if p > 1 and divergent if  $p \le 1$ . The special case where p = 1 is called the **harmonic series**, which diverges.

A few remarks: Since not all series begin with an index of 1, we can often write  $\sum_{n=1}^{\infty} a_n$  simply as  $\sum a_n$ . We are typically only concerned where the index of the series begins if we are interested in the *sum* of the series. Since convergence of a series is dependent on the *end behavior* of its terms, we can use the notation  $\sum a_n$  if we are only interested in the convergence or divergence of the series but not the sum.

2. Determine whether the following series converges or diverges and support your answer.

a.) 
$$\sum_{n=3}^{\infty} \frac{10}{n^{\sqrt{2}}}$$

b.) 
$$\sum_{n=3}^{\infty} \frac{1}{n\sqrt{n}}$$

c.) 
$$\frac{1}{\sqrt[4]{2}} + \frac{1}{\sqrt[4]{3}} + \frac{1}{\sqrt[4]{4}} + \frac{1}{\sqrt[4]{5}} + \dots$$

**Remainder Estimate for The Integral Test**: Suppose  $s_n = \sum_{i=1}^n a_i = a_1 + a_2 + ... + a_n$  is the  $n^{th}$  partial sum of the convergent series  $\sum_{n=1}^{\infty} a_n$ . Then the **remainder** in using  $s_n$  to approximate the sum S is defined to be  $R_n = S - s_n = \sum_{i=n+1}^{\infty} a_i = a_{n+1} + a_{n+2} + ...$ 

Moreover, if  $\sum_{n=1}^{\infty} a_n$  was shown to be convergent by the integral test where  $a_n = f(n)$ , then

$$R_n = \sum_{i=n+1}^{\infty} a_i \le \int_n^{\infty} f(x) \, dx.$$



3. For the series  $\sum_{n=1}^{\infty} \frac{1}{n^2}$ , Find  $s_6$ , the sum of the first 6 terms. Using the Remainder Estimate for the Integral Test, estimate the error,  $R_6$ , in using the sum of the first 6 terms as an approximation to  $\sum_{n=1}^{\infty} \frac{1}{n^2}$ .

4. Consider the series  $\sum_{n=1}^{\infty} \frac{1}{n^6}$ . Using the Remainder Estimate for the Integral Test, find a value of n that will ensure the error in the approximation,  $s_n$ , is less than 0.00001. Express your answer as  $n > \underline{\qquad }'$ .

5. Consider the series  $\sum_{n=1}^{\infty} \frac{1}{n^4}$ . Using the Remainder Estimate for the Integral Test, find a value of *n* that will ensure the error in the approximation,  $s_n$ , is less than  $\frac{1}{95}$ . Express your answer as '*n* >\_\_\_\_\_'.

Once this value of n is found, approximate  $\sum_{n=1}^{\infty} \frac{1}{n^4}$  to within  $\frac{1}{95}$ .