## Exercise on the modular group

The modular group is the group of linear fractional transformations

$$z \mapsto \frac{az+b}{cz+d}, \qquad a, b, c, d \in \mathbb{Z}, \quad ad-bc=1.$$

The group may be viewed as a group of matrices  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  with the understanding that a matrix and its negative are identified (since  $\frac{az+b}{cz+d} = \frac{-az-b}{-cz-d}$ ). The group is known as  $PSL(2,\mathbb{Z})$ , where the letter L stands for "linear", the letter S stands for "special" (meaning determinant equal to 1), and the letter P stands for "projective" (referring to the identification of a matrix and its negative).

In the textbook you read about the *congruence subgroup*, which consists of matrices that are congruent to the identity matrix modulo 2. This exercise addresses instead the full modular group.

**1.** Show that the translation  $z \mapsto z+1$  (corresponding to the matrix  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ ) and the inversion  $z \mapsto -1/z$  (corresponding to the matrix  $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ ) generate the modular group.

The elements of the modular group map the upper half-plane to itself. When z is an arbitrary point in the upper half-plane, the *orbit* of z is the set of all points  $\varphi(z)$  obtained as  $\varphi$  runs over the elements of the modular group. A *fundamental domain* for the modular group is a set consisting of one point from each orbit.<sup>1</sup>



2. Show that the shaded region indicated in the figure is a fundamental domain for the modular group. The region is the union of

$$\{ z : \text{Im } z > 0 \text{ and } |z| > 1 \text{ and } -\frac{1}{2} < \text{Re } z \le \frac{1}{2} \}$$

and the circular arc  $\{ e^{i\theta} : \pi/3 \le \theta \le \pi/2 \}.$ 

Theory of Functions of a Complex Variable II

 $<sup>^{1}</sup>$ The meaning of the word "domain" here differs from our standard usage: a fundamental domain is not necessarily either connected or open.