

THE GEOMETRY OF RANK DECOMPOSITIONS OF MATRIX MULTIPLICATION II: 3×3 MATRICES

GREY BALLARD, CHRISTIAN IKENMEYER, J.M. LANDSBERG, AND NICK RYDER

ABSTRACT. This is the second in a series of papers on rank decompositions of the matrix multiplication tensor. We present new rank 23 decompositions for the 3×3 matrix multiplication tensor $M_{(3)}$. All our decompositions have symmetry groups that include the standard cyclic permutation of factors but otherwise exhibit a range of behavior. One of them has 11 cubes as summands and admits an unexpected symmetry group of order 12.

We establish basic information regarding symmetry groups of decompositions and outline two approaches for finding new rank decompositions of $M_{(\mathbf{n})}$ for larger \mathbf{n} .

1. INTRODUCTION

This is the second in a planned series of papers on the geometry of rank decompositions of the matrix multiplication tensor $M_{(\mathbf{n})} \in \mathbb{C}^{\mathbf{n}^2} \otimes \mathbb{C}^{\mathbf{n}^2} \otimes \mathbb{C}^{\mathbf{n}^2}$. Our goal is to obtain new rank decompositions of $M_{(\mathbf{n})}$ by exploiting symmetry. For a tensor $T \in \mathbb{C}^m \otimes \mathbb{C}^m \otimes \mathbb{C}^m$, the *rank* of T is the smallest r such that $T = \sum_{j=1}^r a_j \otimes b_j \otimes c_j$, with $a_j, b_j, c_j \in \mathbb{C}^m$. The rank of $M_{(\mathbf{n})}$ is a standard complexity measure of matrix multiplication, in particular, it governs the total number of arithmetic operations needed to multiply two matrices.

In this paper we present rank 23 decompositions of $M_{(3)}$ that have large symmetry groups, in particular all admit the standard cyclic \mathbb{Z}_3 -symmetry of permuting the three tensor factors. Although many rank 23 decompositions of $M_{(3)}$ are known [15, 13], none of them was known to admit the standard cyclic \mathbb{Z}_3 -symmetry.

We describe techniques to determine symmetry groups of decompositions and to determine if two decompositions are in the same *family*, as defined in §4. We also develop a framework for using representation theory to write down new rank decompositions for $M_{(\mathbf{n})}$ for all \mathbf{n} . Similar frameworks are also being developed implicitly and explicitly in [5] and [12].

As discussed below, decompositions come in families. DeGroot [8] has shown, in contrast to $M_{(3)}$, the family generated by Strassen's decomposition is the unique family of rank seven decompositions of $M_{(2)}$. Unlike $M_{(2)}$, it is still not known if the tensor rank of $M_{(3)}$ is indeed 23. The best lower bound on the rank is 16 [17]. There have been substantial unsuccessful efforts to find smaller decompositions by numerical methods. While some researchers have taken this as evidence that 23 might be optimal, it just might be the case that rank 22 (or smaller) decompositions might be much rarer than border rank 22 decompositions, and so using numerical search methods, and given initial search points, when they converge, with probability nearly one would converge to border rank decompositions.

Key words and phrases. matrix multiplication complexity, alternating least squares, MSC 68Q17, 14L30, 15A69.

Landsberg supported by NSF grant DMS-1405348. Ballard supported in part by an appointment to the Sandia National Laboratories Truman Fellowship in National Security Science and Engineering, sponsored by Sandia Corporation (a wholly owned subsidiary of Lockheed Martin Corporation) as Operator of Sandia National Laboratories under its U.S. Department of Energy Contract No. DE-AC04-94AL85000.

In this paper we focus on decompositions with *standard cyclic symmetry*: viewed as a trilinear map, matrix multiplication is $(X, Y, Z) \mapsto \text{trace}(XYZ)$, where X, Y, Z are $\mathbf{n} \times \mathbf{n}$ matrices. Since $\text{trace}(XYZ) = \text{trace}(YZX)$, the matrix multiplication tensor $M_{(\mathbf{n})} \in \mathbb{C}^{\mathbf{n}^2} \otimes \mathbb{C}^{\mathbf{n}^2} \otimes \mathbb{C}^{\mathbf{n}^2} = \text{Mat}_{\mathbf{n} \times \mathbf{n}}^{\otimes 3}$ has a \mathbb{Z}_3 symmetry by cyclically permuting the three factors. If one applies this cyclic permutation to a tensor decomposition of $M_{(\mathbf{n})}$, it is transformed to another decomposition of $M_{(\mathbf{n})}$. When a decomposition is transformed to itself, we say the decomposition has *cyclic symmetry* or $\mathbb{Z}_3^{\text{std}}$ -*invariance*.

In §2 we present three of our examples. We explain the search methods used in §3. We then discuss techniques for determining symmetry groups in §4 and determine the symmetry groups of our decompositions in §5. Our searches sometimes found equivalent decompositions but in different coordinates, and the same techniques enabled us to identify when two decompositions are equivalent. Further techniques for studying decompositions are presented in §6 and §8, respectively in terms of configurations of points in \mathbb{P}^{n-1} and eigenvalues. In §7, we precisely describe the subspace of \mathbb{Z}_3 -invariant tensors in $\text{Mat}_{\mathbf{n} \times \mathbf{n}}^{\otimes 3}$, as well as the subspaces invariant under other finite groups. In an appendix §A, we present additional decompositions that we found.

Why search for decompositions with symmetry? There are many examples where the optimal decompositions (or expressions) of tensors with symmetry have some symmetry. This is true of $M_{(2)}$ [6, 4], the monomial $x_1 \cdots x_n$ [19], also see [16, §7.1], the optimal determinantal expression of perm_3 [18], and other cases. In any case, imposing the symmetry i) reduces the size of the search space, and ii) provides a guide for constructing decompositions through the use of “building blocks”.

Notation and conventions. A, B, C, U, V, W are vector spaces, A^* is the dual vector space to A , $GL(A)$ denotes the group of invertible linear maps $A \rightarrow A$, $SL(A)$ the maps with determinant one, and $PGL(A) = GL(A)/\{\text{CId}_A \setminus 0\}$ the group of projective transformations of projective space $\mathbb{P}A$. The action of $GL(A)$ on $A \otimes A^*$ descends to an action of $PGL(A)$. If $a \in A$, $[a]$ denotes the corresponding point in projective space. \mathfrak{S}_d denotes the permutation group on d elements and \mathbb{Z}_d denotes the cyclic group of order d . For $X \subset \mathbb{P}V$, $\hat{X} \subset V$ denotes its pre-image under the projection map union the origin. $X^{(\times r)}$ denotes the quotient $X^{\times r}/\mathfrak{S}_r$. We write $SL_m = SL(\mathbb{C}^m)$, $GL_m = GL(\mathbb{C}^m)$, and $H^{SL_m} \subset SL_m$ is the subgroup of diagonal matrices. For a matrix a , a_j^i denotes the entry in the i -th row and j -th column.

Acknowledgements. Work on this paper began during the fall 2014 semester program *Algorithms and Complexity in Algebraic Geometry* at the Simons Institute for the Theory of Computing. We thank the Institute for bringing us together and making this paper possible.

2. EXAMPLES

2.1. A rank 23 decomposition of $M_{(3)}$ with $\mathbb{Z}_4 \times \mathbb{Z}_3$ symmetry. Let $\mathbb{Z}_4^{a_0} \subset GL_3 \subset GL_3^{\times 3}$ be generated by

$$(1) \quad a_0 = \begin{pmatrix} 0 & 0 & -1 \\ 1 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix},$$

where the action on $x \otimes y \otimes z$ where x, y, z are 3×3 -matrices is $x \otimes y \otimes z \mapsto a_0 x a_0^{-1} \otimes a_0 y a_0^{-1} \otimes a_0 z a_0^{-1}$ and let $\mathbb{Z}_3^{\text{std}}$ denote the standard cyclic symmetry.

Here is the decomposition, we call it $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$:

$$\begin{aligned}
 (2) \quad M_{\langle 3 \rangle} &= - \begin{pmatrix} 0 & 0 & -1 \\ 1 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix}^{\otimes 3} \\
 (3) \quad &+ \mathbb{Z}_4^{a_0} / \mathbb{Z}_2^{a_0^2} \cdot \begin{pmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} \\
 (4) \quad &+ \mathbb{Z}_4^{a_0} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} \\
 (5) \quad &+ \mathbb{Z}_4^{a_0} \cdot \begin{pmatrix} 0 & -1 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} \\
 (6) \quad &+ \mathbb{Z}_3^{std} \times \mathbb{Z}_4^{a_0} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 1 & -1 \end{pmatrix}.
 \end{aligned}$$

The decomposition is a sum of 23 terms, each of which is a trilinear form on matrices. For example, the term $\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 1 & -1 \end{pmatrix}$ sends a triple of matrices (x, y, z) to the number $x_{23}(y_{32} - y_{31})(z_{22} - z_{23} + z_{32} - z_{33})$. This expresses $M_{\langle 3 \rangle}$ in terms of five $\mathbb{Z}_4 \times \mathbb{Z}_3$ -orbits, of sizes 1, 2, 4, 4, 12.


2.2. An element of the $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$ family with a_0 diagonalized. It is also illuminating to diagonalize a_0 , in other words decompose the decomposition with respect to the \mathbb{Z}_4 action: In the following plot each row of 3×3 matrices forms an orbit under the \mathbb{Z}_4 -action. In the first four rows are the eleven 3×3 matrices that appear as cubes in the decomposition. In remaining 4 rows each of the four columns forms three rank one tensors by tensoring the three matrices in the column in three different orders: 1-2-3, 2-3-1, and 3-1-2.


Each complex number in each matrix is depicted by plotting its position in the complex plane. To help identify the precise position, a square is drawn with vertices $(-\frac{1}{2}, -\frac{1}{2})$, $(\frac{1}{2}, -\frac{1}{2})$, $(-\frac{1}{2}, \frac{1}{2})$, $(\frac{1}{2}, \frac{1}{2})$. To quickly identify the absolute value of a complex number they are color coded:

Symbol for the complex number					
absolute value	1	$\frac{1}{\sqrt{2}}$	$\frac{1}{2\sqrt{2}}$	$\frac{1}{2}$	$\frac{\sqrt{5}}{2\sqrt{2}}$

Numbers with a blue background are the sum of two numbers with a yellow background. Numbers with a purple background are twice the numbers with a yellow background. Numbers with a red background are twice the numbers with a blue background. Numbers with a green background are the sum of a number with a yellow background and a number with a blue background.

Matrix cells with zeros are left empty.





2.3. A cyclic-invariant decomposition in the Laderman family. Let $\mathcal{S}_{Laderman}$ denote the rank 23 Laderman decomposition of $M_{(3)}$. By [4] the symmetry group of this decomposition (see §4) is $\Gamma_{\mathcal{S}_{Laderman}} = (\mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes (\mathbb{Z}_3 \rtimes \mathbb{Z}_2) \simeq \mathfrak{S}_4$, where $\mathbb{Z}_2 \times \mathbb{Z}_2 \subset SL_3^{\times 3}$. We found a decomposition, which we call $\mathcal{S}_{Lader-\mathbb{Z}_3^{std}}$ that we identified as a \mathbb{Z}_3^{std} -invariant member of the Laderman *family* in the sense of §4.

Let

$$\tau_{23} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \epsilon_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \tau_{13} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix},$$

Let $\phi(x \otimes y \otimes z) = \tau_{13} x \tau_{13} \otimes \tau_{13} y \otimes z \tau_{13}$, and let $\zeta(x \otimes y \otimes z) = \epsilon_2 y^T \epsilon_2 \otimes \epsilon_2 x^T \epsilon_2 \otimes \epsilon_2 z^T \epsilon_2$. Let π be the generator of \mathbb{Z}_3^{std} . Let Γ denote the group generated by π, ϕ , and ζ .

$$(7) \quad M_{\{3\}} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3}$$

$$(8) \quad + \Gamma/(\mathbb{Z}_3^\pi \times \mathbb{Z}_2^\zeta) \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}^{\otimes 3}$$

$$(9) \quad + \Gamma/(\mathbb{Z}_3^\pi \times \mathbb{Z}_2^\zeta) \cdot \begin{pmatrix} -1 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3}$$

$$(10) \quad + \Gamma/(\mathbb{Z}_2^\phi \times \mathbb{Z}_2^\zeta) \cdot \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & -1 & 0 \\ 1 & -1 & -1 \\ 0 & 1 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$(11) \quad + \Gamma/\mathbb{Z}_3^\pi \cdot \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3}$$

These five orbits are respectively of sizes 1, 4, 4, 6, 8. There are five \mathbb{Z}_3^{std} invariant terms, one from each of (7),(8),(9) and two from (11) because ζ preserves \mathbb{Z}_3^{std} -invariance.

We originally found this decomposition by numerical methods. The incidence graphs discussed in §4.3 gave us evidence that it should be in the Laderman family, and then it was straightforward to find the transformation that exchanged $\mathcal{S}_{Laderman}$ and $\mathcal{S}_{Lader-\mathbb{Z}_3^{std}}$, namely $x \otimes y \otimes z \mapsto \tau_{12}x \otimes y \epsilon_2 \tau_{12} \otimes \tau_{12} \epsilon_2 z \tau_{12}$. See §4.3 for more discussion. As shown in in [4], Γ is isomorphic to \mathfrak{S}_4 and these are all the symmetries of the Laderman family. Translated to $\mathcal{S}_{Laderman}$ as presented in [4], these five orbits are respectively $\{19\}$, $\{20, 21, 22, 23\}$, $\{4, 7, 12, 16\}$, $\{1, 3, 6, 10, 11, 14\}$ and $\{2, 5, 8, 9, 13, 15, 17, 18\}$.

Remark 2.1. As pointed out in [20], there are similarities between this decomposition and Strassen's.

Remark 2.2. The decompositions of Johnson-McLoughlin [13] cannot have \mathbb{Z}_3 -invariant decompositions for rank reasons: The first space of decompositions has five terms (those numbered 3,12,16,22,23 in [13]) where there are two matrices of rank one and one of rank greater than one, while to have any external \mathbb{Z}_3 -symmetry, the number of such would have to be a multiple of three. The second space has a unique matrix of rank three appearing (23b), so is ruled out for the same reason.

2.4. Decomposition $\mathcal{S}_{2fix-\mathbb{Z}_3}$. Here is a decomposition with two \mathbb{Z}_3 -fixed points:

$$\begin{aligned}
 (12) \quad M_{\langle 3 \rangle} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix}^{\otimes 3} \\
 (13) \quad &+ \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} \\
 (14) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix} \\
 (15) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \\
 (16) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \\
 (17) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix} \\
 (18) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & -1 & 0 \end{pmatrix} \\
 (19) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
 (20) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} -1 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}
 \end{aligned}$$

An interesting feature of this decomposition is that it “nearly” has a transpose-like symmetry, as discussed in later sections.

3. DISCUSSION ON THE NUMERICAL METHODS USED

Our techniques for discovering cyclic-invariant decompositions use numerical optimization methods that are designed to compute approximations rather than exact decompositions. We also use heuristics in order to encourage sparsity in the solutions, and a fortunate by-product of the sparsity is that the nonzero values often tend towards a discrete set of values from which an exact decomposition can be recognized. Our methods are based on techniques that have proved successful in discovering generic exact decompositions (those that have no noticeable symmetries) [1, 22]; we summarize this approach in Section 3.1. Our search process can be divided into two phases. First, as we discuss in Section 3.2, we find a dense, cyclic-invariant, approximate solution using nonlinear optimization methods. Then, as we describe in Section 3.3, we transform the dense approximate solution to an exact cyclic-invariant decomposition using heuristics that encourage sparsity.

3.1. Alternating Least Squares. Computing low-rank approximations of tensors is a common practice in data analysis when the data represents multi-way relationships. The most popular algorithm for computing approximations with CANDECOMP/PARAFAC (CP) structure, i.e., rank decompositions, is known as alternating least squares (ALS) [14]. In particular, in the case of the matrix multiplication tensor, the objective function of the optimization problem is given by

$$(21) \quad \arg \min_{X,Y,Z} \left\| M_{(\mathbf{n})} - \sum_{r=1}^R x_r \otimes y_r \otimes z_r \right\|,$$

where X , Y , and Z are $\mathbf{n}^2 \times R$ *factor matrices* with r th columns given by x_r , y_r , and z_r and this and all norms correspond to the square root of the sum of squares of the entries of the tensor (or matrix). The objective function itself is nonlinear and non-convex and cannot be solved in closed form. However, if two of the three factor matrices are held fixed, then the resulting objective function is a linear least squares problem and can be solved using linear algebra. Thus, ALS works by alternating over the factor matrices, holding two fixed and updating the third, and iterating until convergence.

In general, ALS iterations are performed in floating point arithmetic. While an objective function value of 0 corresponds to a rank decomposition, with ALS we can hope for an objective function value only as small as the finite precision allows. In the case of the matrix multiplication tensor, there are multiple pitfalls that make it difficult to find approximations that approach objective function values of 0. The most successful technique was proposed by Smirnov and uses regularization, with objective function given by

$$(22) \quad \arg \min_{X,Y,Z} \left\| M_{(\mathbf{n})} - \sum_{r=1}^R x_r \otimes y_r \otimes z_r \right\| + \lambda (\|X - \tilde{X}\| + \|Y - \tilde{Y}\| + \|Z - \tilde{Z}\|),$$

for judicious choices of scalar λ and matrices \tilde{X} , \tilde{Y} , and \tilde{Z} [22]. We discuss effective choices for the regularization parameters in Section 3.3. This method works for the cases of non-square matrix multiplication and has been used to discover exact rank decompositions for many small cases [1, 22], but it does not encourage solutions to reflect any symmetries.

3.2. Nonlinear Optimization for Dense Approximations. To enforce cyclic invariance on rank decompositions, we can impose structure on the factor matrices X , Y , and Z . In particular, if a cyclic-invariant approximation includes the component $x_r \otimes y_r \otimes z_r$, then it must also include $z_r \otimes x_r \otimes y_r$ and $y_r \otimes z_r \otimes x_r$. This implies either that all three components appear or that $x_r = y_r = z_r$, which means the factor matrices have the following structure:

$$(23) \quad \begin{aligned} X &= (A \ B \ C \ D) \\ Y &= (A \ D \ B \ C) \\ Z &= (A \ C \ D \ B), \end{aligned}$$

where A is an $\mathbf{n}^2 \times P$ matrix and B, C, D are $\mathbf{n}^2 \times Q$ matrices with $P + 3Q = R$. With this structure, (21) becomes

$$(24) \quad \arg \min_{A,B,C,D} \left\| M_{(\mathbf{n})} - \sum_{p=1}^P a_p \otimes a_p \otimes a_p - \sum_{q=1}^Q (b_q \otimes c_q \otimes d_q + d_q \otimes b_q \otimes c_q + c_q \otimes d_q \otimes b_q) \right\|.$$

Note that the total number of variables (now spread across 4 matrices) is reduced by a factor of 3.

Again, this objective function is nonlinear and non-convex. It is possible to use an ALS approach to drive the objective function value to 0; however, while (24) is linear in B , C , and D , it is not linear in A . Thus, the optimal update for A , for fixed B , C , and D , cannot be computed in closed form. While there are numerical optimization techniques for A , we were not successful in using an ALS approach to drive the objective function value close to zero.

Instead, we used generic nonlinear optimization software to search for cyclic-invariant approximations. In particular, we used the LOQO software package [23], which relies on the AMPL [9] modeling language to specify the optimization problem. In order to find solutions, we try all possible values of P and Q , use multiple random starting points, and constrain the variable entries to be no greater than 1 in absolute value. For fixed R , there are $\lfloor R/3 \rfloor$ possible values for P and Q . Multiple starting points are required because the objective function is non-convex: numerical optimization techniques are sensitive to starting points in this case, with approximations often getting stuck at local minima. Constraining the variables to be no greater than 1 in absolute value is a technique to avoid converging numerically to border rank decompositions, in which case some variable values must grow in magnitude to continue to improve the objective function value. Driving the objective function value to zero with bounded variable values ensures that the approximation corresponds to a rank decomposition.

When we are successful, we obtain matrices A, B, C, D that correspond to an objective function value very close to zero. Thus, the approximation has the cyclic invariance we desire. However, these matrices are dense and have floating point values throughout the $[-1, 1]$ range. Rounding these floating point values, even to a large set of discrete rational values, typically does not yield an exact rank decomposition. The next section describes the techniques we use to convert dense approximate solutions to exact solutions.

3.3. Heuristics to Encourage Sparsity for Exact Decompositions. In order to obtain exact decompositions, we use the ALS regularization heuristics that proved effective for the non-invariant case, given in equation (22). However, those techniques ignore the cyclic-invariant structure in the dense approximations we obtain from the techniques described in §3.2. The heuristic we used to discover cyclic-invariant rank decomposition consists of alternating between ALS iterations with regularization and projection of the approximation back to the set of cyclic-invariant solutions.

We now describe the specifics of the regularization terms from (22). The scalar parameter λ determines the relative importance between an accurate approximation of the matrix multiplication tensor and adherence to the regularization terms. In the context of ALS, only one of the regularization terms affects the optimization problem when updating one factor matrix. The *target matrices* $\tilde{X}, \tilde{Y}, \tilde{Z}$ are designed to encourage the corresponding factor matrices to match a desired structure, and they can be defined differently for each iteration of ALS. Here is the technique proposed by Smirnov [22]. Consider the update of factor matrix X . By default, \tilde{X} is set to have the values of X from the previous iteration. If any of the values are larger than 1 (or any specified maximum value), then the corresponding value of \tilde{X} is set to magnitude 1 with the corresponding sign. Then, for a given number z of desired zeros, the smallest z values of \tilde{X} are set to exactly 0. Thus, the regularization term $\|X - \tilde{X}\|_F$ will encourage any large values of X to tend towards ± 1 and the smallest z values to tend toward 0. The parameter z can be varied over iterations and also across factor matrices (though in the case of cyclic-invariant approximations the set of values within each factor matrix is the same as those of the other factor matrices).

Because ALS does not enforce cyclic invariance, the approximation will tend to deviate from the structure given in equation (23). We project back to a cyclic-invariant approximation by

setting all three values that should be the same in each of the factor matrices to their average value.

Fortunately, perhaps miraculously, by encouraging sparsity and bounding the variable values, when many entries are driven to zero, the nonzero values tend towards a discrete set of values (usually ± 1). However, the process of converting a dense approximation to an exact decomposition involves manual tinkering and much experimentation. The basic approach we have used successfully is to maintain a tight approximation to the matrix multiplication tensor, start with $z = 0$, gradually increase z , frequently project back to cyclic invariance, and play with the λ parameter between values of 10^{-3} and 1. ALS iterations are relatively cheap, so often 100 or 1000 iterations can be taken with a given parameter setting before changing the configuration. While this process is artful and lacks any guarantees of success, we have nearly always been able to convert cyclic-invariant dense approximations of $M_{(3)}$ to exact decompositions.

4. SYMMETRY GROUPS OF TENSORS AND DECOMPOSITIONS

In this section we explain how we found the additional symmetries beyond the \mathbb{Z}_3^{std} that was built into the search, and describe the full symmetry groups of the decompositions. We establish additional properties regarding symmetries for use in future work. We begin with a general discussion of symmetry groups of tensors and their decompositions.

4.1. Symmetry groups of tensors. Let V be a complex vector space and let $T \in V^{\otimes k}$. Define the *symmetry group* of T , $G_T \subset GL(V)^{\times k} \rtimes \mathfrak{S}_k$ to be the subgroup preserving T , where \mathfrak{S}_k acts by permuting the factors.

For a rank decomposition $T = \sum_{j=1}^r t_j$, where each t_j has rank one, i.e., $t_j = v_{1j} \otimes \cdots \otimes v_{kj}$, define the set $\mathcal{S} := \{t_1, \dots, t_r\}$, which we also call the decomposition, and the *symmetry group of the decomposition* $\Gamma_{\mathcal{S}} := \{g \in G_T \mid g \cdot \mathcal{S} = \mathcal{S}\}$. We also consider \mathcal{S} as a point of the variety of r -tuples of unordered points on the Segre variety of rank one tensors, denoted $\hat{Seg}(\mathbb{P}V \times \cdots \times \mathbb{P}V)^{(xr)}$.

If $g \in G_T$, then $g \cdot \mathcal{S} := \{gt_1, \dots, gt_r\}$ is also a rank decomposition of T , and $\Gamma_{g \cdot \mathcal{S}} = g\Gamma_{\mathcal{S}}g^{-1}$ (see [6]), so decompositions come in *families* parametrized by $G_T/\Gamma_{\mathcal{S}}$, and each member of the family has the same abstract symmetry group. We reserve the term *family* for G_T -orbits. The quasi-projective subvariety $\Sigma_r^T \subset \hat{Seg}(\mathbb{P}A_1 \times \cdots \times \mathbb{P}A_k)^{(xr)}$ of all rank r decompositions is a union of G_T orbits.

If one is not concerned with the rank of a decomposition, then for any finite subgroup $\Gamma \subset G_T$, T admits rank decompositions \mathcal{S} with $\Gamma \subseteq \Gamma_{\mathcal{S}}$, by taking any rank decomposition of T and then averaging it with its Γ -translates. We will be concerned with rank decompositions that are minimal or close to minimal, so only very special groups Γ can occur.

4.2. Matrix multiplication. The symmetry groups of matrix multiplication decompositions are useful for determining if two decompositions lie in the same family, and the groups that appear in known decompositions will be a guide for constructing decompositions in future work.

The symmetry groups of many of the decompositions that have already appeared in the literature are determined in [5]. Burichenko does not use the associated graphs discussed below. One can recover the results of [5] with shorter proofs by using them.

Let $T = M_{(\mathbf{n})} \in \mathbb{C}^{\mathbf{n}^2} \otimes \mathbb{C}^{\mathbf{n}^2} \otimes \mathbb{C}^{\mathbf{n}^2} =: A \otimes B \otimes C = (U^* \otimes V) \otimes (V^* \otimes W) \otimes (W^* \otimes U)$ where $U, V, W = \mathbb{C}^{\mathbf{n}}$. Recall that $M_{(\mathbf{n})}$ is the re-ordering of $\text{Id}_U \otimes \text{Id}_V \otimes \text{Id}_W$ and

$$(25) \quad G_{M_{(\mathbf{n})}} = [(PGL(U) \times PGL(V) \times PGL(W)) \rtimes \mathbb{Z}_3] \rtimes \mathbb{Z}_2 \subset GL(A) \times GL(B) \times GL(C) \rtimes \mathfrak{S}_3,$$

see, e.g., [7, Thms. 3.3,3.4], [6, Prop. 4.1], [11, Thm. 2.9] or [5, Prop. 4.7]. The $\mathbb{Z}_2 \subset GL(A) \times GL(B) \times GL(C) \rtimes \mathfrak{S}_3$ may be generated by e.g., $(a \otimes b \otimes c) \mapsto (a^T \otimes c^T \otimes b^T)$, and the \mathbb{Z}_3

by cyclically permuting the factors A, B, C . The $\mathbb{Z}_3 \rtimes \mathbb{Z}_2$ is isomorphic to \mathfrak{S}_3 , but it is more naturally thought of as $\mathbb{Z}_3 \rtimes \mathbb{Z}_2$, since it is not the \mathfrak{S}_3 appearing in (25).

Thus if \mathcal{S} is a rank decomposition of $M_{\langle \mathbf{n} \rangle}$, then $\Gamma_{\mathcal{S}} \subset [(PGL(U) \times PGL(V) \times PGL(W)) \rtimes \mathbb{Z}_3] \rtimes \mathbb{Z}_2$.

Remark 4.1. As pointed out by Burichenko [4], for matrix multiplication rank decompositions one can define an *extended symmetry group* $\hat{\Gamma}_{\mathcal{S}}$ by viewing $A \otimes B \otimes C = (U^* \otimes U)^{\otimes 3} = (U^*)^{\otimes 3} \otimes (U^{\otimes 3})$. We do not study such groups in this paper.

We call a $\mathbb{Z}_2 \subset \Gamma_{\mathcal{S}}$ a *transpose like symmetry* if it corresponds to the symmetry of $M_{\langle \mathbf{n} \rangle}$ given by $x \otimes y \otimes z \mapsto x^T \otimes z^T \otimes y^T$, or a cyclic variant of it such as $x \otimes y \otimes z \mapsto y^T \otimes x^T \otimes z^T$, composed with an element of $PGL(U) \times PGL(V) \times PGL(W)$ such that the total map is an involution on the elements of \mathcal{S} . Transpose like symmetries where the elements of $PGL(U) \times PGL(V) \times PGL(W)$ are all the identity (which we will call *convenient transpose symmetries*) do not appear to be compatible with standard cyclic symmetries in minimal decompositions, at least this is the case for rank seven decompositions of $M_{\langle 2 \rangle}$ and the known rank 23 decompositions of $M_{\langle 3 \rangle}$.

Example 4.2. Let x_j^i , $1 \leq i, j \leq n$ be a basis of A , y_j^i a basis of B , and z_j^i a basis of C . Consider the standard decomposition \mathcal{S}_{std} of $M_{\langle \mathbf{n} \rangle}$ of size \mathbf{n}^3 :

$$(26) \quad M_{\langle \mathbf{n} \rangle} = \sum_{i,j,k=1}^{\mathbf{n}} x_j^i \otimes y_k^j \otimes z_i^k.$$

Let $H^{SL_n} \subset SL_n$ denote the maximal torus (diagonal matrices). It is clear $\Gamma_{\mathcal{S}_{std}} \supseteq (H^{SL_n} \times \mathfrak{S}_n)^{\times 3} \rtimes (\mathbb{Z}_3 \rtimes \mathbb{Z}_2)$ because for $((\lambda_1, \dots, \lambda_n), (\mu_1, \dots, \mu_n), (\nu_1, \dots, \nu_n)) \in (H^{SL_n})^{\times 3}$, we have

$$(\lambda_i \mu_j^{-1} x_j^i) \otimes (\mu_j \nu_k^{-1} y_k^j) \otimes (\nu_k \lambda_i^{-1} z_i^k) = x_j^i \otimes y_k^j \otimes z_i^k$$

and for all $\sigma, \tau, \eta \in \mathfrak{S}_n$, we have the equality of sets $\{\cup_{i,j,k} x_j^i \otimes y_k^j \otimes z_i^k\} = \{\cup_{i,j,k} x_{\tau(j)}^{\sigma(i)} \otimes y_{\eta(k)}^{\tau(j)} \otimes z_{\sigma(i)}^{\eta(k)}\}$, the cyclic symmetry is evident, and the \mathbb{Z}_2 may be generated e.g., by $x \otimes y \otimes z \mapsto x^T \otimes z^T \otimes y^T$.

The *Comon conjecture* [2, 3], in its original form, asserts that a tensor $T \in (\mathbb{C}^N)^{\otimes d}$ that happens to be symmetric, will have an optimal rank decomposition consisting of rank one symmetric tensors, that is, the symmetric tensor rank of T equals the usual tensor rank. An explicit counter-example to this when $N = 800$ has been asserted in [21]. Nevertheless, there appear to be many instances where it is known to hold, see, e.g., [10, 3] so we pose the following question:

Question 4.3 (Generalized Comon Question). Given $T \in (\mathbb{C}^N)^{\otimes d}$ that is invariant under some $\Gamma \subset \mathfrak{S}_d$, when does there exist an optimal rank decomposition \mathcal{S} of T that is Γ -invariant, i.e., $\Gamma \subseteq \Gamma_{\mathcal{S}}$?

4.3. Invariants associated to a decomposition of $M_{\langle \mathbf{n} \rangle}$. Let $M_{\langle \mathbf{n} \rangle} = \sum_{j=1}^r t_j$ be a rank decomposition \mathcal{S} for $M_{\langle \mathbf{n} \rangle}$ and write $t_j = a_j \otimes b_j \otimes c_j$. Partition \mathcal{S} by rank triples into disjoint subsets: $\mathcal{S} := \{\mathcal{S}_{1,1,1}, \mathcal{S}_{1,1,2}, \dots, \mathcal{S}_{n,n,n}\}$, where for $s \leq t \leq u$ we set $\mathcal{S}_{s,t,u} = \{t_j \mid \{\text{rank}(a_j), \text{rank}(b_j), \text{rank}(c_j)\} = \{s, t, u\}\}$. Then $\Gamma_{\mathcal{S}}$ preserves each $\mathcal{S}_{s,t,u}$.

We can say more about $\Gamma_{\mathcal{S}_{1,1,1}}$: If $a \in U^* \otimes V$ and $\text{rank}(a) = 1$, then there are unique points $[\mu] \in \mathbb{P}U^*$ and $[v] \in \mathbb{P}V$ such that $[a] = [\mu \otimes v]$, so define $\mathcal{S}_{U^*} \subset \mathbb{P}U^*$ and $\mathcal{S}_U \subset \mathbb{P}U$ to correspond to the elements in $\mathbb{P}U^*$ (resp. $\mathbb{P}U$) appearing in $\mathcal{S}_{1,1,1}$. Let $\tilde{\mathcal{S}}_{U^*} \subset \mathbb{P}U^*$ and $\tilde{\mathcal{S}}_U \subset \mathbb{P}U$ correspond to the elements appearing in some rank one matrix in \mathcal{S} .

Since we are concerned with decompositions with a standard cyclic symmetry, we will have $\mathcal{S}_U \simeq \mathcal{S}_V \simeq \mathcal{S}_W$ and $\mathcal{S}_{U^*} \simeq \mathcal{S}_{V^*} \simeq \mathcal{S}_{W^*}$ and similarly for the tilded spaces.





FIGURE 1. Incidence graph of standard decomposition

FIGURE 2. Incidence graph of decomposition $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$

Define a bipartite graph $\mathcal{IG}_{\mathcal{S}}$, the *incidence graph* where the top vertex set is given by elements in $\tilde{\mathcal{S}}_{U^*}$ (or \mathcal{S}_{U^*}) and the bottom vertex set by elements in $\tilde{\mathcal{S}}_U$. Draw an edge between elements $[\mu]$ and $[v]$ if they are *incident*, i.e., $\mu(v) = 0$. Geometrically, $[v]$ belongs to the hyperplane determined by $[\mu]$ (and vice-versa). One can weight the vertices of this graph in several ways, the simplest is just by the number of times the element appears in the decomposition. In practice (see the examples below) this has been enough to determine the symmetry group $\Gamma_{\mathcal{S}}$, in the sense that it cuts the possible size of the group down and it becomes straightforward to determine $\Gamma_{\mathcal{S}}$ as a subgroup of the symmetry group of $\mathcal{IG}_{\mathcal{S}}$.

Incidence graphs for the four decompositions are given in Figures 1, 2, 3, and 4. Here interpret the top set of vectors as column vectors and the bottom set as row vectors. Then μ from the top set is incident to v from the bottom set if the scalar $v\mu$ is zero, and then μ and v are joined by an edge in the graph.

For decompositions where the three copies of $Mat_{n \times n}$ have been identified, such as our \mathbb{Z}_3^{std} -invariant decompositions, consider the *restricted family* $PGL_n \rtimes (\mathbb{Z}_3 \rtimes \mathbb{Z}_2) \cdot \mathcal{S}$, where $PGL_n \subset PGL_n^{\times 3}$ is diagonally embedded and consider the corresponding restricted symmetry groups $\Gamma_{\mathcal{S}}^{res} \subset PGL_n \rtimes (\mathbb{Z}_3 \rtimes \mathbb{Z}_2)$. We define a second graph $\mathcal{PG}_{\mathcal{S}}$ that is an invariant of this restricted




FIGURE 3. Incidence graph of decomposition $\mathcal{S}_{Lader-\mathbb{Z}_3^{std}}$




FIGURE 4. Incidence graph of decomposition $\mathcal{S}_{2fix-\mathbb{Z}_3}$

family, the *pairing graph* which has an edge between $[\mu]$ and $[v]$ if $\mu \otimes v$ appears in the decomposition, and triples of edges that appear in the same summands are grouped by color, and one can weight the edge by the number of times it appears. Pairing graphs for the four decompositions are given in Figures 5, 6, 7, and 8. The dashed black lines correspond to cubes, so should be interpreted as edges with multiplicity three.

As is clear from this discussion, one can continue labeling and coloring to get additional information about the decomposition.

5. SYMMETRY GROUPS OF OUR DECOMPOSITIONS

With the graphs in hand, it is straightforward to determine the symmetry groups.

5.1. Symmetries of $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$.

Proposition 5.1. *The symmetry group of $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$ is $\Gamma_{\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}} = \mathbb{Z}_4^{a_0} \times \mathbb{Z}_3^{std}$, where $\mathbb{Z}_4^{a_0} \subset PGL_3 \subset PGL_3^{\times 3}$ is generated by a_0 of (1).*




FIGURE 5. Pairing graph of standard decomposition




FIGURE 6. Pairing graph of decomposition $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$





FIGURE 7. Pairing graph of decomposition $\mathcal{S}_{Lader-\mathbb{Z}_3^{std}}$


 FIGURE 8. Pairing graph of decomposition $\mathcal{S}_{2fix-\mathbb{Z}_3}$

Proof. The incidence graph shows no transpose-like symmetry is possible as points occur with different frequencies in the two spaces. Since we already know the \mathbb{Z}_3^{std} symmetry, we are reduced to determining $\Gamma_{\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}} \cap PGL_3^{\times 3}$.

Say $\gamma = (g, h, k) \in PGL_3^{\times 3}$ preserves $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$. Since a_0 is the unique term in the decomposition of full rank, $a_0^{\otimes 3}$ is fixed by γ . Since a_0 is fixed up to scale, $[a_0] = [ga_0h^{-1}] = [ha_0k^{-1}] = [ka_0g^{-1}]$. This implies, up to a scale that we can ignore, that $a_0^3 = ga_0h^{-1}ha_0k^{-1}ka_0g^{-1} = ga_0^3g^{-1}$. Hence a_0^3 and g commute. (We will argue similarly several times in what follows.) Two matrices A, B , with B invertible, commute if and only if A commutes with B^{-1} . Since $a_0^3 = a_0^{-1}$, we have g, g^{-1} commute with a_0 . This reasoning holds for h and k as well.

Since g, h, k all commute with a_0 , γ commutes with the \mathbb{Z}_4 action. Since γ preserves rank, the orbit (5) must be fixed. Since γ commutes with the \mathbb{Z}_4 action, it suffices to determine it up to a power of the \mathbb{Z}_4 action. Namely we can expect γ to fix one of the elements of the 4 orbits of rank 2 matrices, e.g., the matrix in (5)

$$(27) \quad M = \begin{pmatrix} 0 & -1 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Case 1: $M = gMh^{-1} = hMk^{-1} = kMg^{-1}$, which implies g, h, k all commute with M^3 (using the same reasoning as above). The only matrices which commute with both M^3 and a_0 are scalar multiples of the identity and we conclude.

Case 2: $a_0Ma_0^{-1} = gMh^{-1} = hMk^{-1} = kMg^{-1}$ which implies $a_0M^3a_0^{-1} = gM^3g$ so $a_0^{-1}gM^3 = (a_0^{-1}g)^{-1}$. Hence $a_0^{-1}g$ commutes with M^3 and a_0 , thus is a scalar multiple of the identity and $g = h = k = a_0$, which has already been accounted for.

The other two cases, like case 2, only with a_0^2 and a_0^3 playing the role of a_0 , are similar. \square

Proposition 5.2. *In the family of decompositions $PGL_3^{\times 3} \rtimes (\mathbb{Z}_3 \rtimes \mathbb{Z}_2) \cdot \mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$, the set of \mathbb{Z}_3^{std} -invariant decompositions is the image of the diagonal PGL_3 -action on $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$ times the standard transpose \mathbb{Z}_2 .*

Proof. We need to show any $\gamma = (g, h, k) \in PGL_3^{\times 3}$ that takes $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$ to another decomposition invariant under the same standard \mathbb{Z}_3 satisfies $g = h = k$.

Since γ fixes rank and there is only one tensor consisting of three matrices of rank 3, $\gamma \cdot a_0^{\otimes 3}$ must be a \mathbb{Z}_3 -fixed point. Thus $ga_0h^{-1} = ha_0k^{-1} = ka_0g^{-1}$. Write $a = ga_0h^{-1}$. This implies

$a^3 = ga_0^3g^{-1} = ha_0^3h^{-1} = ka_0^3k^{-1}$. Equivalently $a^{-3} = ga_0g^{-1}$, and the same for h, k . Thus

$$a = a^{-3}gh^{-1} = gh^{-1}a^{-3}.$$

Equivalently

$$a^4 = gh^{-1}.$$

We have $g^{-1}ag$ is a cube root of a_0^{-1} . Recall that a_0^{-1} has distinct eigenvalues $-1, i, -i$ so all cube roots must likewise have distinct eigenvalues. This uniquely determines the finite choices of eigensystems for our cube roots, yielding 27 possibilities. The \mathbb{Z}_3^{std} action commutes with the diagonal PGL_3 , so we can assume $g = \text{Id}$. Then a is a cube root of a_0^{-1} , so a has order 12. Combining this with the observations $a^4 = h^{-1} = hk^{-1}$ shows $h = a^{-4}$ and $k = a^{-8}$. This leaves us with 27 total candidate restricted families that have a rank three $\mathbb{Z}_4 \times \mathbb{Z}_3$ -fixed summand.

Since g, h, k all commute with $a_0 = a^{-3}$, γ commutes with the \mathbb{Z}_4 action. Consider the action of γ on the two terms in (3). Since the \mathbb{Z}_4 action sends \mathbb{Z}_3 fixed points to \mathbb{Z}_3 fixed points, these summands each map to \mathbb{Z}_3 fixed points.

For γ to send M of (27) to another \mathbb{Z}_3 fixed point we must have

$$Ma^4 = a^{-4}Ma^8 = a^{-8}M = a^4M.$$

Testing all 27 cube roots of a_0^{-1} we see that M commuting with a^{-4} implies a is a scalar times a_0 , and we may assume $a = a_0$, so $a^4 = \text{Id}$. This implies that there are no other \mathbb{Z}_3^{std} -invariant subfamilies in the family. \square

5.2. Symmetries of $\mathcal{S}_{Lader-\mathbb{Z}_3}$. The symmetry group of $G_{\mathcal{S}_{Lader-\mathbb{Z}_3}}$ has already been discussed.

Proposition 5.3. *In Laderman the family of decompositions, the set of \mathbb{Z}_3^{std} -invariant decompositions is the image of the diagonal PGL_3 -action on $\mathcal{S}_{Lader-\mathbb{Z}_3}$ times the standard transpose \mathbb{Z}_2 .*

The proof is very similar to that of Proposition 5.2, so is omitted.

5.3. Symmetries of $\mathcal{S}_{2fix-\mathbb{Z}_3}$.

Proposition 5.4. *The symmetry group of $\mathcal{S}_{2fix-\mathbb{Z}_3}$ is $\Gamma_{\mathcal{S}_{2fix-\mathbb{Z}_3}} = \mathbb{Z}_3$.*

Proof. The incidence and pairing graphs have no joint automorphisms, which shows there are no additional diagonal $GL_3 \subset GL_3^{\times 3}$ symmetries.

To show, despite the symmetry of the graphs, that there is no transpose-like symmetry, i.e., a symmetry of the form $x \otimes y \otimes z \mapsto gx^T k^{-1} \otimes kz^T h^{-1} \otimes hy^T g^{-1}$, first note that the first matrix in (14), call this M_{14a} , must satisfy $gM_{14a}^T = M_{14a}h$ as (14) is the only triple with ranks $(2, 1, 1)$, and similarly (using the \mathbb{Z}_3 -action), $hM_{14a}^T = M_{14a}k$ and $kM_{14a}^T = M_{14a}g$. Moreover the second and third matrices in this triple, call them M_{14b}, M_{14c} must satisfy $hM_{14c}^T k^{-1} = M_{14b}$ and $kM_{14b}^T g^{-1} = M_{14c}$, which forces

$$(28) \quad g = h = k = \begin{pmatrix} s & s & s \\ s & s & 0 \\ s & 0 & 0 \end{pmatrix}$$

and we may normalize $s = 1$. Now apply this to the triple in (17), we get the triple

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix}$$

which is not a triple appearing in the decomposition.

It remains to show there are no additional symmetries coming from $GL_3^{\times 3}$. There are only two \mathbb{Z}_3 fixed points in $\mathcal{S}_{2fix-\mathbb{Z}_3}$, M_{12} appearing in (12) of rank one and M_{13} appearing in (13), so these must be fixed by any $(g, h, k) \in PGL(U) \times PGL(V) \times PGL(W)$ -symmetry. Since both of these matrices are idempotent, we get, by an argument as in the proof of Proposition 5.2,

$$(29) \quad \begin{aligned} M_{12} &= M_{12}^3 = gM_{12}^3g^{-1} = gM_{12}g^{-1}, \\ M_{13} &= M_{13}^3 = gM_{13}^3g^{-1} = gM_{13}g^{-1}. \end{aligned}$$

There is only one tensor with ranks $(2, 1, 1)$ in the decomposition, so this also must be fixed. Since it is \mathbb{Z}_3 invariant, we get the $(1, 2, 1)$ tensor and the $(1, 1, 2)$ tensor also must be fixed. These matrices are $M_{14a}, M_{14b}, M_{14c}$. Then $M_{14a} = gM_{14a}h^{-1}$, $M_{14b} = hM_{14b}k^{-1}$, $M_{14c} = kM_{14c}g^{-1}$. Combining this we get $M_{14a}, M_{14b}, M_{14c}$ commutes with g .

Finally we check that the only matrix which commutes with $M_{12}, M_{13}, M_{14a}, M_{14b}, M_{14c}$ is the identity matrix. \square

Remark 5.5. Remarkably, the group element (28) splits $M_{(3)}$ into the sum of two tensors: T_1 , the sum of (12),(13),(14),(15),(16) which is invariant under the transpose like action, and the sum of the others, call it T_2 , which is sent to a different decomposition of T_2 under the action.

Proposition 5.6. *In the family of decompositions $PGL_3^{\times 3} \rtimes (\mathbb{Z}_3 \rtimes \mathbb{Z}_2) \cdot \mathcal{S}_{2fix-\mathbb{Z}_3}$, the set of \mathbb{Z}_3^{std} -invariant decompositions is the image of the diagonal PGL_3 -action on $\mathcal{S}_{2fix-\mathbb{Z}_3}$ times the standard transpose \mathbb{Z}_2 .*

Proof. The two \mathbb{Z}_3 fixed points M_{12}, M_{13} must be fixed by any symmetry, and as above, for any triple (g, h, k) , equation (29) still holds (in fact for g replaced by h or k as well).

Similarly the triple (14) and its \mathbb{Z}_3 -translates also must be fixed. Then $M_{14a} = gM_{14a}h^{-1}$, $M_{14b} = hM_{14b}k^{-1}$, $M_{14c} = kM_{14c}g^{-1}$. Combining this we get $M_{14a}M_{14b}M_{14c}$ commutes with g . We conclude as above. \square

6. CONFIGURATIONS OF POINTS IN PROJECTIVE SPACE

In the decompositions, vectors appear tensored with other vectors, so they are really only defined up to scale (there is only a ‘‘global scale’’ for each term). This suggests using points in projective space, and only later taking scales into account.

Towards our goal of building new decompositions, we would like to describe existing decompositions in terms of simple building blocks. The standard cyclic \mathbb{Z}_3 invariant decompositions of $M_{(n)}$ naturally come in the restricted families parameterized by the diagonal $PGL_n \subset PGL_n^{\times 3}$. Thus, when we examine, e.g., the points in $\mathbb{P}U = \mathbb{P}^{n-1}$ appearing in the rank one terms in a decomposition, we should really study the set of points up to PGL_n -equivalence, call such a *configuration*. Identifying configurations will also facilitate comparisons between known decompositions.

The simplest configuration is n points in \mathbb{C}^n that form a basis, as occurs with the standard decomposition. All known decompositions of size less than n^3 use more than n points. The next simplest is a collection of $n + 1$ points in general linear position, i.e., a collection of points such that any subset of n of them forms a basis. Call such a *framing* of \mathbb{P}^{n-1} . Just as all bases are PGL_n -equivalent to the standard basis, all framings, as points in projective space, are equivalent

to:

$$\begin{pmatrix} 1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \dots, \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ 1 \end{pmatrix}.$$

We focus on the case of \mathbb{P}^2 .

6.1. The case of \mathbb{P}^2 . The group PGL_3 acts simply transitively on the set of 4-plets of points in \mathbb{P}^2 in general linear position (i.e. such that vectors associated to any three of them form a basis).

Start with any 4-ple of points in general linear position. We will call the following choice, the *default framing*:

$$u_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, u_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, u_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, u_4 = \begin{pmatrix} -1 \\ -1 \\ -1 \end{pmatrix}.$$

Note that u_1, u_2, u_3 is the standard basis and u_4 is chosen such that $u_1 + u_2 + u_3 + u_4 = 0$.

The $\{[u_j]\}$ determine 6 lines in $\mathbb{P}U$, those going through pairs of points, that we consider as points in $\mathbb{P}U^*$.

For the default framing, representatives of these are:

$$\begin{aligned} v_{12} &= (0, 0, 1), & v_{13} &= (0, 1, 0), & v_{14} &= (0, 1, -1), \\ v_{23} &= (-1, 0, 0), & v_{24} &= (-1, 0, 1), & v_{34} &= (1, -1, 0). \end{aligned}$$

Here $[v_{ij}]$ is the line in \mathbb{P}^2 , considered as a point in the dual space \mathbb{P}^{2*} , through the points $[u_i]$ and $[u_j]$ in \mathbb{P}^2 (or dually, the point of intersection of the two lines $[u_i], [u_j]$ in \mathbb{P}^{2*}). Algebraically this means $v_{ij}(u_i) = 0 = v_{ij}(u_j)$.

The choices of scale made here are useful for the decomposition $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$ because they make the \mathbb{Z}_4 action easier to write down. They are such that $v_{i,i+1}(u_{i+2}) = 1$, $v_{i,i+1}(u_{i+3}) = -1$ (indices considered mod four). This has the advantage of $v_{i+1,i+2} = a_0^{-i} v_{12}$ where a_0 is as in (1). For v_{13} there was no obvious choice of sign, and we chose $v_{24} = a_0^{-1}(v_{13})$.

The v_{ij} 's constitute two \mathbb{Z}_4 -orbits: the $v_{i,i+1}$'s which consist of four vectors, and the $v_{i,i+2}$'s of which there are two.

The $v_{i,j}$ in turn determine their new points of intersection:

$$u_{12,34} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, u_{13,24} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, u_{14,23} = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}.$$

These determine further lines

$$v_{(12,34),(13,24)} = (-1, 1, 1), \quad v_{(12,34),(14,23)} = (1, -1, 1), \quad v_{(13,24),(14,23)} = (1, 1, -1),$$

which determine

$$\begin{aligned} u_{34,(13,24|14,23)} &= \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}, & u_{24,(12,34|14,23)} &= \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}, & u_{23,(12,34|13,24)} &= \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix}, \\ u_{12,(12,34|13,24)} &= \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, & u_{13,(12,34|13,24)} &= \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, & u_{14,(12,34|13,24)} &= \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}. \end{aligned}$$




FIGURE 9. Points and lines from the default framing of \mathbb{P}^2 . Original four points are black dots, three new are shaded dots

This process continues, giving rise to an infinite collection of points but in practice only vectors from the first 3 rounds appeared in decompositions.

6.2. Point-line configuration for $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$. The rank one elements appearing in $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$ consist of points from three rounds of points obtained from the default configuration. All points appear except that $u_{13|24}$ is missing (the orbit under \mathbb{Z}_4 of $u_{12|34}$ is $\{u_{12|34}, u_{14|23}\}$).

Here is the decomposition $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$ in terms of the points from §6.1:

$$\begin{aligned}
 (30) \quad & M_{(3)} = -a_0^{\otimes 3} \\
 (31) \quad & + \mathbb{Z}_4 \cdot (u_1 v_{23})^{\otimes 3} \\
 (32) \quad & + \mathbb{Z}_2 \cdot (u_2 v_{23} - u_{12|34} v_{23})^{\otimes 3} \\
 (33) \quad & + \mathbb{Z}_4 \cdot (u_2 v_{24})^{\otimes 3} \\
 (34) \quad & + \mathbb{Z}_3 \times \mathbb{Z}_4 \cdot (u_2 v_{23} \otimes u_4 v_{14} \otimes u_{12|34} v_{13}).
 \end{aligned}$$

6.3. Point-line configuration for $\mathcal{S}_{Lad-\mathbb{Z}_3}$. Thanks to the transpose-like symmetry, it is better to label points in the dual space by their image under transpose rather than annihilators, to make the transpose-like symmetry more transparent. Points:

$$u_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad u_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad u_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad u_{12} = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}, \quad u_{23} = \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}.$$

$$\begin{aligned} v_1 &= (1, 0, 0), \quad v_2 = (0, 1, 0), \quad v_3 = (0, 0, 1), \\ v_{12} &= (1, 1, 0), \quad v_{23} = (0, 1, 1). \end{aligned}$$

This collection of points has a \mathbb{Z}_2 -symmetry generated by τ_{13} which swaps the two lines.




FIGURE 10. Rank one $\mathbb{P}U$ -points appearing in $\mathcal{S}_{Lad-\mathbb{Z}_3}$

In order to express $\mathcal{S}_{Lad-\mathbb{Z}_3}$ in terms of just these points, we write the decomposition in terms of the $\mathbb{Z}_3 \times \mathbb{Z}_2^\zeta$ -orbits.

$$\begin{aligned} (35) \quad M_{\{3\}} &= (u_2 v_2)^{\otimes 3} \\ (36) \quad &+ (u_3 v_3)^{\otimes 3} \\ (37) \quad &+ (u_{12} v_1)^{\otimes 3} \\ (38) \quad &+ (u_1 v_{12})^{\otimes 3} \\ (39) \quad &+ (u_2 v_1 - u_1 v_{12})^{\otimes 3} \\ (40) \quad &+ \mathbb{Z}_3 \cdot (u_1 v_3) \otimes (u_3 v_1) \otimes (u_1 v_1) \\ (41) \quad &+ \mathbb{Z}_3 \cdot (u_{23} v_1) \otimes (u_{12} v_3) \otimes (u_{23} v_3) \\ (42) \quad &+ \mathbb{Z}_3 \cdot (u_3 v_{12}) \otimes (u_1 v_{23}) \otimes (u_3 v_{23}) \\ (43) \quad &+ \mathbb{Z}_3 \cdot (u_2 v_3 - u_{23} v_1) \otimes (u_1 v_2 - u_{12} v_3) \otimes (u_3 v_2 - u_{23} v_3) \\ (44) \quad &+ \mathbb{Z}_3 \times \mathbb{Z}_2^\zeta \cdot (u_{23} v_{12} + u_2 v_3 - u_1 v_{23}) \otimes (u_2 v_3) \otimes (u_3 v_2). \end{aligned}$$

6.4. Point-line arrangement for $\mathcal{S}_{2fix-\mathbb{Z}_3}$. Despite the lack of a transpose-like symmetry, both sets of points are the 6 points corresponding to lines dual to the standard configuration of four points. Again, this illustrates how the decomposition nearly has such a symmetry, and could likely be modified to have such.

7. SPACES OF Γ -INVARIANTS FOR VARIOUS Γ

The space of \mathbb{Z}_3 -invariants in $A^{\otimes 3}$ is $S^3 A \oplus \Lambda^3 A$. To see this, as a $GL(A) \times \mathfrak{S}_3$ -module we have the decomposition $A^{\otimes 3} = S^3 A \otimes [3] \oplus S_{21} A \otimes [21] \oplus \Lambda^3 A \otimes [111]$. Then \mathbb{Z}_3 acts trivially on the trivial representation [3] and trivially on the sign representation [111] (as the three cycle is even), and [21] decomposes into its $e^{\frac{2\pi i}{3}}, e^{\frac{4\pi i}{3}}$ eigenspaces under \mathbb{Z}_3 . Thus, if $\dim A = m$, the




FIGURE 11. Points from $\mathcal{S}_{2\text{fix-}\mathbb{Z}_3}$ rank one matrices appearing in $\mathbb{P}U$ and $\mathbb{P}U^*$, with the latter expressed as lines in $\mathbb{P}U$

space of invariants $(A^{\otimes 3})^{\mathbb{Z}_3}$ has dimension $\frac{1}{3}(m^3 + 2m)$, so restricting the search to \mathbb{Z}_3 -invariant decompositions reduces the search size by about a factor of 3. (In our case, $m = n^2$.)

The \mathbb{Z}_3 -fixed triples all lie in $S^3 A$, so one cannot have all terms of a decomposition individually \mathbb{Z}_3 -fixed. *A-priori* there could be $2 + 3k$ \mathbb{Z}_3 -fixed points for $0 \leq k < 7$. We found decompositions with 2, 5, and 11 \mathbb{Z}_3 -fixed points, i.e., $k = 0, 1, 3$.

7.1. \mathbb{Z}_{n+1} -invariants. Consider the diagonal $\mathbb{Z}_{n+1} \subset PGL_n \subset PGL_n^{\times 3}$ invariants: Each of U^*, V decomposes into n one-dimensional representations for \mathbb{Z}_{n+1} , corresponding to the eigenvalues $\omega, \omega^2, \dots, \omega^n$ where $\omega = e^{\frac{2\pi i}{n+1}}$. Write $(U^*)_j, V_j$ for the eigenspace corresponding to ω^j . Then, adding indices modulo $n + 1$, and letting $u \in [n]$,

$$A_0 = \bigoplus_{t=1}^n U_t^* \otimes V_{n+1-t},$$

$$A_u = \bigoplus_{i=1, i \neq u}^n U_i^* \otimes V_{u-i},$$

so $\dim A_0 = n$, and $\dim A_u = n - 1$, where each A_i is a \mathbb{Z}_{n+1} -isotypic component of A . Note that A_0 is spanned by the powers of a_0 of (1).

The space of \mathbb{Z}_{n+1} invariants in $A \otimes B \otimes C$ is spanned by the spaces $A_\alpha \otimes B_\beta \otimes C_\gamma$ with $\alpha + \beta + \gamma \equiv 0 \pmod{n + 1}$. Let $1 \leq i, j \leq n$, we have the following dimensions:

space	dimension	number of such	total contribution
$A_0 \otimes B_0 \otimes C_0$	n^3	1	n^3
$A_0 \otimes B_j \otimes C_{n+1-j}$ plus cyclic perms	$n(n-1)^2$	$3n$	$3n^2(n-1)^2$
$A_i \otimes B_j \otimes C_{n+1-j-i}$, $i + j \neq n + 1$	$(n-1)^3$	$n(n-1)$	$n(n-1)^4$

Thus

Proposition 7.1. *The space of $\mathbb{Z}_{n+1} \subset PGL_n \subset PGL_n^{\times 3}$ invariants for the diagonal \mathbb{Z}_{n+1} in $\mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2}$ is $n^5 - n^4 + n^3 - n^2 + n$. In particular, when $n = 2, 3, 4$ these dimensions are respectively 22, 183, 820.*

7.2. $\mathbb{Z}_{n+1} \times \mathbb{Z}_3$ -invariants. We look inside $S^3 A \oplus \Lambda^3 A$ for \mathbb{Z}_{n+1} -invariants. Write

$$\begin{aligned} S^3 A = S^3(A_0 + \cdots + A_n) = S^3 A_0 &\oplus \bigoplus_{j=1}^n S^3 A_j \oplus \bigoplus_{j=1}^n A_0 \otimes S^2 A_j \oplus \bigoplus_{j=1}^n A_j \otimes S^2 A_0 \\ &\oplus \bigoplus_{j,k=1, j \neq k}^n A_j \otimes S^2 A_k \oplus \bigoplus_{1 \leq j < k \leq n} A_0 \otimes A_j \otimes A_k \oplus \bigoplus_{1 \leq i < j < k \leq n} A_i \otimes A_j \otimes A_k. \end{aligned}$$

The subspace of \mathbb{Z}_{n+1} -invariants is (here all indices run from 1 to n)

$$\begin{aligned} (S^3 A)^{\mathbb{Z}_{n+1}} = S^3 A_0 &\oplus \bigoplus_{j|3j \equiv 0 \pmod{n+1}} S^3 A_j \oplus \bigoplus_{j|2j \equiv 0 \pmod{n+1}} A_0 \otimes S^2 A_j \oplus 0 \\ &\oplus \bigoplus_{(j,k)|j+2k \equiv 0 \pmod{n+1}; j \neq k} A_j \otimes S^2 A_k \oplus \bigoplus_{i|n+1-i} A_0 \otimes A_i \otimes A_{n+1-i} \oplus \bigoplus_{\substack{(i,j)|i < j < n+1-i-j \\ \text{or } i < j < 2n+2-i-j < n+1}} A_i \otimes A_j \otimes A_{n+1-i-j}. \end{aligned}$$

The dimensions of the summands of the various types are respectively

$$\binom{n+2}{3}, \binom{n+1}{3}, n \binom{n}{2}, 0, (n-1) \binom{n}{2}, n(n-1)^2, (n-1)^3.$$

Similarly, the space of \mathbb{Z}_{n+1} -invariants in $\Lambda^3 A$ is

$$\begin{aligned} (\Lambda^3 A)^{\mathbb{Z}_{n+1}} = \Lambda^3 A_0 &\oplus \bigoplus_{j|3j \equiv 0 \pmod{n+1}} \Lambda^3 A_j \oplus \bigoplus_{j|2j \equiv 0 \pmod{n+1}} A_0 \otimes \Lambda^2 A_j \oplus 0 \\ &\oplus \bigoplus_{(i,j)|i+2j \equiv 0 \pmod{n+1}; i \neq j} A_i \otimes \Lambda^2 A_j \oplus \bigoplus_{i|n+1-i} A_0 \otimes A_i \otimes A_{n+1-i} \oplus \bigoplus_{\substack{(i,j)|i < j < n+1-i-j \\ \text{or } i < j < 2n+2-i-j < n+1}} A_i \otimes A_j \otimes A_{n+1-i-j}. \end{aligned}$$

The dimensions of the summands of the various types are respectively

$$\binom{n}{3}, \binom{n-1}{3}, n \binom{n-1}{2}, 0, (n-1) \binom{n-1}{2}, n(n-1)^2, (n-1)^3.$$

In both cases, the number of terms of each type depends on divisibility properties of n . When $n = 2$, the summands are

$$S^3 A_0 \oplus S^3 A_1 \oplus S^3 A_2 \oplus (A_0 \otimes A_1 \otimes A_2)^S \oplus (A_0 \otimes A_1 \otimes A_2)^\Lambda,$$

for a total dimension of $8 + 2 = 10$. (Here the S, Λ superscripts are whether the factor is in S^3 or Λ^3 .)

When $n = 3$, the summands are

$$\begin{aligned} S^3 A_0 \oplus A_0 \otimes S^2 A_2 \oplus A_2 \otimes S^2 A_1 \oplus A_2 \otimes S^2 A_3 \oplus (A_0 \otimes A_1 \otimes A_3)^S \\ \oplus \Lambda^3 A_0 \oplus A_0 \otimes \Lambda^2 A_2 \oplus A_2 \otimes \Lambda^2 A_1 \oplus A_2 \otimes \Lambda^2 A_3 \oplus (A_0 \otimes A_1 \otimes A_3)^\Lambda, \end{aligned}$$

for a total dimension of $43 + 20 = 63$.

$M_{(3)}$ has a nonzero projection onto each factor except the one-dimensional $\Lambda^3 A_0$.

When $n = 4$ the summands are

$$\begin{aligned} S^3 A_0 \oplus A_1 \otimes S^2 A_2 \oplus A_2 \otimes S^2 A_4 \oplus A_3 \otimes S^2 A_1 \oplus A_4 \otimes S^2 A_3 \oplus (A_0 \otimes A_1 \otimes A_4)^S \oplus (A_0 \otimes A_2 \otimes A_3)^S \\ \oplus \Lambda^3 A_0 \oplus A_1 \otimes \Lambda^2 A_3 \oplus A_2 \otimes \Lambda^2 A_4 \otimes A_3 \otimes \Lambda^2 A_1 \oplus A_4 \otimes \Lambda^2 A_3 \oplus (A_0 \otimes A_1 \otimes A_4)^\Lambda \oplus (A_0 \otimes A_2 \otimes A_3)^\Lambda, \end{aligned}$$

for a total dimension of $164 + 112 = 276$, compared with the naïve search space dimension of $16^3 = 4096$ and the \mathbb{Z}_3 -invariant search space of dimension 1376.

In summary:

Proposition 7.2. *The dimension of the space of $\mathbb{Z}_{n+1} \times \mathbb{Z}_3$ invariants in $(U^* \otimes U)^{\otimes 3}$ is respectively of dimensions 10, 63, and 276 when $n = 2, 3, 4$.*

$\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$	Char. Poly.	Count
symmetric	$t^2(t-1)$	6
	$t+t^2+t^3$	4
	$t-t^2+t^3-1$	1
triples	$\{t^3, t^3, t^3\}$	4

TABLE 1. Characteristic polynomials of matrices appearing in $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$

$\mathcal{S}_{Lader-\mathbb{Z}_3}$	Char. Poly.	Count
symmetric	$t^2(t-1)$	4
	$t+t^2+t^3$	1
triples	$\{t^2(t-1), t^3, t^3\}$	3
	$\{t^3, t^3, t+t^2+t^3\}$	1
	$\{t^3, t^3, t-t^2+t^3-1\}$	2

TABLE 2. Characteristic polynomials of matrices appearing in $\mathcal{S}_{Lader-\mathbb{Z}_3}$

$\mathcal{S}_{2fix-\mathbb{Z}_3}$	Char. Poly.	Count
symmetric	$t(t-1)^2$	1
	$t^2(t-1)$	1
triples	$\{t^3, t^3, t^3\}$	1
	$\{t^2(t-1), t^2(t-1), t+t^2+t^3\}$	1
	$\{t^2(t-1), t^3, t^3\}$	2
	$\{t^2(t-1), t^2(t-1), t^3\}$	2
	$\{t^2(t-1), t^2(t-1), t^2(t+1)\}$	1

TABLE 3. Characteristic polynomials of matrices appearing in $\mathcal{S}_{2fix-\mathbb{Z}_3}$

8. EIGENVALUES

When we deal with a restricted family $PGL_n \cdot \mathcal{S}$, it makes sense to discuss eigenlines and eigenvalues of the terms appearing. These also facilitate determining if two decompositions lie in the same family, beyond the graphs.

REFERENCES

1. Austin R. Benson and Grey Ballard, *A framework for practical parallel fast matrix multiplication*, Proceedings of the 20th ACM SIGPLAN Symposium on Principles and Practice of Parallel Programming (New York, NY, USA), PPOPP 2015, ACM, February 2015, pp. 42–53.
2. Jerome Brachat, Pierre Comon, Bernard Mourrain, and Elias Tsigaridas, *Symmetric tensor decomposition*, Linear Algebra Appl. **433** (2010), no. 11–12, 1851–1872. MR 2736103
3. Jaroslaw Buczyński, Adam Ginensky, and J. M. Landsberg, *Determinantal equations for secant varieties and the Eisenbud-Koh-Stillman conjecture*, J. Lond. Math. Soc. (2) **88** (2013), no. 1, 1–24. MR 3092255
4. Vladimir P. Burichenko, *On symmetries of the strassen algorithm*, CoRR **abs/1408.6273** (2014).
5. ———, *Symmetries of matrix multiplication algorithms. I*, CoRR **abs/1508.01110** (2015).
6. Luca Chiantini, Christian Ikenmeyer, J. M. Landsberg, and Giorgio Ottaviani, *The geometry of rank decompositions of matrix multiplication I: 2x2 matrices*, CoRR **abs/1610.08364** (2016).
7. Hans F. de Groote, *On varieties of optimal algorithms for the computation of bilinear mappings. I. The isotropy group of a bilinear mapping*, Theoret. Comput. Sci. **7** (1978), no. 1, 1–24. MR 0506377
8. ———, *On varieties of optimal algorithms for the computation of bilinear mappings. II. Optimal algorithms for 2 x 2-matrix multiplication*, Theoret. Comput. Sci. **7** (1978), no. 2, 127–148. MR 509013 (81i:68061)

9. Robert Fourer, David M. Gay, and Brian W. Kernighan, *Ampl: A modeling language for mathematical programming*, second ed., 2002.
10. Shmuel Friedland, *Remarks on the symmetric rank of symmetric tensors*, SIAM J. Matrix Anal. Appl. **37** (2016), no. 1, 320–337. MR 3474849
11. Fulvio Gesmundo, *Geometric aspects of iterated matrix multiplication*, J. Algebra **461** (2016), 42–64. MR 3513064
12. J. A. Grochow and C. Moore, *Matrix multiplication algorithms from group orbits*, ArXiv e-prints (2016).
13. Rodney W. Johnson and Aileen M. McLoughlin, *Noncommutative bilinear algorithms for 3×3 matrix multiplication*, SIAM J. Comput. **15** (1986), no. 2, 595–603. MR 837607
14. T. G. Kolda and B. W. Bader, *Tensor decompositions and applications*, SIAM Review **51** (2009), no. 3, 455–500.
15. Julian D. Laderman, *A noncommutative algorithm for multiplying 3×3 matrices using 23 multiplications*, Bull. Amer. Math. Soc. **82** (1976), no. 1, 126–128. MR MR0395320 (52 #16117)
16. J. M. Landsberg, *Geometry and complexity theory*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 2017.
17. J. M. Landsberg and Mateusz Michalek, *On the geometry of border rank decompositions for matrix multiplication and other tensors with symmetry*, SIAM J. Appl. Algebra Geom. **1** (2017), no. 1, 2–19. MR 3633766
18. J.M. Landsberg and Nicolas Ressayre, *Permanent v. determinant: an exponential lower bound assuming symmetry and a potential path towards valiant’s conjecture*, arXiv:1508.05788 (2015).
19. Hwangrae Lee, *Power sum decompositions of elementary symmetric polynomials*, Linear Algebra Appl. **492** (2016), 89–97. MR 3440150
20. A. Sedoglavic, *Laderman matrix multiplication algorithm can be constructed using Strassen algorithm and related tensor’s isotropies*, ArXiv e-prints (2017).
21. Y. Shitov, *A counterexample to Comon’s conjecture*, ArXiv e-prints (2017).
22. A.V. Smirnov, *The bilinear complexity and practical algorithms for matrix multiplication*, Computational Mathematics and Mathematical Physics **53** (2013), no. 12, 1781–1795 (English).
23. Robert J Vanderbei, *LOQO user’s manual-version 3.10*, Optimization methods and software **11** (1999), no. 1-4, 485–514.

APPENDIX A. ADDITIONAL DECOMPOSITIONS

What follows are three additional decompositions with their graphs and eigenvalue tables.

A.1. A decomposition with five \mathbb{Z}_3 -fixed points and no diagonal GL_3 -symmetry. The lack of extra symmetry may be easily deduced from the incidence graph.

$$(45) \quad M_{(3)} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 1 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 1 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}^{\otimes 3}$$

$$(46) \quad + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} -1 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^{\otimes 3}$$

$$(47) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$(48) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & -1 & 1 \\ 0 & -1 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 & -1 \\ -1 & -1 & 1 \\ 0 & -1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}$$

$$(49) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & -1 \\ 1 & 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} -1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$$(50) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & -1 \\ 0 & 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}$$

$$(51) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$$(52) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & -1 & 0 \\ 1 & -1 & 0 \\ 0 & -1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$




FIGURE 12. Incidence graph of Addtl. Dec. #1




FIGURE 13. Pairing graph of Addtl. Dec. #1

Addtl. Dec. #1	Char. Poly.	Count
symmetric	$t^2(t-1)$	4
	$t+t^2+t^3$	1
triples	$\{t^3, t^3, t^3\}$	3
	$\{t^2(t-1), t^2(t+1), t^3\}$	1
	$\{t^2(t-1), t^3, t-t^2+t^3\}$	1
	$\{t^2(t-1), t^2(t-1), t^3\}$	1

TABLE 4. Characteristic polynomials of matrices appearing in Addtl. Dec. #1

A.2. Another decomposition with five \mathbb{Z}_3 -fixed points.




FIGURE 14. Incidence graph of Addtl. Dec. #2




FIGURE 15. Pairing graph of Addtl. Dec. #2

$$(53) \quad M_{(3)} = \begin{pmatrix} 0 & 1 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 0 & 0 & 0 \\ -1 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3}$$

$$(54) \quad + \begin{pmatrix} 0 & -1 & 1 \\ 1 & -1 & -1 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}^{\otimes 3}$$

$$(55) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} -1 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$(56) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ -1 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & -1 & 0 \\ 1 & -1 & -1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$(57) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & -1 & 0 \\ 0 & -1 & 0 \\ 0 & -1 & 0 \end{pmatrix}$$

$$(58) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$(59) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 1 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & -1 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ -1 & 0 & 1 \\ -1 & 0 & 0 \end{pmatrix}$$

$$(60) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & -1 & 1 \\ -1 & -1 & 1 \\ -1 & -1 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

A.3. Another decomposition with 2 \mathbb{Z}_3 -fixed points.

Addtl. Dec. #2	Char. Poly.	Count
symmetric	$t^2(t-1)$	4
	$t+t^2+t^3$	1
triples	$\{t^2(t-1), t^3, t^3\}$	1
	$\{t^3, t^3, t^3\}$	2
	$\{t^2(t+1), t^2(t+1), t^3\}$	2
	$\{t^2(t-1), t^3, t-t^2+t^3\}$	1

TABLE 5. Characteristic polynomials of matrices appearing in Addtl. Dec. #2




FIGURE 16. Incidence graph of Addtl. Dec. #3

$$\begin{aligned}
 (61) \quad M_{(3)} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} \\
 (62) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} \\
 (63) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} -1 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
 (64) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & -1 & -1 \\ 0 & 0 & 0 \end{pmatrix} \\
 (65) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ -1 & -1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \\
 (66) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & -1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 & 0 \\ -1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \\
 (67) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} -1 & 0 & -1 \\ 1 & 0 & 1 \\ -1 & 0 & -1 \end{pmatrix} \\
 (68) \quad &+ \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 & 1 \\ -1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & -1 \end{pmatrix}
 \end{aligned}$$

APPENDIX B. TRIPLETS

For the reader's convenience, we write out all the matrices appearing $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$ and $\mathcal{S}_{Lader-\mathbb{Z}_3^{std}}$.




FIGURE 17. Pairing graph of Addtl. Dec. #3

Addtl. Dec. #3	Char. Poly.	Count
symmetric	$t(t-1)^2$	1
	$t^2(t-1)$	1
triples	$\{t^2(t-1), t^2(t-1), t^2(t+1)\}$	1
	$\{t^2(t-1), t^2(t+1), t^3\}$	2
	$\{t^2(t+1), t^3, t^3\}$	1
	$\{t^3, t^3, t^3\}$	1
	$\{t^2(t-1), t^2(t-1), 2t^2+t^3\}$	1
	$\{t^2(t+1), t^2(t+1), t^3-t^2-t\}$	1

TABLE 6. Characteristic polynomials of matrices appearing in Addtl. Dec. #3

Matrix triplets for $\mathcal{S}_{\mathbb{Z}_4 \times \mathbb{Z}_3}$:

$$(69) \quad M_{(3)} = \begin{pmatrix} 0 & 0 & 1 \\ -1 & 0 & 1 \\ 0 & -1 & 1 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 0 & 0 & 0 \\ -1 & 0 & 1 \\ -1 & 0 & 1 \end{pmatrix}^{\otimes 3}$$

$$(70) \quad + \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 0 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 1 \end{pmatrix}^{\otimes 3}$$

$$(71) \quad + \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 0 & -1 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix}^{\otimes 3}$$

$$(72) \quad + \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 0 & 0 & -1 \\ 1 & 0 & -1 \\ 1 & 0 & -1 \end{pmatrix}^{\otimes 3}$$

$$(73) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 1 & -1 \end{pmatrix}$$

$$(74) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & -1 \\ 0 & 1 & -1 \\ 0 & 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$(75) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} -1 & 1 & 0 \\ -1 & 1 & 0 \\ -1 & 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

$$(76) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & -1 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Matrix triplets for $\mathcal{S}_{Lader-\mathbb{Z}_3}$:

$$(77) \quad M_{(3)} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} -1 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3}$$

$$(78) \quad + \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3} + \begin{pmatrix} 1 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}^{\otimes 3}$$

$$(79) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

$$(80) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & -1 & -1 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$(81) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & -1 & 0 \\ 1 & -1 & -1 \\ 0 & 1 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$(82) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & 1 \\ -1 & 1 & 1 \\ 1 & -1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$(83) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ -1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$(84) \quad + \mathbb{Z}_3^{std} \cdot \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & -1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

PO Box 7311, DEPARTMENT OF COMPUTER SCIENCE, WAKE FOREST UNIVERSITY, WINSTON-SALEM, NC 27109

E-mail address: ballard@wfu.edu

MAX PLANCK INSTITUTE FOR INFORMATICS, SAARLAND INFORMATICS CAMPUS, BUILDING E1.4, D-66123 SAARBRÜCKEN, GERMANY

E-mail address: cikenmey@mpi-inf.mpg.de

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, MAILSTOP 3368, COLLEGE STATION, TX 77843-3368, USA

E-mail address: jml@math.tamu.edu

DEPARTMENT OF MATHEMATICS, UC BERKELEY

E-mail address: nick.ryder@berkeley.edu