


Section 16.1: Vector Fields

Definition: Let D be a set of points in either \mathbb{R}^2 or \mathbb{R}^3 . A vector field is a function \mathbf{F} that assigns to each point in D to a vector.

in \mathbb{R}^2 : $\mathbf{F}(x, y) = \langle P_1(x, y), P_2(x, y) \rangle$

in \mathbb{R}^3 : $\mathbf{F}(x, y, z) = \langle P_1(x, y, z), P_2(x, y, z), P_3(x, y, z) \rangle$

Example: Sketch the vector field $\mathbf{F}(x, y) = \langle 1, 2x \rangle$



$$F(0,0) = \langle 1, 0 \rangle$$

$$F(0,1) = \langle 1, 0 \rangle$$

$$F(0,2) = \langle 1, 0 \rangle$$


$$F(0,*) = \langle 1, 0 \rangle$$

$$F(1,*) = \langle 1, 2 \rangle$$

$$F(2,*) = \langle 1, 4 \rangle$$

$$F(-1,*) = \langle 1, -2 \rangle$$

$$F(-2,*) = \langle 1, -4 \rangle$$



Definition: If f is a scalar function then the gradient of f , $\nabla f = \langle f_x, f_y \rangle$, is called a gradient vector field.

A vector field F is called a conservative vector field if it is the gradient of some scalar function f , i.e. $F(x, y) = \nabla f(x, y)$. The function f is called a potential function for F .

Example: Find the gradient vector field for $f(x, y, z) = x \ln(y - z)$.

$$\nabla f = \langle f_x, f_y, f_z \rangle$$

$$= \left\langle \ln(y-z), x \cdot \frac{1}{y-z}, x \cdot \frac{-1}{y-z} \right\rangle$$

$$\nabla f = \left\langle \ln(y-z), \frac{x}{y-z}, \frac{-x}{y-z} \right\rangle$$