

Two Results on Separating Vectors

W. GONG, D. R. LARSON & W. R. WOGEN

ABSTRACT. We show that a linear space of bounded operators with denumerable Hamel basis has a separating vector if and only if its subspace of finite rank operators has a separating vector, and also that the weakly closed algebra generated by a triangular operator has a separating vector. Moreover, we prove density of the set of all separating vectors.

A *separating vector* for a linear subspace \mathcal{S} of linear transformations on a vector space V is a vector $x \in V$ for which the map $S \rightarrow Sx$ from \mathcal{S} to V is injective. The purpose of this note is to settle two separating vector problems which arose in our earlier work and which have been open for several years.

In Theorem 8 we answer a question in [L], which arose in a study of reflexivity, and show that if \mathcal{S} is a linear subspace of $\mathcal{B}(X)$ for a Banach space X , and if \mathcal{S} has a denumerable Hamel basis, then \mathcal{S} has a separating vector if and only if the linear subspace \mathcal{S}_F of finite rank operators in \mathcal{S} has a separating vector. Moreover, in this case $\text{sep}(\mathcal{S})$ is dense in X , where $\text{sep}(\mathcal{S})$ denotes the set of *all* separating vectors for \mathcal{S} . In the types of applications we have in mind, X will often be an operator algebra.

In Corollary 12 we show that if T is a Hilbert space operator which is triangular in the sense that there is an orthonormal basis $\{e_i\}$, such that $Te_n \in \text{span}\{e_1, \dots, e_n\}$ for each n , then the weak operator topology closed algebra $\mathcal{W}(T)$ generated by T and I has a separating vector. (In fact, we prove the stronger result that $\text{sep}(\mathcal{W}(T))$ is dense, and we obtain this in a somewhat wider setting.) The question of validity of this was raised formally in [LW1] and [HLW], although it had been considered earlier, and was motivated by a counterexample constructed in [W] of a Hilbert space operator S for which $\mathcal{W}(S)$ fails to have a separating vector. Theorems 8 and 11 are related in that the ideas that led to a proof of the first inspired the proof of the second.

We refer the reader to [H] for an extensive exposition of triangular operators, and to [LW1] for a discussion of some open questions. In this article all vector spaces will be assumed complex. If V is a vector space we use $\mathcal{L}(V)$ to denote

all linear transformations on V , and if X is a Banach space, we use $\mathcal{B}(X)$ to denote all *bounded* linear transformations on X . By a *complex line* we mean a one dimensional affine subspace.

We prove a sequence of results leading to Theorem 8. Some have independent interest.

Lemma 1. *Let \mathcal{S} be a linear subspace of $\mathcal{L}(V)$ and let $x \in \text{sep}(\mathcal{S})$. Let $\{S_1, \dots, S_n\}$ be a finite set of nonzero elements of \mathcal{S} . If there is a vector $y \in V$ and distinct scalars $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ such that*

$$S_i(y + \lambda_i x) = 0, \quad i = 1, \dots, n,$$

then $\{S_1, \dots, S_n\}$ is linearly independent.

Proof. Suppose $c_1 S_1 + \dots + c_n S_n = 0$ for scalars $c_i \in \mathbb{C}$. We will show $c_i = 0$, $1 \leq i \leq n$. The hypothesis implies that

$$c_1 S_1(y + \lambda_1 x) + \dots + c_n S_n(y + \lambda_n x) = 0.$$

Rewrite this as

$$(c_1 S_1 + \dots + c_n S_n)y + (c_1 \lambda_1 S_1 + \dots + c_n \lambda_n S_n)x = 0.$$

The first term vanishes by hypothesis so, since x separates \mathcal{S} , this implies that

$$c_1 \lambda_1 S_1 + \dots + c_n \lambda_n S_n = 0.$$

Repeating the argument above with $c_i \lambda_i$ in place of c_i , we obtain

$$c_1 \lambda_1^2 S_1 + \dots + c_n \lambda_n^2 S_n = 0.$$

Proceeding inductively, this implies that

$$c_1 \lambda_1^\ell S_1 + \dots + c_n \lambda_n^\ell S_n = 0$$

for all positive integers ℓ . It follows that

$$c_1 p(\lambda_1) S_1 + c_2 p(\lambda_2) S_2 + \dots + c_n p(\lambda_n) S_n = 0$$

for all complex polynomials $p(z)$. Fix i and choose $p(z)$ so that $p(\lambda_i) = 1$ and $p(\lambda_j) = 0$ for $j \neq i$. This shows that $c_i S_i = 0$, and hence $c_i = 0$ since $S_i \neq 0$. \square

Lemma 2. *Let \mathcal{S} be a linear subspace of $\mathcal{L}(V)$ of finite dimension n . Let $y \in V$, and suppose $x \in \text{sep}(\mathcal{S})$. If there exist distinct nonzero scalars $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ such that $y + \lambda_i x$ fails to separate \mathcal{S} for each $1 \leq i \leq n$, then $y \in \text{sep}(\mathcal{S})$.*

Proof. By hypothesis there are nonzero elements S_1, \dots, S_n of \mathcal{S} such that

$$S_i(y + \lambda_i x) = 0$$

for each $i = 1, \dots, n$. Then $\{S_1, \dots, S_n\}$ are linearly independent by Lemma 1, so span the space \mathcal{S} since \mathcal{S} has dimension n . If $(c_1 S_1 + \dots + c_n S_n) y = 0$ for some $c_i \in \mathbb{C}$, it follows from the equation

$$c_1 S_1(y + \lambda_1 x) + \dots + c_n S_n(y + \lambda_n x) = 0$$

that $(c_1 \lambda_1 S_1 + \dots + c_n \lambda_n S_n)x = 0$. By hypothesis x separates \mathcal{S} , so this implies that $c_1 \lambda_1 S_1 + \dots + c_n \lambda_n S_n = 0$ and hence $c_i \lambda_i = 0$, $1 \leq i \leq n$. Since the λ_i are nonzero, this shows that $c_i = 0$, $1 \leq i \leq n$. Since $(c_1 S_1 + \dots + c_n S_n)$ represents a generic element of \mathcal{S} , this shows that y separates \mathcal{S} .

Proposition 3. *Let \mathcal{S} be a linear subspace of $\mathcal{L}(V)$ of finite dimension n . If L is a complex line in V , then either L contains no separating vector for \mathcal{S} or all but at most n vectors in L separate \mathcal{S} .*

Proof. Suppose $x \in L$ separates \mathcal{S} . By way of contradiction, suppose x_1, \dots, x_{n+1} are distinct elements of L which fail to separate \mathcal{S} . Let $y = x_{n+1}$. Since $y \neq x$ and both lie in L , we have

$$L = \{(1 - \alpha)y + \alpha x : \alpha \in \mathbb{C}\}.$$

For each $1 \leq i \leq n$ let α_i be the unique scalar with $x_i = (1 - \alpha_i)y + \alpha_i x$. For each i we have $\alpha_i \neq 1$ since $x_i \neq x$. Let $\lambda_i = \alpha_i(1 - \alpha_i)^{-1}$, $1 \leq i \leq n$. The λ_i are distinct because the α_i are distinct. We have $y + \lambda_i x = (1 - \alpha_i)^{-1} x_i$ so $y + \lambda_i x$ fails to separate \mathcal{S} , $1 \leq i \leq n$. An application of Lemma 2 now implies that y must separate \mathcal{S} , our contradiction. \square

Lemma 2 and Proposition 3 have generalizations to subspaces \mathcal{S} of algebraic dimension \aleph_0 . No topological assumptions are needed for this, as opposed to the case of some of the results in [L].

Proposition 4. *Let \mathcal{S} be a linear subspace of $\mathcal{L}(V)$ of denumerable Hamel basis. Then:*

- (i) *If x separates \mathcal{S} and $y \in V$ is arbitrary, there is at most countably many complex numbers λ for which $y + \lambda x$ fails to separate \mathcal{S} .*
- (ii) *If L is a complex line in V , then either L contains no separating vectors for \mathcal{S} or all but countably many vectors in L separate \mathcal{S} .*

Proof. Write $\mathcal{S} = \text{span}\{S_n\}_{n=1}^\infty$ and let $\mathcal{S}_n = \text{span}\{S_1, \dots, S_n\}$. If $x \in V$, then x separates \mathcal{S} iff x separates \mathcal{S}_n for all n . So (ii) follows from Proposition 3. Then (i) follows from the fact that for $\lambda \neq -1$, $y + \lambda x = (1 - \alpha)^{-1}[(1 - \alpha)y + \alpha x]$ for $\alpha = \lambda(1 + \lambda)^{-1}$. □

Corollary 5. *Let \mathcal{S} be a linear subspace of continuous linear transformations on a topological vector space X , and suppose \mathcal{S} has a denumerable Hamel basis. Then $\text{sep}(\mathcal{S})$ is either empty or dense in X .*

Proof. If $x \in \text{sep}(\mathcal{S})$ and $y \in X$ is arbitrary, let L be the complex line containing x and y . By Proposition 4, $\text{sep}(\mathcal{S}) \cap L$ is dense in L so its closure contains y . □

Proposition 6. *Let X be a Banach space and let \mathcal{S} be a linear subspace of $\mathcal{B}(X)$ of finite dimension. Then $\text{sep}(\mathcal{S})$ is an open subset of X .*

Proof. We will show that $X \setminus \text{sep}(\mathcal{S})$ is closed. Suppose $x_n \in X \setminus \text{sep}(\mathcal{S})$ and $x_n \rightarrow x$. For each n there exists $S_\ell \in \mathcal{S}$ with $\|S_\ell\| = 1$ such that $S_\ell x_\ell = 0$. By compactness there exists a subsequence $\{S_{n_\ell}\}_{\ell=1}^\infty$ which converges to a nonzero operator $S \in \mathcal{S}$. We have

$$Sx = \lim_{\ell} S_{n_\ell} x_{n_\ell} = 0.$$

So $x \in X \setminus \text{sep}(\mathcal{S})$. □

Lemma 7. *Let X be a Banach space and let $\mathcal{S}_n \subseteq \mathcal{S}_{n+1}$ be an increasing sequence of finite dimensional subspaces. If each \mathcal{S}_n has a separating vector, then $S := \bigcup_{n=1}^\infty \mathcal{S}_n$ has a separating vector.*

Proof. We have $\text{sep}(\mathcal{S}) = \bigcap_{n=1}^\infty \text{sep}(\mathcal{S}_n)$. By Corollary 5 and Proposition 6, $\text{sep}(\mathcal{S}_n)$ is open and dense in X . So, since X is complete, the Baire Category Theorem implies that $\bigcap_{n=1}^\infty \text{sep}(\mathcal{S}_n)$ is nonempty. □

Theorem 8. *Let X be a Banach space and let \mathcal{S} be a linear subspace of $\mathcal{B}(X)$ with denumerable Hamel basis. Then \mathcal{S} has a separating vector if and only if \mathcal{S}_F has a separating vector.*

Proof. Only one direction requires proof. Assume \mathcal{S}_F has a separating vector. By [L, Proposition 2.3] the theorem is true if \mathcal{S} has finite dimension. If \mathcal{S} has infinite dimension, let $\{S_n\}_{n=1}^\infty$ be a countable spanning set for \mathcal{S} and let $\mathcal{S}_n = \text{span}\{S_1, \dots, S_n\}$, $1 \leq n < \infty$. Our hypothesis implies that $(\mathcal{S}_n)_F$ has a separating vector and thus, by the finite dimensional result, that \mathcal{S}_n has a separating vector. Then \mathcal{S} has a separating vector by Lemma 7. □

We can extend the notion of separating vector in the following fashion: If $\mathcal{S} \subseteq \mathcal{L}(V)$, define a (right) *separator* for \mathcal{S} to be an operator T in $\mathcal{L}(V)$ for which the map $S \rightarrow ST$ is injective. Define *left separator* accordingly. (Then \mathcal{S} has a right separator iff \mathcal{S}^* has a left separator.)

Proposition 9. *The following are equivalent:*

- (i) \mathcal{S} has a right separator of rank n .
- (ii) There is an n -tuple (u_1, \dots, u_n) of vectors in V such that no nonzero element of \mathcal{S} annihilates every u_i (a separating n -tuple).
- (iii) The n -fold inflation $\mathcal{S}^{(n)} := \{S \oplus \dots \oplus S : S \in \mathcal{S}\}$ acting on $V^{(n)} = V \oplus \dots \oplus V$ has a separating vector.

Proof. If T is a right separator, then any n -tuple (u_1, \dots, u_n) which spans $\text{ran}(T)$ separates \mathcal{S} , and conversely, given (u_1, \dots, u_n) any T whose range contains $\{u_1, \dots, u_n\}$ will do. It is clear that (u_1, \dots, u_n) separates \mathcal{S} iff the vector $u_1 \oplus \dots \oplus u_n$ separates $\mathcal{S}^{(n)}$. □

Remark 10. Using Proposition 9, most of the previous results easily generalize to separators and separating n -tuples, and relate to inflations. The other results easily adapt using the observation that T is a right separator for \mathcal{S} iff T is a separating vector for the left regular representation of \mathcal{S} on $\mathcal{L}(V)$. The statements of Lemmas 1 and 2 and Propositions 3 and 4 are valid with “separating vector $x \in V$ ” replaced with “right separator $x \in \mathcal{L}(V)$ ” and “vector $y \in V$ ” replaced with “operator $y \in \mathcal{L}(V)$ ”. Here “complex line in V ” should be replaced with “complex line in $\mathcal{L}(V)$.” Let $\text{SEP}^r(\mathcal{S})$ denote the set of right separators of \mathcal{S} in $\mathcal{L}(V)$. Then, for X a Banach space, the analogues of Corollary 5 and Proposition 6 show that for finite dimensional subspaces $\mathcal{S} \subseteq \mathcal{B}(X)$, $\text{SEP}^r(\mathcal{S})$ is always open and dense in $\mathcal{B}(X)$. Part (iii) of Proposition 9 implies that Lemma 7 and Theorem 8 remain valid with “separating vector” replaced with either “separator of rank $\leq n$ ” or “separating n -tuple.”

Recall that a subspace $E \subseteq H$ is *semiinvariant* for an algebra $\mathcal{A} \subseteq \mathcal{B}(H)$ if $E = M \oplus N$ with $M, N \in \text{Lat } \mathcal{A}$ and $N \subseteq M$. The property we use is that if E is semiinvariant, the compression map $A \mapsto PA|_E$ is a homomorphism, where $P = \text{proj}(E)$.

Theorem 11. *Let H be a separable Hilbert space and let $\mathcal{A} \subseteq \mathcal{B}(H)$ be a weakly closed unital operator algebra for which there is an increasing sequence $E_n \subseteq E_{n+1}$ of semiinvariant subspaces with closed union H , such that each compression algebra $\mathcal{A}_n := P_n \mathcal{A}|_{E_n}$ has a separating vector and is finite dimensional, where $P_n = \text{proj}(E_n)$. Then $\text{sep}(\mathcal{A})$ is dense in H . So in particular, $\text{sep}(\mathcal{A})$ is nonempty.*

Proof. Suppose $\text{sep}(\mathcal{A})$ fails to be dense. Then there exists a closed ball J of positive radius with $J \cap \text{sep}(\mathcal{A}) = \emptyset$. For each pair of positive integers n, m let

$$\mathcal{K}_{nm} := \{y \in J : \exists A \in \mathcal{A}, Ay = 0, \|P_n A P_n\| = 1, \|A\| \leq m\}.$$

Then our assumption implies that $\bigcup_{n,m=1}^\infty \mathcal{K}_{nm} = J$. We first show that \mathcal{K}_{nm} is closed. Suppose $y_k \in \mathcal{K}_{nm}, y_k \rightarrow y$. For each k let A_k be an operator in \mathcal{A} such that $A_k y_k = 0, \|A_k\| \leq m, \|P_n A_k P_n\| = 1$. By weak compactness and metrizable-ability of the unit ball of $\mathcal{B}(H)$, there is a subsequence $\{A_{k_\ell}\}$ which converges weakly to an operator $A \in \mathcal{A}$. We have $\|A\| \leq m$. For each n , the subsequence $P_n A_{k_\ell}|_{E_n}$ converges weakly to $P_n A|_{E_n}$, and since \mathcal{A}_n is finite dimensional this convergence is in *norm*. So $\|P_n A P_n\| = 1$. We have $\|A_{k_\ell} y\| = \|A_{k_\ell}(y - y_{k_\ell})\| \leq m\|y - y_{k_\ell}\|$ so $A_{k_\ell} y \rightarrow 0$. Hence $\|Ay\|^2 = \langle Ay, Ay \rangle = \text{Lim}_\ell \langle A_{k_\ell} y, Ay \rangle = 0$. This shows that $y \in \mathcal{K}_{nm}$ as required.

Since the sets \mathcal{K}_{nm} are closed and $\bigcup_{n,m} \mathcal{K}_{nm} = J$, by the Baire Category Theorem some \mathcal{K}_{nm} must contain a ball $B_\epsilon(y) := \{x \in H: \|x - y\| < \epsilon\}$. Density for $\bigcup_n P_n H$ implies that $B_\epsilon(y) \cap E_q$ is not empty for some $q \geq n$. Since \mathcal{A}_q has a separating vector and is a finite dimensional subalgebra of $\mathcal{B}(E_q)$, Corollary 5 implies that $\text{sep}(\mathcal{A}_q)$ is dense in E_q . Hence $B_\epsilon(y) \cap \text{sep}(\mathcal{A}_q)$ is nonempty. Fix $z \in B_\epsilon(y) \cap \text{sep}(\mathcal{A}_q)$. We have $\mathcal{A}_q \neq \{0\}$ so $z \neq 0$. Since $z \in \mathcal{K}_{nm}$ there exists $A \in \mathcal{A}, Az = 0, \|P_n A P_n\| = 1$. Then $\|P_q A P_q\| \geq 1$. So $P_q A|_{E_q}$ is a nonzero element of \mathcal{A}_q which annihilates z . This is our required contradiction. \square

Theorem 11 yields an affirmative answer to Question 1 in [HLW].

Corollary 12. *Let H be a separable Hilbert space, and let $T \in \mathcal{B}(H)$ be a triangular operator. Then $\text{sep}(\mathcal{W}(T))$ is dense in H . More generally, if $\text{Lat}(T)$ contains a complete multiplicity one nest order isomorphic to a subset of the extended integers, then $\text{sep}(\mathcal{W}(T))$ is dense in H .*

Proof. It is well known that a singly generated algebra acting on a finite dimensional space has a separating vector. So if T has a triangular basis $\{e_n\}$, then the invariant subspaces $E_n = [e_1, \dots, e_n]$ satisfy the hypotheses of Theorem 11 for $\mathcal{A} = \mathcal{W}(T)$. Similarly, if $\text{Lat}(T)$ contains a nest of the prescribed type, then there is a sequence of finite dimensional interval projections for the nest which satisfy the hypotheses of Theorem 11 for $\mathcal{W}(T)$. \square

Remark 13. As pointed out in [HLW], a Hilbert space operator T is triangular iff the set of algebraic vectors for T is dense in H , where a vector x is called algebraic for T if $p(T)x = 0$ for some nontrivial polynomial $p(z)$. This makes sense in a separable Banach space setting. Density of the algebraic vectors for $T \in \mathcal{B}(X)$ is equivalent to the existence of a sequence $\{x_n\}$ in X with dense span such that $E_n := [x_1, \dots, x_n]$ is invariant under T for each n , so this is a natural extension of the notion of triangularity. If X is reflexive Corollary 12 extends to this case. To adapt the proof of Theorem 11 let $\{P_n\}$ be a commuting sequence of bounded projections from X onto the finite dimensional spaces $\{E_n\}$. Although $\{P_n\}$ may not be uniformly bounded, this poses no obstruction, and the proof carries through with only minor modifications.

Questions. Our proofs suggest some new directions. Suppose $\text{sep}(\mathcal{S}) \neq \emptyset$. If L is a complex line meeting $\text{sep}(\mathcal{S})$, define

$$i(\mathcal{S}; L) = \text{card}\{y \in L: y \notin \text{sep}(\mathcal{S})\},$$

and for $x \in \text{sep}(\mathcal{S})$ define

$$j(\mathcal{S}; x) = \sup\{i(\mathcal{S}; L): L \text{ is a complex line containing } x\}.$$

Then define $k(\mathcal{S}) = \sup_x j(\mathcal{S}; x)$. This gives a way of associating natural *indices of separation* to lines meeting $\text{sep}(\mathcal{S})$, points in $\text{sep}(\mathcal{S})$, and to subspaces \mathcal{S} . By Proposition 3, $k(\mathcal{S}) \leq \dim(\mathcal{S})$. It might be worthwhile to investigate this further. If T acts on a finite dimensional space, it is not hard to show that $k(\mathcal{W}(T))$ is the cardinality of the spectrum of T . So the index $k(\mathcal{S})$ somehow generalizes this concept. For another direction, let us say that a nonempty set \mathcal{E} of vectors is *linearly dense* if $\mathcal{E} \cap L$ is dense in L for all complex lines L that meet \mathcal{E} . This is a stronger property than density. Proposition 4 implies that if \mathcal{S} is a linear subspace of operators with denumerable Hamel basis, then $\text{sep}(\mathcal{S})$ is either empty or linearly dense. Also, it is not hard to show (see [LW2], Lemma 8) that if \mathcal{S} is a m.a.s.a. on a separable Hilbert space H , then $\text{sep}(\mathcal{S})$ is linearly dense in H . Is linear density common? More to the point, if T is triangular, is $\text{sep}(\mathcal{W}(T))$ linearly dense?

REFERENCES

- [H] D. A. HERRERO, *Triangular operators*, Bull. London Math. Soc. **23** (1991), 513–554.
- [HLW] D. A. HERRERO, D. R. LARSON & W. R. WOGEN, *Semitriangular operators*, Houston J. Math. **17** (1991), 477–499.
- [L] D. R. LARSON, *Reflexivity, algebraic reflexivity and linear interpolation*, American J. Math. **110** (1988), 283–299.
- [LW1] D. R. LARSON & W. R. WOGEN, *Some problems on triangular and semitriangular operators*, Contemporary Mathematics **120** (1991), 97–100.
- [LW2] ———, *Extensions of normal operators*, (Preprint).
- [W] W. R. WOGEN, *Some counterexamples in nonselfadjoint algebras*, Ann. of Math. **126** (1987), 415–427.

The second author was partially supported by NSF Grant DMS-9107137.

W. GONG
Department of Mathematics
Qufu Normal University
Shandong, China

D. R. LARSON
Department of Mathematics
Texas A&M University
College Station, Texas 77843
larson@math.tamu.edu

W. R. WOGEN
Department of Mathematics
University of North Carolina
Chapel Hill, North Carolina 27599
wrw@math.unc.edu

Received: January 18, 1994.