

# L'Enseignement Mathématique

**DeVore, Ronald**

*MULTIPLIERS OF UNIFORM CONVERGENCE*

L'Enseignement Mathématique, Vol.14 (1968)

PDF erstellt am: Nov 26, 2008

## **Nutzungsbedingungen**

Mit dem Zugriff auf den vorliegenden Inhalt gelten die Nutzungsbedingungen als akzeptiert. Die angebotenen Dokumente stehen für nicht-kommerzielle Zwecke in Lehre, Forschung und für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrücke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und unter deren Einhaltung weitergegeben werden. Die Speicherung von Teilen des elektronischen Angebots auf anderen Servern ist nur mit vorheriger schriftlicher Genehmigung des Konsortiums der Schweizer Hochschulbibliotheken möglich. Die Rechte für diese und andere Nutzungsarten der Inhalte liegen beim Herausgeber bzw. beim Verlag.

## **SEALS**

Ein Dienst des *Konsortiums der Schweizer Hochschulbibliotheken*  
c/o ETH-Bibliothek, Rämistrasse 101, 8092 Zürich, Schweiz

[retro@seals.ch](mailto:retro@seals.ch)

<http://retro.seals.ch>

# MULTIPLIERS OF UNIFORM CONVERGENCE

by Ronald DeVORE

1. *Introduction.* If  $A$  and  $B$  are two classes of  $2\pi$ -periodic integrable functions we say that  $(\lambda_k)$  is a multiplier sequence from  $A$  into  $B$  and we write  $(\lambda_k) \in (A, B)$  if whenever

$$\sum_0^{\infty} (a_n \cos nx + b_n \sin nx)$$

is the Fourier series of a function in  $A$

$$\sum_0^{\infty} \lambda_n (a_n \cos nx + b_n \sin nx)$$

is the Fourier series of a function in  $B$ . Let  $C$  denote the class of  $2\pi$ -periodic continuous functions and  $C_F$  the subclass of those functions in  $C$  whose Fourier series converges uniformly. Karamata [1] has shown that  $(\lambda_k) \in (C, C_F)$  if and only if

$$(1.1) \quad \int_0^{2\pi} |A_n(t)| dt = O(1) \quad (n \rightarrow \infty)$$

where

$$A_n(t) = \sum_0^n \lambda_k \cos kt.$$

This theorem contains as a special case an earlier result of Tomić [2] who showed that if  $(\lambda_k)$  is monotone decreasing and convex (i.e.  $\Delta^2 \lambda_k = \lambda_k - 2\lambda_{k-1} + \lambda_{k-2} \geq 0$ ) or more generally quasi-convex (i.e.  $\sum_0^{\infty} (k+1) |\Delta^2 \lambda_k| < \infty$ ) then  $(\lambda_k) \in (C, C_F)$  if and only if  $\lambda_n \log n = O(1)$  ( $n \rightarrow \infty$ ).

It is interesting to see to what extent condition (1.1) can be relaxed if we restrict our attention to a sub-class of  $C$  determined by some structural property. For example, let  $\omega$  be a modulus of continuity and  $C_\omega$  the subclass of  $C$  consisting of those functions whose modulus of continuity  $\omega(f, h)$  satisfies

$$\omega(f, h) = O(\omega(h)) \quad (h \rightarrow 0).$$

Then Tomić [3] has shown that for a quasi-convex sequence  $(\lambda_k)$  to be in  $(C_\omega, C_F)$  it is sufficient that

$$(1.2) \quad \omega \left( \frac{1}{n} \right) \lambda_n \log n = o(1) \quad (n \rightarrow \infty).$$

Also Bojanic [4] has shown that sufficient conditions for  $(\lambda_k)$  to be in  $(C_\omega, C_F)$  are

$$(1.3) \quad \int_0^{2\pi} \left| \sum_0^n A_k(t) \right| dt = O(n) \quad (n \rightarrow \infty)$$

and

$$(1.4) \quad \omega \left( \frac{1}{n} \right) \int_0^{2\pi} |A_n(t)| dt = o(1) \quad (n \rightarrow \infty).$$

Of course, condition (1.3) is equivalent to  $(\lambda_k)$  being a Fourier Stieljes sequence which in particular characterizes the class of multipliers  $(C, C)$ .

No necessary conditions have been given for  $(\lambda_k)$  to be in  $(C_\omega, C_F)$  and sufficient conditions have been restricted to quasi-convex and Fourier-Stieljes sequences. In order to obtain necessary and sufficient conditions for  $(\lambda_k)$  to be in  $(C_\omega, C_F)$ , it is natural to attempt to make  $C_\omega$  a Banach space in which trigonometric polynomials are dense and then invoke the Banach-Steinhaus theorem as Karamata did in characterizing  $(C, C_F)$ . The most natural norm is to define for  $f \in C_\omega$

$$\|f\|_\omega = \max \left( \|f\|_\infty, \sup_{h>0} \frac{\omega(f, h)}{\omega(h)} \right)$$

where  $\|f\|_\infty$  is the usual supremum norm.

The normed space  $(C_\omega, \|\cdot\|_\omega)$  is a Banach space. However, trigonometric polynomials are not dense in  $(C_\omega, \|\cdot\|_\omega)$ . For if  $\omega(h) \neq O(h)$  ( $h \rightarrow 0$ ), then whenever  $(T_n)$  is a sequence of trigonometric polynomials which converge in  $\|\cdot\|_\omega$  to  $f$ ,  $f$  satisfies

$$\omega(f, h) = o(\omega(h)) \quad (h \rightarrow 0).$$

In the case that  $\omega(h) = O(h)$  ( $h \rightarrow 0$ ), then a sequence of trigonometric polynomials  $(T_n)$  converge in  $\|\cdot\|_\omega$  if and only both  $T_n$  and  $T'_n$  converge uniformly and therefore  $f$  is the limit of the sequence  $(T_n)$  only if  $f$  is contin-

uously differentiable. Accordingly, when  $\omega(h) \neq O(h)$  ( $h \rightarrow 0$ ), we define  $c_\omega$  as the class of those functions in  $C_\omega$  for which

$$\omega(f, h) = o(\omega(h)) \quad (h \rightarrow 0)$$

and when  $\omega(h) = O(h)$  ( $h \rightarrow 0$ ) we define  $c_\omega$  as the class of all continuously differentiable functions.  $c_\omega$  is then a closed subspace of  $C_\omega$  and it is easy to see that if  $f \in c_\omega$ , the Fejer sums of  $f$

$$\sigma_n(f) = \int_0^{2\pi} f(t) F_n(t-x) dt$$

with

$$F_n(t) = \frac{1}{2\pi(n+1)} \left( \frac{\sin(n+1)\frac{1}{2}t}{\sin\frac{1}{2}t} \right)^2$$

converges in  $\|\cdot\|_\omega$  to  $f$ . Thus,  $c_\omega$  is precisely the closure of the class of trigonometric polynomials in  $\|\cdot\|_\omega$ . It therefore appears some what more natural to consider the class  $c_\omega$  rather than the class  $C_\omega$  in terms of problems involving multiplier sequences. For we then have

PROPOSITION 1. *The sequence  $(\lambda_k) \in (c_\omega, C_F)$  if and only if*

$$\| \| A_n \| \|_\omega \equiv \sup_{\substack{f \in c_\omega \\ \|f\|_\omega \leq 1}} \left\| \int_0^{2\pi} f(t) A_n(t-x) dt \right\|_\omega = O(1) \quad (n \rightarrow \infty).$$

This is an immediate application of the Banach-Steinhaus theorem [5, p. 60] and the fact that the operators

$$L_n(f)(x) = \int_0^{2\pi} f(t) A_n(t-x) dt$$

converge in  $\|\cdot\|_\omega$  for each trigonometric polynomial  $T$ .

We shall find it convenient to use the following proposition which follows immediately from the fact that any function  $f$  in  $C_\omega$  with  $\|f\|_\omega \leq 1$  is the uniform limit of sequence of functions from the unit ball of  $(c_\omega, \|\cdot\|_\omega)$  (e.g.  $\sigma_n(f)$  provides such a sequence of functions).

PROPOSITION 2. *If  $A(t)$  is an integrable function then*

$$\| \| A \| \|_\omega = \sup_{\substack{f \in C_\omega \\ \|f\|_\omega \leq 1}} \left\| \int_0^{2\pi} f(t) A_n(t-x) dt \right\|_\omega$$

In section 2, we shall consider quasi-convex sequences and show that in this case  $(\lambda_k) \in (c_\omega, C_F)$  if and only if

$$\lambda_n \omega\left(\frac{1}{n}\right) \log n = O(1) \quad (n \rightarrow \infty).$$

In section 3, we shall give a necessary condition that  $(\lambda_k)$  be in  $(c_\omega, C_F)$  with no restrictions on  $(\lambda_k)$ . We shall show that  $(\lambda_k) \in (c_\omega, C_F)$  only if

$$\omega\left(\frac{1}{n}\right) \int_0^{2\pi} |A_n(t)| dt = O(1) \quad (n \rightarrow \infty).$$

It is easy to see that this condition is in general not sufficient. For example, if  $\omega(h) = h$ , then simple integration by parts (see theorem 4.2) shows that

$$\| \| A_n \| \|_\omega = \int_0^{2\pi} \left| \int_0^t A_n(x) dx \right| dt + O(1) \quad (n \rightarrow \infty)$$

thus, if we let

$$\lambda_n = \begin{cases} n, & n = 2^k \\ 0, & n \neq 2^k \end{cases} \quad k = 0, 1, 2, \dots$$

then

$$\int_0^{2\pi} |A_n(t)| dt = \int_0^{2\pi} \left| \sum_0^{[\log_2 n]} 2^k \cos 2^k t \right| dt = O(n) \quad (n \rightarrow \infty).$$

Whereas,

$$\int_0^{2\pi} \left| \int_0^t A_n(x) dx \right| dt = \int_0^{2\pi} \left| \sum_0^{[\log_2 n]} \sin 2^k t \right| dt$$

and it follows from a theorem of Helson [6] that

$$\int_0^{2\pi} \left| \int_0^t A_n(x) dx \right| dt \neq O(1) \quad (n \rightarrow \infty).$$

In section 4, we shall examine sufficient conditions for  $(\lambda_k)$  to be in  $(c_\omega, C_F)$ . First we shall obtain the result analogous to that of Bojanic. In particular, using the necessary condition given in Section 3, we shall prove that if  $(\lambda_k)$  is a Stieltjes sequence then  $(\lambda_k) \in (c_\omega, C_F)$  if and only if

$$\omega\left(\frac{1}{n}\right) \int_0^{2\pi} |A_n(t)| dt = O(1) \quad (N \rightarrow \infty)$$

Finally, we shall give a sufficient condition for  $(\lambda_k)$  to be in  $(c_\omega, C_F)$  with no restrictions on  $(\lambda_k)$ . We shall show that  $(\lambda_k) \in (c_\omega, C_F)$  if

$$(1.5) \quad \omega(\mu_n) \int_0^{2\pi} |A_n(t)| dt = O(1)$$

where

$$\mu_n = \frac{\int_0^{2\pi} \left| \int_0^t A_n(x) dx \right| dt}{\int_0^{2\pi} |A_n(t)| dt}.$$

This condition is also necessary in the case that  $\omega(h) = O(h)$  ( $h \rightarrow 0$ ). However, it is generally not necessary. For example, if  $F(x)$  is the classical Lebesgue function (see [7, p. 195]), then  $F(x) - \frac{x}{2\pi}$  is continuous, of

bounded variation, and its Fourier coefficients are not  $o\left(\frac{1}{n}\right)$  ( $n \rightarrow \infty$ ). Thus,

if  $(\lambda_k)$  is the sequence of Fourier-Stieljes coefficients of  $d\left(F(t) - \frac{t}{2\pi}\right)$  we have using the theorem of Dirichlet-Jordan [7, p. 57] that

$$\lim_{n \rightarrow \infty} \int_0^{2\pi} \left| \sum_0^n \frac{\lambda_k}{k} \sin kt \right| dt = \int_0^{2\pi} \left| F(t) - \frac{t}{2\pi} \right| dt > 0.$$

while by the result of Helson [6]

$$\int_0^{2\pi} \left| \sum_0^n \lambda_k \cos kt \right| dt \neq O(1) \quad (n \rightarrow \infty).$$

Also,

$$\int_0^{2\pi} \left| \sum_0^n \lambda_k \cos kt \right| dt = O(\log n) \quad (n \rightarrow \infty)$$

since it is a Fourier-Stieljes series. So that, if we choose  $\omega$  to satisfy the conditions

$$\omega\left(\frac{1}{n}\right) \int_0^{2\pi} \left| \sum_0^n \lambda_k \cos kt \right| dt = O(1) \quad (n \rightarrow \infty)$$

and

$$\omega(\mu_n) \int_0^{2\pi} \left| \sum_0^n \lambda_k \cos kt \right| dt \neq O(1) \quad (n \rightarrow \infty)$$

with

$$\mu_n = \frac{\int_0^{2\pi} \left| \sum_0^{2n} \frac{\lambda_k}{k} \sin kt \right| dt}{\int_0^{2\pi} \sum_0^n \lambda_k \cos kt \left| dt \right|}$$

we see that (1.5) is in general not necessary.

Although, we give necessary and sufficient conditions for  $(\lambda_k)$  to be in  $(C_\omega, C_F)$  in the case that  $(\lambda_k)$  is quasi-convex or a Stieljes sequence in general no conditions that are both necessary and sufficient are known.

2. *Quasi-convex sequences.* We consider first the simplest case of quasi convex sequences. If we apply Abel summation twice we find

$$A_n(t) = \sum_0^n (k+1) \Delta^2 \lambda_k F_k(t) + n \Delta \lambda_{n-1} F_n(t) + \lambda_n D_n(t)$$

where  $D_n$  is the Dirichlet kernel

$$D_n(t) = \frac{1}{2\pi} \frac{\sin((n + \frac{1}{2})t)}{\sin \frac{1}{2}t}.$$

From the quasi-convexity and the fact that  $\int_0^{2\pi} |F_n(t)| dt = 1$ , we have

$$||| \sum_0^n (k+1) \Delta^2 \lambda_k F_k |||_\omega \leq \int_0^{2\pi} \left| \sum_0^n (k+1) \Delta^2 \lambda_k F_k(t) \right| dt = O(1) \quad (n \rightarrow \infty)$$

for any modulus of continuity  $\omega$ . Thus

$$(2.1) \quad ||| A_n |||_\omega = O(1) + ||| n \Delta \lambda_{n-1} F_n + \lambda_n D_n |||_\omega \quad (n \rightarrow \infty)$$

It follows from standard estimates that there exist positive constants  $C_1, C_2$  such that

$$(2.2) \quad C_1 \omega\left(\frac{1}{n}\right) \log n \leq ||| D_n |||_\omega \leq C_2 \omega\left(\frac{1}{n}\right) \log n.$$

This result is contained in theorems (3.1) and (4.1) so we shall not supply an independent proof.

The main result of this section is

THEOREM 2.1. *If  $(\lambda_k)$  is a quasi-convex sequence then  $(\lambda_k) \in (c_\omega, C_F)$  if and only if*

$$(2.1) \quad \lambda_n \omega \left( \frac{1}{n} \right) \log n = O(1) \quad (n \rightarrow \infty).$$

Proof: We first consider the case when  $(\lambda_n)$  is a bounded sequence. Then by a result of Tomić [3]

$$n \Delta \lambda_{n-1} = o(1).$$

Thus from (2.1) we have

$$||| A_n |||_\omega = O(1) + ||| \lambda_n D_n |||_\omega$$

and the theorem follows immediately from the inequalities (2.2).

We shall now show that the case  $(\lambda_k)$  unbounded does not arise. Tomić [3] has shown that if  $(\lambda_k)$  is quasi convex and unbounded then

$$(2.3) \quad \lambda_n = An + B + o(1) \quad (n \rightarrow \infty)$$

and

$$(2.4) \quad n \Delta \lambda_{n-1} = -An + o\left(\frac{1}{n}\right). \quad (n \rightarrow \infty)$$

thus if

$$\lambda_n \omega \left( \frac{1}{n} \right) \log n = O(1) \quad (n \rightarrow \infty)$$

we must have

$$\frac{\lambda_n}{n} \log n = O(1) \quad (n \rightarrow \infty)$$

and therefor  $(\lambda_n)$  cannot satisfy (2.3) and the conditions (2.1) and  $(\lambda_k)$  unbounded are not compatible. Secondly, if  $(\lambda_k)$  is unbounded then by virtue of (2.1)

$$||| A_n |||_\omega = O(1) + ||| n \Delta \lambda_{n-1} F_n + \lambda_n D_n |||_\omega$$

and thus by (2.2) (2.3), and (2.4) we must have

$$(2.5) \quad ||| A_n |||_\omega \geq An - A C_2 n \omega \left( \frac{1}{n} \right) \log n.$$



For  $\omega(h) = h$ , (2.5) fails and thus  $(\lambda_k) \notin (c_\omega, C_F)$  for any  $\omega$ . Thus,  $(\lambda_k)$  unbounded and  $(\lambda_k) \in (c_\omega, C_F)$  are also incompatible.

3. *A necessary condition for  $(\lambda_k)$  to be in  $(c_\omega, C_F)$ .* In this section, we shall give a necessary condition for  $(\lambda_k)$  to be in  $(c_\omega, C_F)$ . Our main result is the following theorem.

**THEOREM 3.1.** *There exists an absolute constant  $C > 0$  such that for any trigonometric polynomial  $T$  of degree  $n$  we have*

$$\|T\|_\omega \geq C\omega\left(\frac{1}{n}\right) \int_0^{2\pi} |T| dt \quad n = 1, 2, \dots$$

An immediate corollary of this theorem and Proposition 1 is

**COROLLARY 3.1.** *A necessary condition for the sequence  $(\lambda_k)$  to be in  $(c_\omega, C_F)$  is that*

$$\omega\left(\frac{1}{n}\right) \int_0^{2\pi} |A_n| dt = O(1), \quad (n \rightarrow \infty)$$

We shall need some preliminary results concerning representations of trigonometric polynomials. Let  $x_k = \frac{2k\pi}{3n}$ ,  $k = 0, 1, 2, \dots, 3n-1$ . Then if  $T$  is a trigonometric polynomial of degree  $n$ , we have (see [8, p. 33])

$$(3.1) \quad T(x) = \frac{2}{3n} \sum_0^{3n-1} T(x_k) K_n(x - x_k)$$

where

$$(3.2) \quad K_n(t) = \frac{1}{\pi} \frac{\sin\left(\frac{3n}{2}t\right) \sin\left(\frac{n}{2}t\right)}{2n \left(\sin\frac{t}{2}\right)^2}.$$

Also [8, p. 33]

$$(3.3) \quad \int_0^{2\pi} |T(x)| dx \leq \frac{1}{n} \sum_0^{3n-1} |T(x_k)|.$$

Now to the proof of theorem (3.1). Let  $0 < \delta < \frac{1}{4}$ . We wish to estimate

$$\int_{-\frac{\pi\delta}{3n}}^{\frac{\pi\delta}{3n}} K_n(t) dt$$

from below. We have for  $|t| \leq \frac{\pi\delta}{3n}$

$$K_n(t) \geq \frac{1}{\pi} \left( \frac{\binom{2}{\frac{\pi}{3n}} \binom{3nt}{2} \binom{2}{\frac{\pi}{3n}} \binom{nt}{2}}{2n \left(\frac{t}{2}\right)^2} \right) = \frac{6}{\pi^3} n.$$

So that,

$$(3.4) \quad \int_{-\frac{\pi\delta}{3n}}^{\frac{\delta\pi}{3n}} K_n(t) dt \geq \frac{6n}{\pi^3} \cdot \frac{2\pi\delta}{3n} = \frac{4}{\pi^2} \delta.$$

Secondly, for  $k \neq 0$  we estimate  $\int_{x_k - \frac{\pi}{3n}}^{x_k + \frac{2\pi\delta}{3n}} K_n(t) dt$  from above. For

$|t - x_k| \leq \frac{2\pi\delta}{3n}$ , we have

$$K_n(t) \leq \frac{\sin \frac{\delta\pi}{2}}{2n \left(\frac{2\pi}{3n} \left(k - \frac{1}{2}\right)\right)^2} \leq \frac{\delta\pi}{4n} \frac{1}{\left(\frac{2\pi}{3n} \left(k - \frac{1}{2}\right)\right)^2} = \frac{9\delta}{8\pi} \frac{n}{\left(k - \frac{1}{2}\right)^2}.$$

Thus

$$(3.5) \quad \int_{x_k - \frac{2\pi\delta}{3n}}^{x_k + \frac{2\pi}{3n}} |K_n(t)| dt \leq \frac{4\delta\pi}{3n} \cdot \frac{9\delta}{8\pi} \frac{n}{\left(k - \frac{1}{2}\right)^2} = \frac{3}{2} \frac{\delta^2}{\left(k - \frac{1}{2}\right)^2}.$$

Let  $g_\delta(x)$  be the  $2\pi$ -periodic continuous function which has the value one on the interval  $\left[\frac{-\pi\delta}{3n}, \frac{\pi\delta}{3n}\right]$  has the value zero on  $[-\pi, \pi] - \left[\frac{-2\pi\delta}{3n}, \frac{2\pi\delta}{3n}\right]$  and is linear on the intervals  $\left[\frac{-\pi\delta}{3n}, \frac{-\pi\delta}{3n}\right]$  and  $\left[\frac{\pi\delta}{3n}, \frac{2\pi\delta}{3n}\right]$ .

The function

$$\bar{g}_\delta(x) = \omega \left(\frac{\delta\pi}{3n}\right) \sum_{k=0}^{3n-1} \text{Sgn}(T(x_k)) g_\delta(x - x_k)$$

is in  $C_\omega$  and  $\|\bar{g}_\delta\|_\omega \leq 1$ . Also,

$$T(x_k) \int_0^{2\pi} \bar{g}_\delta(x) K_n(x-x_k) dx \geq \omega\left(\frac{\delta\pi}{3n}\right) |T(x_k)| \int_{x_k - \frac{\pi\delta}{3n}}^{x_k + \frac{\pi\delta}{3n}} |K_n(x-x_k)| dx$$

$$- \omega\left(\frac{\delta\pi}{3n}\right) |T(x_k)| \sum_{\substack{j=0 \\ j \neq k}}^{3n-1} \int_{x_j - \frac{2\pi\delta}{3n}}^{x_j + \frac{2\pi\delta}{3n}} |K_n(x-x_k)| dx$$

which by virtue of (3.4) and (3.5) is

$$\geq \omega\left(\frac{\delta\pi}{3n}\right) |T(x_k)| \left( \frac{4}{\pi^2} \delta - \frac{3}{2} \delta^2 \sum_{\substack{j=0 \\ j \neq k}}^{3n-1} \frac{1}{(j-k-\frac{1}{2})^2} \right)$$

$$\geq \omega\left(\frac{\delta\pi}{3n}\right) |T(x_k)| \left( \frac{4}{\pi^2} \delta - \frac{3}{2} \delta^2 \sum_{j=0}^{\infty} \frac{1}{(j-\frac{1}{2})^2} \right)$$

Thus if we choose  $\delta_0 > 0$  such that

$$\left( \frac{4}{\pi^2} \delta_0 - \frac{3}{2} \delta_0^2 \sum_{j=0}^{\infty} \frac{1}{(j-\frac{1}{2})^2} \right) = C_0 > 0$$

We have, using the elementary properties of a modulus of continuity that

$$T(x_k) \int_0^{2\pi} \bar{g}_{\delta_0}(x) K_n(x-x_k) dx \geq C\omega\left(\frac{1}{n}\right) |T(x_k)| \quad k = 0, 1, 2, \dots, 3n-1$$

where  $C$  is an absolute positive constant. Finally,

$$\int_0^{2\pi} \bar{g}_{\delta_0}(x) T(x) dx = \frac{2}{3n} \sum_{k=0}^{3n-1} T(x_k) \int_0^{2\pi} \bar{g}_{\delta_0}(x) K_n(x-x_k) dx \geq$$

$$\geq C\omega\left(\frac{1}{3n}\right) \cdot \frac{2}{3n} \sum_{k=0}^{3n-1} |T(x_k)|$$

which by virtue of (3.3.) is

$$\geq \frac{2}{3} C\omega\left(\frac{1}{n}\right) \int_0^{2\pi} |T(x)| dx.$$

Thus, using Proposition 2,

$$\| \| T_n \| \|_\omega \geq \int_0^{2\pi} \bar{g}_{\delta_0}(x) T(x) dx \geq \frac{2}{3} C \omega \left( \frac{1}{n} \right) \int_0^{2\pi} |T(x)| dx$$

and the theorem is proved.

4. *Sufficient conditions for  $(\lambda_k)$  to be in  $(c_\omega, C_F)$ .* We first establish the result analogous to that of Bojanic (1.3) and (1.4). The proof is essentially that of Haršiladze [9].

THEOREM 4. 1. *If  $(\lambda_k)$  is a Stieljes sequence and if*

$$\omega \left( \frac{1}{n} \right) \int_0^{2\pi} |A_n(x)| dx = O(1) \quad (n \rightarrow \infty)$$

then  $(\lambda_k) \in (c_\omega, C_F)$ .

Proof: Let  $V_n(f)$  be the de la Vallée Poussin sums of  $f$

$$V_n(f) = \int_0^{2\pi} f(t) (2F_{2n}(t-x) - F_n(t-x)) dt.$$

It is well known [10, p. 92] that

$$(4.1) \quad \| f - V_n(f) \|_\infty \leq C \omega \left( f, \frac{1}{n} \right)$$

where  $C$  is a constant independent of  $f$  and  $n$ . Also if  $T$  is a trigonometric polynomial of degree  $n$  then

$$V_n(T) = T.$$

Thus if  $f \in C_\omega$ ,  $\| f \|_\omega \leq 1$

$$\begin{aligned} \int_0^{2\pi} f(t) A_n(t-x) dt &= \int_0^{2\pi} (f(t) - V_n(f)(t)) A_n(t-x) dt + \\ &+ \int_0^{2\pi} V_n(f)(t) A_n(t-x) dt. \end{aligned}$$

We have

$$\int_0^{2\pi} \left| \int_0^{2\pi} (2F_{2n}(t) - F_n(t)) A_n(t-x) dt \right| dx = O(1) \quad (n \rightarrow \infty).$$

Since  $(\lambda_k)$  is a Stieltjes sequence. Thus

$$\begin{aligned} \left\| \int_0^{2\pi} f(t) A_n(t-x) dt \right\|_{\infty} &\leq \left\| \int_0^{2\pi} (f(t) - V_n(f)(t)) A_n(t-x) dt \right\|_{\infty} + \\ &+ \|f\|_{\infty} \int_0^{2\pi} \left| \int_0^{2\pi} (2F_{2n}(t) - F_n(t)) A_n(t-x) dt \right| dx \leq \\ &\leq \left\| \int_0^{2\pi} (f(t) - V_n(f)(t)) A_n(t-x) dt \right\|_{\infty} + O(1) \quad (n \rightarrow \infty) \end{aligned}$$

which by virtue of (4.1) is

$$\leq C \omega\left(\frac{1}{n}\right) \int_0^{2\pi} |A_n(t)| dt + O(1) \quad (n \rightarrow \infty).$$

As a corollary of theorem 4.1 and theorem 3.1, we have

COROLLARY 4.1. *A Stieljes Sequence  $(\lambda_k)$  is in  $(c_{\omega}, C_F)$  if and only if*

$$\omega\left(\frac{1}{n}\right) \int_0^{2\pi} |A_n(t)| dt = O(1) \quad (n \rightarrow \infty).$$

We shall now give a sufficient condition for  $(\lambda_k)$  to be in  $(c_{\omega}, C_F)$  which requires no special restriction on  $(\lambda_k)$ .

THEOREM 4.2. *A sufficient condition for  $(\lambda_k)$  to be in  $(c_{\omega}, C_F)$  is that*

$$(4.2) \quad \omega(\mu_n) \int_0^{2\pi} |A_n(t)| dt = O(1) \quad (n \rightarrow \infty)$$

where

$$\mu_n = \frac{\int_0^{2\pi} \left| \int_0^x A_n(t) dt \right| dx}{\int_0^{2\pi} |A_n(t)| dt} \quad n = 0, 1, 2, \dots$$

If  $\omega(h) = h$  then (4.2) is also necessary.

Proof: We consider first the case when  $\omega(h) = h$ .

If  $f \in C_{\omega}$  with  $\|f\|_{\omega} \leq 1$  then

$$|f'(x)| \leq 1 \text{ a. e.}$$

So that

$$\left| \int_0^{2\pi} f(t) A_n(t-x) dt \right| = \left| \int_0^{2\pi} f'(t) \bar{A}_n(t-x) dt \right| \leq \int_0^{2\pi} |\bar{A}_n(t)| dt$$

$$\text{with } \bar{A}_n(t) = \int_0^t A_n(u) du.$$

Thus,

$$\| \| A_n \| \|_{\omega} \leq \int_0^{2\pi} |\bar{A}_n(t)| dt,$$

the function  $g(x) = \frac{1}{2\pi} \operatorname{sgn} \int_0^x A_n(t) dt$  is in  $C_{\omega}$  and  $\| \| g \| \|_{\omega} \leq 1$ . Also

$$\int_0^{2\pi} g(t) A_n(t) dt = |g(2\pi) A_n(2\pi) - \int_0^{2\pi} |\bar{A}_n(t)| dt| \geq \int_0^{2\pi} |\bar{A}_n(t)| dt - \lambda_0.$$

Thus,

$$\int_0^{2\pi} |\bar{A}_n(t)| dt - \lambda_0 \leq \| \| A_n \| \|_{\omega} \leq \int_0^{2\pi} |\bar{A}_n(t)| dt \quad n = 1, 2, \dots$$

This shows that (4.2) is necessary and sufficient for  $(\lambda_k)$  to be in  $(c_{\omega}, C_F)$  if  $\omega(h) = h$ .

Finally in the general case, the inequality

$$\| \int_0^{2\pi} f(t) A_n(t-x) dt \|_{\omega} \leq \omega(\mu_n) \int_0^{2\pi} |A_n(t)| dt$$

is a simple modification of Lemma 1 of [11] and we will not give its proof.

#### REFERENCES

- [1] KARAMATA, J., Suite de fonctionnelles linéaires et facteurs de convergence des séries de Fourier. *Journal de Math. P et Appl.*, 35 (1956), 87-95.
- [2] TOMIĆ, M., Sur les Facteurs de convergence des séries de Fourier des fonctions continues. *Publ. Inst. Math. Acad. Serb. Sci.*, VIII (1955), 23-32.
- [3] ——— Sur la sommation de la série de Fourier d'une fonction continue avec le module de continuité donné, *Publ. Inst. Math. Acad. Serb. Sci.*, X (1956), 19-36.
- [4] BOJANIC, R., On uniform convergence of Fourier series. *Publ. Inst. Math. Acad. Serb. Sci.*, X (1956), 153-158.

- [5] DUNFORD, N. and J. SCHWARTZ, *Linear Operators*, Vol. I. Interscience, N.Y., 1957, 858 pp.
- [6] HELSON, H., Proof of a conjecture of Steinhaus. *Proc. Nat. Acad. Sci. U.S.A.*, 40 (1954), 205-206.
- [7] ZYGMUND, A., *Trigonometric Series*, Vol. I, Cambridge Univ. Press, New York, 1959, 383 pp.
- [8] ——— *Trigonometric Series*, Vol. II, Cambridge Univ. Press, New York, 1959,
- [9] HARSILADZE, F., Multipliers of uniform convergence. *Trudi Tbilisk. Mat. Inst.*, 26 (1959), 121-130.
- [10] LORENTZ, G., *Approximation of Functions*. Holt, New York, 1966, 188 pp.
- [11] DEVORE, R., On Jackson's Theorem. *Jour. of App. Theory*, Acad. Press, 1 (1968), 314-318.

(Reçu le 15 novembre 1968)

Dep. of Mathematics  
Oakland University  
Rochester, Mi. 48063.