

# REPRESENTATIONS OF GROUP ALGEBRAS IN SPACES OF COMPLETELY BOUNDED MAPS

ROGER R. SMITH AND NICO SPRONK

ABSTRACT. Let  $G$  be a locally compact group,  $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$  be a strongly continuous unitary representation, and  $\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$  the space of normal completely bounded maps on  $\mathcal{B}(\mathcal{H})$ . We study the range of the map

$$\Gamma_\pi : M(G) \rightarrow \mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H})), \quad \Gamma_\pi(\mu) = \int_G \pi(s) \otimes \pi(s)^* d\mu(s)$$

where we identify  $\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$  with the extended Haagerup tensor product  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$ . We use the fact that the  $C^*$ -algebra generated by integrating  $\pi$  to  $L^1(G)$  is unital exactly when  $\pi$  is norm continuous, to show that  $\Gamma_\pi(L^1(G)) \subset \mathcal{B}(\mathcal{H}) \otimes^h \mathcal{B}(\mathcal{H})$  exactly when  $\pi$  is norm continuous. For the case that  $G$  is abelian, we study  $\Gamma_\pi(M(G))$  as a subset of the Varopoulos algebra. We also characterise positive definite elements of the Varopoulos algebra in terms of completely positive operators.

---

2000 *Mathematics Subject Classification*: Primary 46L07, 22D20; Secondary 22D10, 22D25, 22B05.

*Key words and phrases*: Group algebra, completely bounded map, extended Haagerup tensor product.

The first author was supported by a grant from the National Science Foundation.

The second author was supported by an NSERC Postdoctoral Fellowship.

## 1. INTRODUCTION

In [24], Størmer conducted an interesting study of spaces of completely bounded maps on  $\mathcal{B}(\mathcal{H})$ . For subalgebras  $\mathcal{A}$  and  $\mathcal{B}$  of  $\mathcal{B}(\mathcal{H})$  he defined what is now known as the Haagerup tensor product  $\mathcal{A} \otimes^h \mathcal{B}$ , as a completion of the set of elementary operators of the form  $x \mapsto \sum_{i=1}^n a_i x b_i$  where each  $a_i \in \mathcal{A}$  and each  $b_i \in \mathcal{B}$ . This approach gives the same tensor product norm as that in the more standard approach (see [8], for example), as shown in [21].

If  $G$  is an abelian group and  $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$  is a strongly continuous unitary representation, the homomorphism  $\Gamma_\pi$  from the measure algebra  $M(G)$  to the space  $\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$  of normal completely bounded maps on  $\mathcal{B}(\mathcal{H})$ , defined by

$$(1.1) \quad \Gamma_\pi(\mu) = \int_G \pi(s) \otimes \pi(s)^* d\mu(s)$$

was studied by Størmer. (We identify  $\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$  with the extended Haagerup tensor product  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$  from [4] and [9].) He used this homomorphism to generate many examples of regular and non-regular Banach subalgebras of  $\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$ . It was shown in [24, Lem. 5.6] that if  $\pi$  is norm continuous (i.e. continuous when the norm topology is placed on  $\mathcal{U}(\mathcal{H})$ ) then for any  $f$  in  $L^1(G)$

$$(1.2) \quad \Gamma_\pi(f) = \int_G f(s) \pi(s) \otimes \pi(s)^* ds \in C_\pi^* \otimes^h C_\pi^*$$

where  $C_\pi^*$  is the  $C^*$ -algebra generated by  $\{\int_G f(s) \pi(s) ds : f \in L^1(G)\}$ .

We note that for an arbitrary locally compact group  $G$ , the map  $\Gamma_\lambda$  as in (1.1), where  $\lambda$  is the left regular representation, was studied in [11] and [16].

In this paper we will make use of the theory of completely bounded normal maps on  $\mathcal{B}(\mathcal{H})$  from [21] to study the range of  $\Gamma_\pi$ . We show that, for a general locally compact group  $G$ ,

$$\Gamma_\pi(L^1(G)) \subset C_\pi^* \otimes^{eh} C_\pi^*$$

where  $\otimes^{eh}$  denotes the extended Haagerup tensor product from [9], [7] and [4]. Moreover, using the fact that  $C_\pi^*$  is unital exactly when the representation  $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$  is norm continuous, we show that the validity of (1.2) for every  $f$  in  $L^1(G)$  gives a characterisation of the norm continuity of  $\pi$ .

In the case that  $G$  is abelian, we develop the “Fourier-Stieltjes transform” for  $\Gamma_\pi(M(G))$ . The range of this transform is a Varopoulos type algebra  $V^b(E_\pi)$ , which will be defined below. We use some general results on completely positive maps to characterise complete positivity of elements of  $V^b(E_\pi)$ , as operators on  $\mathcal{B}(\mathcal{H})$ , extending some results from [24]. In particular, we characterise those  $\mu$  in  $M(G)$  for which  $\Gamma_\pi(\mu)$  is completely positive.

## 2. SPACES OF NORMAL COMPLETELY BOUNDED MAPS

Let  $\mathcal{H}$  be a Hilbert space, let  $\mathcal{B}(\mathcal{H})$  be the space of bounded operators on  $\mathcal{H}$  and let  $\mathcal{V}$  and  $\mathcal{W}$  be closed subspaces of  $\mathcal{B}(\mathcal{H})$ . The *Haagerup tensor product*  $\mathcal{V} \otimes^h \mathcal{W}$  is defined in [13] and [6]. The *extended Haagerup tensor product*  $\mathcal{V} \otimes^{eh} \mathcal{W}$  is developed in [9] and [7]; and also in [4], but in the context of dual spaces where it is called the “weak\* Haagerup tensor product” and denoted  $\mathcal{V} \otimes^{w^*h} \mathcal{W}$ . It is shown in [22] that the approach of [4] can be modified to develop the extended Haagerup tensor product in general.

Following [22], we thus define  $\mathcal{V} \otimes^{eh} \mathcal{W}$  to be the space of all (formal) series  $\sum_{i \in I} v_i \otimes w_i$  where each  $v_i \in \mathcal{V}$ , each  $w_i \in \mathcal{W}$ , and each of the series  $\sum_{i \in I} v_i v_i^*$  and  $\sum_{i \in I} w_i^* w_i$  converges weak\* in  $\mathcal{B}(\mathcal{H})$ . The index set  $I$  is established to have cardinality  $|I| = \dim \mathcal{H}$ . Two series  $\sum_{i \in I} v_i \otimes w_i$  and  $\sum_{i \in I} v'_i \otimes w'_i$  define the same element of  $\mathcal{V} \otimes^{eh} \mathcal{W}$  provided  $\sum_{i \in I} v_i x w_i = \sum_{i \in I} v'_i x w'_i$  for each  $x$  in  $\mathcal{B}(\mathcal{H})$ . Then  $\mathcal{V} \otimes^{eh} \mathcal{W}$  is a Banach space when endowed with the norm

$$\|T\|_{eh} = \inf \left\{ \left\| \sum_{i \in I} v_i v_i^* \right\|^{1/2} \left\| \sum_{i \in I} w_i^* w_i \right\|^{1/2} : T = \sum_{i \in I} v_i \otimes w_i \right\}$$

and the infimum is attained. As in [4], note that the Haagerup tensor product  $\mathcal{V} \otimes^h \mathcal{W}$  may be realized as the set of those  $T$  in  $\mathcal{V} \otimes^{eh} \mathcal{W}$  which admit a representation  $T = \sum_{i \in I} v_i \otimes w_i$  where  $\sum_{i \in I} v_i v_i^*$  and  $\sum_{i \in I} w_i^* w_i$  converge in norm. It is easy to see that any element  $T$  of  $\mathcal{V} \otimes^h \mathcal{W}$  may thus be written with a countable index set as  $T = \sum_{i=1}^{\infty} v_i \otimes w_i$ .

The space  $\mathcal{V} \otimes^{eh} \mathcal{W}$  has two natural, though more extrinsic descriptions. First, if  $\mathcal{V}$  and  $\mathcal{W}$  are each weak\* closed subspaces of  $\mathcal{B}(\mathcal{H})$ , they have respective preduals  $\mathcal{V}_*$  and  $\mathcal{W}_*$ . For example,

$$\mathcal{V}_* = \mathcal{B}(\mathcal{H})_* / \{\omega \in \mathcal{B}(\mathcal{H})_* : \omega(v) = 0 \text{ for all } v \text{ in } \mathcal{V}\}$$

which is an operator space when endowed with the quotient structure from the predual operator space structure on  $\mathcal{B}(\mathcal{H})_*$ . Then  $\mathcal{V} \otimes^{eh} \mathcal{W}$  is the dual

space of  $\mathcal{V}_* \otimes^h \mathcal{W}_*$  via the pairing

$$(2.1) \quad \left\langle \sum_{i \in I} v_i \otimes w_i, \omega \otimes \nu \right\rangle = \sum_{i \in I} \omega(v_i) \nu(w_i).$$

A proof of this can be found in [4] or [9]. In particular,  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H}) \cong (\mathcal{B}(\mathcal{H})_* \otimes^h \mathcal{B}(\mathcal{H})_*)^*$ .

Let  $\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$  denote the space of normal completely bounded operators on  $\mathcal{B}(\mathcal{H})$ . The map  $\theta : \mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$  given by

$$\theta \left( \sum_{i \in I} v_i \otimes w_i \right) x = \sum_{i \in I} v_i x w_i, \text{ for } x \text{ in } \mathcal{B}(\mathcal{H})$$

is a surjective isometry by [13] or [21]. Moreover,  $\theta$  is still an isometry when restricted to the spaces  $\mathcal{V} \otimes^{eh} \mathcal{W}$  or  $\mathcal{V} \otimes^h \mathcal{W}$ . For notational ease we will simply identify  $\mathcal{V} \otimes^{eh} \mathcal{W}$  and  $\mathcal{V} \otimes^h \mathcal{W}$  as subspaces of  $\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$  in the sequel, and omit the map  $\theta$ . In particular, we view  $\mathcal{B}(\mathcal{H}) \otimes^h \mathcal{B}(\mathcal{H})$  as being the completion in the completely bounded operator norm of the space of elementary operators  $x \mapsto \sum_{i=1}^n v_i x w_i$  on  $\mathcal{B}(\mathcal{H})$ . The composition of operators in  $\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$  induces a product in  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$ , making it a Banach algebra. This product is given on elementary tensors by

$$(a \otimes b) \circ (c \otimes d) = ac \otimes db.$$

The following is an extension of a theorem from [2], whose proof is much like the one offered there.

**Proposition 2.1.** *If  $\mathcal{A}$  and  $\mathcal{B}$  are norm closed subalgebras of  $\mathcal{B}(\mathcal{H})$ , then  $\mathcal{A} \otimes^{eh} \mathcal{B}$  is a subalgebra of  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$ . If  $\mathcal{V}$  is a (left)  $\mathcal{A}$ -module and  $\mathcal{W}$  is a (right)  $\mathcal{B}$ -module in  $\mathcal{B}(\mathcal{H})$ , then  $\mathcal{V} \otimes^{eh} \mathcal{W}$  is a (left)  $\mathcal{A} \otimes^{eh} \mathcal{B}$ -module in  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$ .*

If  $\Omega \in \mathcal{B}(\mathcal{H})^*$  then the *left* and *right slice maps*  $L_\Omega, R_\Omega : \mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  are given for  $T = \sum_{i \in I} v_i \otimes w_i$  by

$$(2.2) \quad L_\Omega T = \sum_{i \in I} \Omega(v_i) w_i \quad \text{and} \quad R_\Omega T = \sum_{i \in I} \Omega(w_i) v_i.$$

These series each converge in norm as is shown in [22, Thm. 2.2]. Moreover, it is shown there that for any pair of closed subspaces  $\mathcal{V}$  and  $\mathcal{W}$  of  $\mathcal{B}(\mathcal{H})$ ,  $\mathcal{V} \otimes^{eh} \mathcal{W}$  consists exactly of those  $T$  in  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$  for which  $L_\Omega T \in \mathcal{W}$  and  $R_\Omega T \in \mathcal{V}$  for each  $\Omega$  in  $\mathcal{B}(\mathcal{H})^*$  (or for which  $L_\omega T \in \mathcal{W}$  and  $R_\omega T \in \mathcal{V}$  for each  $\omega$  in  $\mathcal{B}(\mathcal{H})_*$ ). These results extend [21, Thm. 4.5].

We will finish this section with a theorem on completely positive maps which will be useful in Section 4. We will first need some general preliminary results which are modeled on results from [21].

A closed subalgebra  $\mathcal{B}$  of  $\mathcal{B}(\mathcal{H})$  is called *locally cyclic* if for each finite dimensional subspace  $\mathcal{L}$  of  $\mathcal{H}$ , there is a vector  $\xi$  in  $\mathcal{H}$  such that  $\overline{\mathcal{B}\xi} \supset \mathcal{L}$ . We note, for example, that if  $\mathcal{B}$  is a maximal abelian self-adjoint subalgebra of  $\mathcal{B}(\mathcal{H})$  then it is locally cyclic. Indeed if  $\xi_1, \dots, \xi_n$  span  $\mathcal{L}$ , consider the orthogonal projections  $p_1, p_2, \dots, p_n$  whose respective ranges are

$$\overline{\mathcal{B}\xi_1}, \overline{\mathcal{B}\xi_2} \ominus \overline{\mathcal{B}\xi_1}, \dots, \overline{\mathcal{B}\xi_n} \ominus \bigoplus_{i=1}^{n-1} \overline{\mathcal{B}\xi_i}.$$

Then each  $p_i \in \mathcal{B}' = \mathcal{B}$ , and  $\xi = \xi_1 + p_2\xi_2 + \dots + p_n\xi_n$  satisfies  $\overline{\mathcal{B}\xi} \supset \mathcal{L}$ .

The following is an adaptation of [21, Thm. 2.1].

**Lemma 2.2.** *If  $\mathcal{B}$  is a locally cyclic  $C^*$ -subalgebra of  $\mathcal{B}(\mathcal{H})$  and  $T : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  is a positive map which is also a  $\mathcal{B}$ -bimodule map, then  $T$  is completely positive.*

**Proof.** Let us fix  $n$ , a positive matrix  $[x_{ij}]$  in  $M_n(\mathcal{B}(\mathcal{H}))$  and a column vector  $\xi = [\xi_1 \cdots \xi_n]^t$  in  $\mathcal{H}^n$  with  $\|\xi\| < 1$ . Then, given  $\varepsilon > 0$ , there is vector  $\xi$  in  $\mathcal{H}$  and elements  $b_1, \dots, b_n$  in  $\mathcal{B}$  such that the vector  $\eta = [b_1\xi \cdots b_n\xi]^t$  satisfies  $\|\xi - \eta\| < \varepsilon$  and  $\|\eta\| < 1$ . Letting  $T^{(n)} : M_n(\mathcal{B}(\mathcal{H})) \rightarrow M_n(\mathcal{B}(\mathcal{H}))$  be the

amplification of  $T$ , we have

$$\begin{aligned} \langle T^{(n)}[x_{ij}]\boldsymbol{\eta}|\boldsymbol{\eta} \rangle &= \left\langle [Tx_{ij}] \begin{bmatrix} b_1\xi \\ \vdots \\ b_n\xi \end{bmatrix} \middle| \begin{bmatrix} b_1\xi \\ \vdots \\ b_n\xi \end{bmatrix} \right\rangle \\ &= \sum_{i,j=1}^n \langle b_i^* T(x_{ij}) b_j \xi | \xi \rangle = \left\langle T \left( \sum_{i,j=1}^n b_i^* x_{ij} b_j \right) \xi | \xi \right\rangle \geq 0 \end{aligned}$$

and

$$\left| \langle T^{(n)}[x_{ij}]\boldsymbol{\eta}|\boldsymbol{\eta} \rangle - \langle T^{(n)}[x_{ij}]\boldsymbol{\xi}|\boldsymbol{\xi} \rangle \right| < \left( \|T^{(n)}\| + 1 \right) \varepsilon.$$

Since  $\varepsilon$  can be chosen arbitrarily small, we conclude that  $\langle T^{(n)}[x_{ij}]\boldsymbol{\xi}|\boldsymbol{\xi} \rangle \geq 0$ .

Hence  $T$  is completely positive.  $\square$

If a family of operators  $\{b_i\}_{i \in I}$  from  $\mathcal{B}(\mathcal{H})$  defines a bounded row matrix  $B = [\cdots b_i \cdots]$ , i.e.  $\sum_{i \in I} b_i b_i^*$  converges weak\* in  $\mathcal{B}(\mathcal{H})$ , then the product  $B \cdot \boldsymbol{\lambda} = \sum_{i \in I} \lambda_i b_i$  converges in norm and thus defines an element of  $\mathcal{B}(\mathcal{H})$  for each  $\boldsymbol{\lambda} = [\cdots \lambda_i \cdots]^t$  in  $\ell^2(I)$ . We say that the set  $\{b_i\}_{i \in I}$  is *strongly independent* if  $B \cdot \boldsymbol{\lambda} = 0$  only when  $\boldsymbol{\lambda} = 0$ . This is an obvious extension of the usual notion of linear independence, and can be easily adapted to column matrices. Elements of  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$  admit many different representations, and strong independence was introduced in [21] to handle the difficulties caused by this.

The following is an adaptation of [21, Thm. 3.1].

**Lemma 2.3.** *If  $\mathcal{A}$  is a  $C^*$ -subalgebra of  $\mathcal{B}(\mathcal{H})$  and  $T \in \mathcal{A} \otimes^{eh} \mathcal{A}$ , then  $T$  is completely positive if and only if there is a strongly independent family  $\{a_i\}_{i \in I}$  from  $\mathcal{A}$  for which  $\sum_{i \in I} a_i a_i^*$  converges weak\* in  $\mathcal{B}(\mathcal{H})$  and  $T = \sum_{i \in I} a_i \otimes a_i^*$ .*

**Proof.** We need only to prove that the first condition implies the second.

If  $T$  is completely positive and normal on  $\mathcal{B}(\mathcal{H})$ , then its restriction to the algebra of compact operators  $T|_{\mathcal{K}(\mathcal{H})}$  is a completely positive map which determines  $T$ . Using Stinespring's theorem and the representation theory

for  $\mathcal{K}(\mathcal{H})$ , just as in [21, Thm. 3.1] or [13], we obtain a family  $\{b_j\}_{j \in J}$  from  $\mathcal{B}(\mathcal{H})$  for which  $\sum_{j \in J} b_j b_j^*$  converges weak\* in  $\mathcal{B}(\mathcal{H})$  and  $T = \sum_{j \in J} b_j \otimes b_j^*$ . We see that  $J$  can be any index set whose cardinality coincides with the Hilbertian dimension of  $\mathcal{H}$ . Let  $B = [\cdots b_j \cdots]$ .

Now we let

$$\mathcal{L} = \{\boldsymbol{\lambda} \in \ell^2(J) : B \cdot \boldsymbol{\lambda} = 0\}$$

and partition  $J = I' \cup I$  in such a way that there is an orthonormal basis  $\{\boldsymbol{\lambda}_j\}_{j \in J}$  of  $\ell^2(J)$  for which

$$\overline{\text{span}}\{\boldsymbol{\lambda}_i\}_{i \in I'} = \mathcal{L} \quad \text{and} \quad \overline{\text{span}}\{\boldsymbol{\lambda}_i\}_{i \in I} = \mathcal{L}^\perp.$$

Let  $U$  denote the  $J \times J$  unitary matrix whose columns are the vectors  $\{\boldsymbol{\lambda}_j\}_{j \in J}$ . Let  $A = [\cdots a_j \cdots] = B \cdot U$ . Note that  $a_j = 0$  for each  $j$  in  $I'$ . Then for any  $x$  in  $\mathcal{B}(\mathcal{H})$ , letting  $x^J$  denote the  $J \times J$  diagonal matrix which is the amplification of  $x$ , we have

$$Tx = \sum_{j \in J} b_j x b_j^* = B x^J B^* = B \cdot U x^J U^* \cdot B^* = A x^J A^* = \sum_{i \in I} a_i x a_i^*.$$

We have that  $\{a_i\}_{i \in I}$  is strongly independent, for if  $\boldsymbol{\alpha} = [\cdots \alpha_i \cdots]^t$  in  $\ell^2(I)$  is such that  $A \cdot \boldsymbol{\alpha} = 0$ , then

$$0 = A \cdot \boldsymbol{\alpha} = \sum_{i \in I} \alpha_i a_i = \sum_{i \in I} \alpha_i B \cdot \boldsymbol{\lambda}_i = B \cdot \left( \sum_{i \in I} \alpha_i \boldsymbol{\lambda}_i \right)$$

so  $\sum_{i \in I} \alpha_i \boldsymbol{\lambda}_i \in \mathcal{L} \cap \mathcal{L}^\perp$ , whence  $\boldsymbol{\alpha} = 0$ . Hence

$$T = \sum_{i \in I} a_i \otimes a_i^*$$

where  $\{a_i\}_{i \in I}$  is strongly independent. It remains to show that  $\{a_i\}_{i \in I} \subset \mathcal{A}$ .

Since  $\{a_i\}_{i \in I}$  is strongly independent, so too is  $\{a_i^*\}_{i \in I}$ . Hence by [1, Lem. 2.2], the space

$$\{[\cdots \Omega(a_i^*) \cdots]^t : \Omega \in \mathcal{B}(\mathcal{H})^*\}$$

is dense in  $\ell^2(I)$ . Thus, given a fixed index  $i_0$  in  $I$ , there is a (not necessarily bounded) sequence  $\{\Omega_n\}_{n=1}^\infty$  from  $\mathcal{B}(\mathcal{H})^*$  such that

$$a_{i_0} = \lim_{n \rightarrow \infty} \sum_{i \in I} \Omega_n(a_i^*) a_i = \lim_{n \rightarrow \infty} R_{\Omega_n} T.$$

Since  $R_\Omega T \in \mathcal{A}$  for each right slice map  $R_\Omega$ , it follows that  $a_{i_0} \in \mathcal{A}$ .  $\square$

If  $E$  is any locally compact space we let

$$\begin{aligned} V_0(E) &= \mathcal{C}_0(E) \otimes^h \mathcal{C}_0(E) \\ (2.3) \quad V^0(E) &= \mathcal{C}_0(E) \otimes^{eh} \mathcal{C}_0(E) \\ \text{and } V^b(E) &= \mathcal{C}_b(E) \otimes^{eh} \mathcal{C}_b(E). \end{aligned}$$

These spaces are discussed in [22]. These all may be regarded as Banach algebras of functions on  $E \times E$  by Proposition 2.1. However, as pointed out in [20], an element  $u$  of  $V^b(E)$  may not be continuous on  $E \times E$ , even if  $E$  is compact. However, if  $\mathcal{C}$  is a closed subalgebra of  $\mathcal{C}_b(E)$  (say  $\mathcal{C} = \mathcal{C}_0(E)$ ), then for each  $u \in \mathcal{C} \otimes^{eh} \mathcal{C} \subset V^b(E)$ , the pointwise slices,  $u(\cdot, x)$  and  $u(x, \cdot)$  for any fixed  $x$  in  $E$ , will always be elements of  $\mathcal{C}$ . In the case where  $E$  is a compact group,  $V_0(E)$  is discussed in [23], and in a profound way in [25]. We note that by Grothendieck's Inequality,  $V_0(E) = \mathcal{C}_0(E) \otimes^\gamma \mathcal{C}_0(E)$  (projective tensor product), up to equivalent norms.

If  $u : E \times E \rightarrow \mathbb{C}$ , we say that  $u$  is *positive definite* if for any finite collection of elements  $x_1, \dots, x_n$  from  $E$ , the matrix  $[u(x_i, x_j)]$  is of positive type.

If  $\mathcal{A}$  is any abelian  $C^*$ -algebra for which there is a locally compact space  $E$  and an injective  $*$ -homomorphism  $F : \mathcal{A} \rightarrow \mathcal{C}_b(E)$ , then there is an isometric algebra homomorphism  $F \otimes F : \mathcal{A} \otimes^{eh} \mathcal{A} \rightarrow V^b(E)$ , by [9] or [22, Cor. 2.3].

The following theorem generalises [24, Thm. 5.1].

**Theorem 2.4.** *Let  $\mathcal{A}$  be an abelian  $C^*$ -subalgebra of  $\mathcal{B}(\mathcal{H})$  for which there is a locally compact space  $E$  and an injective  $*$ -homomorphism  $F : \mathcal{A} \rightarrow \mathcal{C}_b(E)$ . If  $T \in \mathcal{A} \otimes^{eh} \mathcal{A}$  and  $u = (F \otimes F)T$ , so  $u \in F(\mathcal{A}) \otimes^{eh} F(\mathcal{A}) \subset V^b(E)$ , then the following are equivalent:*

- (i)  $T$  is positive.
- (ii)  $T$  is completely positive.
- (iii)  $u$  is positive definite.

**Proof.** (i) $\Rightarrow$ (ii) If  $\mathcal{B}$  is any maximal abelian subalgebra of  $\mathcal{B}(\mathcal{H})$  which contains  $\mathcal{A}$ , then  $T$  is a  $\mathcal{B}$ -bimodule map. The result then follows from Lemma 2.2.

(ii) $\Rightarrow$ (iii) By Lemma 2.3 we have that  $T = \sum_{i \in I} a_i \otimes a_i^*$  for some family of elements from  $\mathcal{A}$  for which  $\sum_{i \in I} a_i a_i^*$  converges weak\* in  $\mathcal{B}(\mathcal{H})$ . Let  $\varphi_i = F(a_i)$  in  $\mathcal{C}_b(E)$ , so

$$u = \sum_{i \in I} \varphi_i \otimes \bar{\varphi}_i \quad \text{and} \quad \left\| \sum_{i \in I} |\varphi_i|^2 \right\|_{\infty} < \infty.$$

Let  $\xi : E \rightarrow \ell^2(I)$  be given by  $\xi(x) = (\varphi_i(x))_{i \in I}$ . Then for each  $(x, y)$  in  $E \times E$ , we have that

$$(2.4) \quad u(x, y) = \langle \xi(x) | \xi(y) \rangle$$

and hence  $u$  is positive definite.

(iii) $\Rightarrow$ (i) Since  $u$  is positive definite function, then by [12, §3.1], there is a Hilbert space  $\mathcal{L}$  and a bounded function  $\xi : E \rightarrow \mathcal{L}$  such that (2.4) holds. Let  $p$  be the orthogonal projection on  $\mathcal{L}$  whose range is  $\overline{\text{span}}\{\xi(x)\}_{x \in E}$ , and let  $\{\xi_i\}_{i \in I}$  be an orthonormal basis for  $p\mathcal{L}$ . Then for each  $i$  the function

$$\varphi_i = \langle \xi(\cdot) | \xi_i \rangle$$

is in  $F(\mathcal{A})$ . Indeed, given  $\varepsilon > 0$  we can find  $\alpha_1, \dots, \alpha_n$  from  $\mathbb{C}$  and  $y_1, \dots, y_n$  from  $E$ , such that

$$\left\| \xi_i - \sum_{k=1}^n \alpha_k \xi(y_k) \right\| < \varepsilon$$

whence

$$\left\| \varphi_i - \sum_{k=1}^n \bar{\alpha}_k u(\cdot, y_k) \right\|_{\infty} = \left\| \langle \xi(\cdot) | \xi_i \rangle - \sum_{k=1}^n \bar{\alpha}_k \langle \xi(\cdot) | \xi(y_k) \rangle \right\|_{\infty} < \|\xi\|_{\infty} \varepsilon.$$

Hence  $\varphi_i$  can be uniformly approximated arbitrarily closely by elements of  $F(\mathcal{A})$ , and our conclusion holds. It then follows by Parseval's Identity that for any  $(x, y)$  in  $E \times E$

$$u(x, y) = \langle p\xi(x) | p\xi(y) \rangle = \sum_{i \in I} \langle \xi(x) | \xi_i \rangle \langle \xi_i | \xi(y) \rangle = \sum_{i \in I} \varphi_i(x) \overline{\varphi_i(y)}.$$

Hence we may write

$$u = \sum_{i \in I} \varphi_i \otimes \bar{\varphi}_i \quad \text{with} \quad \left\| \sum_{i \in I} |\varphi_i|^2 \right\|_{\infty} = \|\xi\|_{\infty}^2 < \infty.$$

Letting  $a_i = F^{-1}(\varphi_i)$  in  $\mathcal{A}$ , we get that  $T = (F \otimes F)^{-1}u = \sum_{i \in I} a_i \otimes a_i^*$  and is thus positive.  $\square$

## 3. REPRESENTATIONS OF GROUPS IN COMPLETELY BOUNDED MAPS

Let  $G$  be a locally compact group, let  $\mathcal{A}$  be a unital Banach algebra which is also a dual space with predual  $\mathcal{A}_*$ , and let  $\alpha : G \rightarrow \mathcal{A}_{\text{inv}}$  be a weak\* continuous bounded homomorphism where  $\mathcal{A}_{\text{inv}}$  denotes the group of invertible elements in  $\mathcal{A}$ . In particular we assume  $\alpha(e)$  is the unit of  $\mathcal{A}$ . Denote the space of bounded complex Borel measures on  $G$  by  $M(G)$ . Recall that  $M(G)$  is the dual space to the space  $\mathcal{C}_0(G)$  of continuous functions vanishing at infinity. Recall too that  $M(G)$  is a Banach algebra via *convolution*: for each  $\mu, \nu$  in  $M(G)$  we define  $\mu * \nu$  by

$$(3.1) \quad \int_G \varphi d\mu * \nu = \int_G \int_G \varphi(st) d\mu(s) d\nu(t)$$

for each  $\varphi$  in  $\mathcal{C}_0(G)$ . We note that since each of  $\mu$  and  $\nu$  can be approximated in norm by compactly supported bounded measures, (3.1) holds for any choice of  $\varphi$  in  $\mathcal{C}_b(G)$  too. If  $\mu \in M(G)$ , let

$$\alpha_1(\mu) = \text{weak}^* \text{-} \int_G \alpha(s) d\mu(s)$$

i.e. if  $\omega \in \mathcal{A}_*$ , then  $\langle \alpha_1(\mu), \omega \rangle = \int_G \langle \alpha(s), \omega \rangle d\mu(s)$ . Then  $\alpha_1 : M(G) \rightarrow \mathcal{A}$  is a bounded linear map for if  $\|\alpha\|_\infty = \sup_{s \in G} \|\alpha(s)\|$ , then

$$(3.2) \quad \|\alpha_1(\mu)\| = \sup_{\omega \in \text{b}_1(\mathcal{A}_*)} \left| \int_G \langle \alpha(s), \omega \rangle d\mu(s) \right| \leq \int_G \|\alpha\|_\infty d|\mu|(s) = \|\alpha\|_\infty \|\mu\|_1.$$

Recall that the dual  $\mathcal{A}^*$  is a contractive  $\mathcal{A}$ -bimodule where for  $b$  in  $\mathcal{A}$  and  $F$  in  $\mathcal{A}^*$  we define  $b \cdot F$  and  $F \cdot b$  in  $\mathcal{A}^*$  by  $\langle a, b \cdot F \rangle = \langle ab, F \rangle$  and  $\langle a, F \cdot b \rangle = \langle ba, F \rangle$ , for each  $a$  in  $\mathcal{A}$ . We say that a subspace  $\Omega$  of  $\mathcal{A}^*$  is a right  $\alpha(G)$ -submodule if  $\omega \cdot \alpha(s) \in \Omega$ , for each  $\omega$  in  $\Omega$  and  $s$  in  $G$ .

**Proposition 3.1.** *Let  $G$ ,  $\mathcal{A}$  and  $\alpha$  be as above. Moreover, suppose that  $\mathcal{A}_*$  is both a left  $\mathcal{A}$ -submodule of  $\mathcal{A}^*$  and a right  $\alpha(G)$ -submodule. Then  $\alpha_1 : M(G) \rightarrow \mathcal{A}$  is a unital algebra homomorphism.*

**Proof.** If  $\mu, \nu \in M(G)$  and  $\omega \in \mathcal{A}^*$  then

$$\begin{aligned}
 \langle \alpha_1(\mu)\alpha_1(\nu), \omega \rangle &= \langle \alpha_1(\mu), \alpha_1(\nu) \cdot \omega \rangle \\
 &= \int_G \langle \alpha(s), \alpha_1(\nu) \cdot \omega \rangle d\mu(s) \\
 &= \int_G \langle \alpha_1(\nu), \omega \cdot \alpha(s) \rangle d\mu(s) \\
 &= \int_G \int_G \langle \alpha(t), \omega \cdot \alpha(s) \rangle d\nu(t) d\mu(s) \\
 &= \int_G \int_G \langle \alpha(st), \omega \rangle d\nu(t) d\mu(s).
 \end{aligned}$$

where the hypotheses guarantee that  $\alpha_1(\nu) \cdot \omega \in \mathcal{A}_*$  and that  $\omega \cdot \alpha(s) \in \mathcal{A}_*$ , for each  $s$ . By Fubini's Theorem we have that

$$\int_G \int_G \langle \alpha(st), \omega \rangle d\nu(t) d\mu(s) = \int_G \int_G \langle \alpha(st), \omega \rangle d\mu(s) d\nu(t) = \langle \alpha_1(\mu * \nu), \omega \rangle$$

where we note that  $(s, t) \mapsto \langle \alpha(st), \omega \rangle$  is continuous and bounded, hence  $\mu \times \nu$ -integrable.

That  $\alpha_1(\delta_e) = \alpha(e)$  follows from that  $\mathcal{A}_*$  is a separating for  $\mathcal{A}$ . Hence  $\alpha_1$  is a unital map.  $\square$

By a symmetric argument, the above proposition also holds if  $\mathcal{A}_*$  is assumed to be both a right  $\mathcal{A}$ -submodule of  $\mathcal{A}^*$  and a left  $\alpha(G)$ -submodule.

**Example 3.2.** (i) Let  $\mathcal{X}$  be a Banach space admitting a predual  $\mathcal{X}_*$ . Then we have that  $\mathcal{A} = \mathcal{B}(\mathcal{X})$  is a dual unital Banach algebra admitting a predual  $\mathcal{A}_* = \mathcal{X} \otimes^\gamma \mathcal{X}_*$ , via the dual pairing

$$\langle T, x \otimes \omega \rangle = \langle Tx, \omega \rangle \text{ for } T \text{ in } \mathcal{A}, x \text{ in } \mathcal{X} \text{ and } \omega \text{ in } \mathcal{X}_*.$$

Here  $\otimes^\gamma$  denotes the *projective tensor product*. We have then that  $\mathcal{A}_*$  is a left  $\mathcal{A}$  submodule of  $\mathcal{A}^*$ . Indeed, for any  $S, T$  in  $\mathcal{A}$  and elementary tensor  $x \otimes \omega$  in  $\mathcal{A}_*$  we have that,

$$\langle ST, x \otimes \omega \rangle = \langle STx, \omega \rangle = \langle S, (Tx) \otimes \omega \rangle$$

so  $T \cdot (x \otimes \omega) = (Tx) \otimes \omega$ .

If  $\mathcal{B}^\sigma(\mathcal{X})$  denotes the weak\*-weak\* continuous bounded linear maps on  $\mathcal{X}$  then  $\mathcal{A}_*$  is a right  $\mathcal{B}^\sigma(\mathcal{X})$ -submodule of  $\mathcal{A}^*$ . Thus we obtain the situation of Proposition 3.1 whenever  $\alpha : G \rightarrow \mathcal{A}_{\text{inv}}$  is a weak\* continuous bounded homomorphism whose range is in  $\mathcal{B}^\sigma(\mathcal{X})$ . In particular, this happens when  $\mathcal{X}$  is reflexive and  $\alpha$  is a non-degenerate strong operator continuous representation on  $\mathcal{X}$ .

(ii) The example above can be easily modified for the case where  $\mathcal{V}$  is a dual operator space and  $\mathcal{A} = \mathcal{CB}(\mathcal{V})$ .

(iii) There is a standard identification  $\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H})) \cong \mathcal{CB}(\mathcal{K}(\mathcal{H}), \mathcal{B}(\mathcal{H}))$ , and thus an identification of  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H}) \cong \mathcal{CB}(\mathcal{K}(\mathcal{H}), \mathcal{B}(\mathcal{H}))$ . In fact, as shown in [4], this latter identification is a weak\* homeomorphism. Indeed, using standard identifications with row and column Hilbert spaces and the *operator projective tensor product*,  $\hat{\otimes}$  (see [3] or [8, II.9.3]), we have

$$\begin{aligned} \mathcal{B}(\mathcal{H})_* \otimes^h \mathcal{B}(\mathcal{H})_* &\cong \left( \overline{\mathcal{H}}_r \otimes^h \mathcal{H}_c \right) \otimes^h \left( \overline{\mathcal{H}}_r \otimes^h \mathcal{H}_c \right) \\ &\cong \overline{\mathcal{H}}_r \otimes^h \left( \mathcal{H}_c \otimes^h \overline{\mathcal{H}}_r \right) \otimes^h \mathcal{H}_c \cong \overline{\mathcal{H}}_r \hat{\otimes} \mathcal{K}(\mathcal{H}) \hat{\otimes} \mathcal{H}_c \\ &\cong \mathcal{K}(\mathcal{H}) \hat{\otimes} \overline{\mathcal{H}}_r \hat{\otimes} \mathcal{H}_c \cong \mathcal{K}(\mathcal{H}) \hat{\otimes} \mathcal{B}(\mathcal{H})_*. \end{aligned}$$

On elementary tensors this identification is given by

$$(\xi^* \otimes \eta) \otimes (\zeta^* \otimes \vartheta) \mapsto (\eta \otimes \zeta^*) \otimes (\xi^* \otimes \vartheta)$$

where for vectors  $\xi, \eta$  in  $\mathcal{H}$  we let  $\xi \otimes \eta^*$  denote the usual rank 1 operator and  $\xi^* \otimes \eta$  the usual vector functional. Now if  $T = \sum_{i \in I} a_i \otimes b_i$  in  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$  then, in the dual pairing (2.1), we have that

$$\langle T, (\xi^* \otimes \eta) \otimes (\zeta^* \otimes \vartheta) \rangle = \sum_{i \in I} \langle a_i \eta | \xi \rangle \langle b_i \vartheta | \zeta \rangle.$$

Meanwhile, in the  $\mathcal{CB}(\mathcal{K}(\mathcal{H}), \mathcal{B}(\mathcal{H}))$ - $\mathcal{K}(\mathcal{H}) \hat{\otimes} \mathcal{B}(\mathcal{H})_*$  duality we have that

$$\begin{aligned} \langle T, (\eta \otimes \zeta^*) \otimes (\xi^* \otimes \vartheta) \rangle &= \left\langle \sum_{i \in I} a_i \eta \otimes (b_i^* \zeta)^*, \xi^* \otimes \vartheta \right\rangle \\ &= \sum_{i \in I} \langle a_i \eta | \xi \rangle \langle b_i \vartheta | \zeta \rangle. \end{aligned}$$

Now for every elementary tensor  $k \otimes \omega$  in  $\mathcal{K}(\mathcal{H}) \hat{\otimes} \mathcal{B}(\mathcal{H})_*$  and  $T$  in  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$ , we have that  $(k \otimes \omega) \cdot T = k \otimes (\omega \cdot T) \in \mathcal{K}(\mathcal{H}) \hat{\otimes} \mathcal{B}(\mathcal{H})_*$ . Hence  $\mathcal{B}(\mathcal{H})_* \otimes^h \mathcal{B}(\mathcal{H})_* \cong \mathcal{K}(\mathcal{H}) \hat{\otimes} \mathcal{B}(\mathcal{H})_*$  is a right module for  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$ . We note that  $\mathcal{B}(\mathcal{H})_* \otimes^h \mathcal{B}(\mathcal{H})_*$  is a left  $\mathcal{B}(\mathcal{H}) \otimes^h \mathcal{B}(\mathcal{H})$ -module.  $\square$

We will identify the group algebra  $L^1(G)$  with the closed ideal in  $M(G)$  of measures which are absolutely continuous with respect to the left Haar measure  $m$  (whose integral we will denote  $\int_G \cdots ds$ ). We will identify the discrete group algebra  $\ell^1(G)$  with the closed subspace of  $M(G)$  generated by all of the Dirac measures  $\{\delta_s : s \in G\}$ . We let

$$M_\alpha = \overline{\alpha_1(M(G))}, \quad C_\alpha = \overline{\alpha_1(L^1(G))} \quad \text{and} \quad D_\alpha = \overline{\alpha_1(\ell^1(G))}$$

where each of the closures is in the norm topology of  $\mathcal{A}$ .

The following proposition is surely well-known, though we have been unable to find it in the literature.

**Proposition 3.3.** *Given  $G$ ,  $\mathcal{A}$  and  $\alpha$  satisfying the hypotheses of Proposition 3.1,  $\alpha$  is norm continuous if and only if  $C_\alpha$  is unital.*

**Proof.** Let  $(e_U)$  be the bounded approximate identity for  $L^1(G)$  given by  $e_U = \frac{1}{m(U)} 1_U$  (normalised indicator function), indexed over the family of all relatively compact neighbourhoods of the identity  $e$  in  $G$ , partially ordered by reverse inclusion.

“ $\Rightarrow$ ” Let  $\varepsilon > 0$ . Let  $V$  be any relatively compact neighbourhood of  $e$  for which  $\|\alpha(s) - \alpha(e)\| < \varepsilon$  for each  $s$  in  $V$ . Then for any relatively compact neighbourhood  $U$  of  $e$  which is contained in  $V$  we have

$$\begin{aligned} \|\alpha_1(e_U) - \alpha(e)\| &= \left\| \frac{1}{m(U)} \int_U \alpha(s) ds - \alpha(e) \right\| \\ &\leq \frac{1}{m(U)} \int_U \|\alpha(s) - \alpha(e)\| ds < \varepsilon \end{aligned}$$

where the second from last inequality is proved just as in (3.2). Thus  $\alpha(e) = \lim_U \alpha_1(e_U)$  in norm, so  $\alpha(e) \in C_\alpha$ . Now  $\alpha(e)$  is the unit for  $\mathcal{A}$ , and hence the unit for  $C_\alpha$ .

“ $\Leftarrow$ ” It is a standard fact that  $\lim_U \alpha_1(e_U) = \alpha(e)$  in the weak\* topology of  $\mathcal{A}$ . Indeed,  $\lim_U \int_G e_U(s)\varphi(s)ds = \varphi(e)$  for any continuous function  $\varphi$ ; set  $\varphi = \langle \alpha(\cdot), \omega \rangle$  for any  $\omega$  in  $\mathcal{A}_*$ . Now let  $E$  be the unit for  $C_\alpha$ . We will establish that  $E = \alpha(e)$ , the unit of  $\mathcal{A}$ . First, the map  $s \mapsto \alpha(s)E$  is norm continuous. Indeed  $E \in C_\alpha$  and can thus be norm approximated by  $\{\alpha_1(f) : f \in L^1(G)\}$ . Moreover, if  $f \in L^1(G)$  then we have that

$$\|\alpha(s)\alpha_1(f) - \alpha_1(f)\| = \|\alpha_1(\delta_s * f - f)\| \leq \|\alpha\|_\infty \|\delta_s * f - f\|_1 \xrightarrow{s \rightarrow e} 0$$

where the inequality follows from (3.2) and limit follows from [15, 20.4].

Next, for any compact neighbourhood  $U$  of  $e$  we have that

$$\alpha_1(e_U) = \alpha_1(e_U)E = \frac{1}{m(U)} \int_U \alpha(s)ds \cdot E = \frac{1}{m(U)} \int_U \alpha(s)E ds$$

where we note that right multiplication is weak\*-continuous in  $\mathcal{A}$ , by hypothesis. Now, let  $\varepsilon > 0$  be given, and find a neighbourhood  $V$  of  $e$  in  $G$  such that  $\|\alpha(s)E - E\| < \varepsilon$  for each  $s$  in  $V$ . Then for any relatively compact neighbourhood  $U$  of  $E$ , contained in  $V$ , we have that

$$\|\alpha_1(e_U) - E\| = \left\| \frac{1}{m(U)} \int_U \alpha(s)E ds - E \right\| \leq \frac{1}{m(U)} \int_U \|\alpha(s)E - E\| ds < \varepsilon$$

where the second from last inequality is proved just as in (3.2). Hence we have that  $\lim_U \alpha_1(e_U) = E$  in norm, so, *a fortiori*, weak\*- $\lim_U \alpha_1(e_U) = E$ . It then follows from above that  $E = \alpha(e)$ , so  $\alpha(e) \in C_\alpha$ . Thus

$$\|\alpha(s) - \alpha(e)\| = \|\alpha(s)E - E\| \xrightarrow{s \rightarrow e} 0.$$

Hence  $\alpha$  is norm continuous at  $e$ , and thus norm continuous on all of  $G$ .  $\square$

**Corollary 3.4.** *For  $G$ ,  $\mathcal{A}$  and  $\alpha$  as above, the following are equivalent:*

- (i)  $\alpha$  is norm continuous      (ii)  $C_\alpha = M_\alpha$       (iii)  $C_\alpha = D_\alpha$ .

**Proof.** (i) $\Leftrightarrow$ (ii) If  $\alpha$  is norm continuous, then  $C_\alpha$  contains the unit  $\alpha(e)$  by Proposition 3.3. Hence,  $C_\alpha$  is an ideal in  $M_\alpha$ , containing the unit. Conversely, if  $C_\alpha = M_\alpha$  then  $C_\alpha$  is unital, and norm continuity of  $\alpha$  follows from Proposition 3.3.

(i) $\Rightarrow$ (iii) Since (ii) holds, the inclusion  $C_\alpha \supset D_\alpha$  is clear. To obtain the opposite inclusion, note that for any continuous function of compact support  $\varphi$  – the family of which is dense in  $L^1(G)$  – the function  $s \mapsto \varphi(s)\alpha(s)$ , from  $G$  to  $D_\alpha$ , can be uniformly approximated by Borel simple functions. Hence  $\alpha_1(\varphi) = \int_G \varphi(s)\alpha(s)ds$  may be regarded as a Bochner integral, and is thus in  $D_\alpha$ , since each  $\alpha(s) \in D_\alpha$ . It then follows that  $\alpha_1(L^1(G)) \subset D_\alpha$  and hence  $C_\alpha \subset D_\alpha$ .

(iii) $\Rightarrow$ (i) Since  $C_\alpha \supset D_\alpha$ ,  $C_\alpha$  is unital, and the result follows from Proposition 3.3.  $\square$

Now suppose that  $\pi : G \rightarrow \mathcal{U}(\mathcal{H}) \subset \mathcal{B}(\mathcal{H})_{\text{inv}}$  is a strongly continuous unitary representation (which is equivalent to it being weak\* continuous). We will define  $\pi_1 : M(G) \rightarrow \mathcal{B}(\mathcal{H})$  as above, but will use the notation

$$M_\pi^* = \overline{\pi_1(M(G))}, \quad C_\pi^* = \overline{\pi_1(L^1(G))} \quad \text{and} \quad D_\pi^* = \overline{\pi_1(\ell^1(G))}$$

to indicate that these are C\*-algebras. Using von Neumann's double commutant theorem, we have that  $C_\pi^*$  and  $D_\pi^*$  each generate the same von Neumann algebra,  $\text{VN}_\pi$ . We note that  $M_\pi^* \subset \text{VN}_\pi$  but  $M_\pi^* \neq \text{VN}_\pi$  in general. Thus, in particular, there is no reason to suspect that  $M_\pi^*$  is a dual space.

**Proposition 3.5.** *If  $\mu \in M(G)$ , then*

$$(3.3) \quad \Gamma_\pi(\mu) = \int_G \pi(s) \otimes \pi(s)^* d\mu(s)$$

*defines an element of  $\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$ , and the integral converges in the weak\* topology, i.e. for each  $x$  in  $\mathcal{B}(\mathcal{H})_* \otimes^h \mathcal{B}(\mathcal{H})_*$ ,*

$$\langle \Gamma_\pi(\mu), x \rangle = \int_G \langle \pi(s) \otimes \pi(s)^*, x \rangle d\mu(s).$$

Moreover,

(i)  $\Gamma_\pi : M(G) \rightarrow \mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$  is a contractive homomorphism whose range is contained in the algebra  $M_\pi^* \otimes^{eh} M_\pi^*$ .

(ii)  $\Gamma_\pi(L^1(G)) \subset C_\pi^* \otimes^{eh} C_\pi^*$ .

(iii)  $\Gamma_\pi(\ell^1(G)) \subset D_\pi^* \otimes^h D_\pi^*$ .

(iv) If  $\pi$  is norm continuous, then  $\Gamma_\pi(\mathbf{M}(G)) \subset \mathbf{D}_\pi^* \otimes^h \mathbf{D}_\pi^*$ .

**Proof. (i)** First, let us see that, for each  $\mu$  in  $\mathbf{M}(G)$ , the integral in (3.3) converges as claimed. This amounts to verifying that  $s \mapsto \pi(s) \otimes \pi(s)^*$  is a weak\* continuous representation from  $G$  into  $(\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H}))_{\text{inv}}$ , i.e. that  $s \mapsto \langle \pi(s) \otimes \pi(s)^*, x \rangle$  is continuous for each  $x$  in  $\mathcal{B}(\mathcal{H})_* \otimes^h \mathcal{B}(\mathcal{H})_*$ , by (2.1). If  $x \in \mathcal{B}(\mathcal{H})_* \otimes^h \mathcal{B}(\mathcal{H})_*$  and  $\varepsilon > 0$ , then there is  $x_\varepsilon$  in  $\mathcal{B}(\mathcal{H})_* \otimes \mathcal{B}(\mathcal{H})_*$  such that  $\|x - x_\varepsilon\|_h < \varepsilon$ . The function  $x_{\varepsilon, \pi}$ , given by  $s \mapsto \langle \pi(s) \otimes \pi(s)^*, x_\varepsilon \rangle$ , is clearly continuous on  $G$ , and  $\|x_\pi - x_{\varepsilon, \pi}\|_\infty \leq \|x - x_\varepsilon\|_h < \varepsilon$ . Thus, taking choices of  $\varepsilon$  tending to 0, we see that  $x_\pi$  is a continuous function on  $G$ .

Since  $\|\pi(s) \otimes \pi(s)^*\|_{w^*h} = 1$  for each  $s$  in  $G$ , the contractivity of  $\Gamma_\pi$  follows from (3.2). That  $\Gamma_\pi$  is a homomorphism follows from Proposition 3.1 and Example 3.2 (iii).

To see that  $\Gamma_\pi(\mu) \in \mathbf{M}_\pi^* \otimes^{eh} \mathbf{M}_\pi^*$ , for any given  $\mu$  in  $\mathbf{M}(G)$ , we will inspect the image of a typical weak\*-weak\* continuous left slice map on  $\Gamma_\pi(\mu)$  and use [22, Thm. 2.2]. If  $\omega \in \mathcal{B}(\mathcal{H})_*$ , then

$$(3.4) \quad L_\omega(\Gamma_\pi(\mu)) = \int_G \langle \pi(s), \omega \rangle \pi(s)^* d\mu(s) = \int_G \pi(s) d(\omega_\pi \mu)^\vee(s) \in \mathbf{M}_\pi^*$$

where  $\omega_\pi \mu$  is the measure with Radon derivative  $d(\omega_\pi \mu)/d\mu = \omega_\pi$  (here  $\omega_\pi(s) = \langle \pi(s), \omega \rangle$ ), and  $\nu^\vee(E) = \nu(E^{-1}) = \overline{\nu^*(E)}$  for any Borel measure  $\nu$ . The computation for any right slice map is similar.

(ii) This follows from a computation similar to (3.4).

(iii) If  $\mu = \sum_{s \in G} \alpha(s) \delta_s$ , where  $\sum_{s \in G} |\alpha(s)| < \infty$ , then since  $\pi(s) \otimes \pi(s)^* \in \mathbf{D}_\pi^* \otimes^h \mathbf{D}_\pi^*$  for each  $s$  in  $G$ , it follows too that

$$\Gamma_\pi(\mu) = \sum_{s \in G} \alpha(s) \pi(s) \otimes \pi(s)^* \in \mathbf{D}_\pi^* \otimes^h \mathbf{D}_\pi^*.$$

(iv) If we let  $\alpha : G \rightarrow (\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H}))_{\text{inv}}$  be given by  $\alpha(s) = \pi(s) \otimes \pi(s)^*$ , then  $\alpha$  is norm continuous. Hence

$$\Gamma_\pi(\mathbf{M}(G)) \subset \mathbf{M}_\alpha = \mathbf{D}_\alpha \subset \mathbf{D}_\pi^* \otimes^h \mathbf{D}_\pi^*$$

by (iii) above and Corollary 3.4.  $\square$

**Remark 3.6.** We note that (3.3) also converges in the  $\mathcal{CB}(\mathcal{B}(\mathcal{H}))\text{--}(\mathcal{B}(\mathcal{H})\widehat{\otimes}\mathcal{B}(\mathcal{H})_*)$  topology. Indeed, if  $a \in \mathcal{B}(\mathcal{H})$  and  $\eta^* \otimes \xi$  is a vector functional in  $\mathcal{B}(\mathcal{H})_*$ , then for any  $s$  in  $G$  we have that

$$\langle \pi(s) \otimes \pi(s)^*, a \otimes (\eta^* \otimes \xi) \rangle = \langle \pi(s)a\pi(s)^*, \eta^* \otimes \xi \rangle = \langle a, (\pi(s)^*\eta)^* \otimes \pi(s)^*\xi \rangle$$

where  $s \mapsto (\pi(s)^*\eta)^* \otimes \pi(s)^*\xi$  is continuous in the norm topology of  $\mathcal{B}(\mathcal{H})_*$ . Hence  $s \mapsto \langle \pi(s) \otimes \pi(s)^*, a \otimes (\eta^* \otimes \xi) \rangle$  is continuous. In particular, for each  $a$  in  $\mathcal{B}(\mathcal{H})$  and  $\mu$  in  $M(G)$  we have that

$$\Gamma_\pi(\mu)a = \int_G \pi(s)a\pi(s)^*d\mu(s)$$

where the integral converges in the weak\* topology of  $\mathcal{B}(\mathcal{H})$

We observe that it is possible, for each  $\mu$  in  $M(G)$ , to see  $\Gamma_\pi(\mu)|_{\mathcal{K}(\mathcal{H})}$  as an integral converging in the point-norm topology. However, our approach for obtaining (3.3) better lends itself to (4.4).  $\square$

We let the *augmentation ideal* in  $L^1(G)$  be given by

$$I_0(G) = \left\{ f \in L^1(G) : \int_G f(s)ds = 0 \right\}.$$

**Theorem 3.7.** *For any strongly continuous representation  $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$ , the following are equivalent:*

- (i)  $\pi$  is norm continuous.
- (ii)  $\Gamma_\pi(L^1(G)) \subset C_\pi^* \otimes^h C_\pi^*$ .
- (iii) there is an  $f$  in  $L^1(G) \setminus I_0(G)$  such that  $\Gamma_\pi(f) \in C_\pi^* \otimes^h C_\pi^*$ .

**Proof.** That (i) implies (ii) follows from Proposition 3.5 (iv) and the fact that  $C_\pi^* = D_\pi^*$ . That (ii) implies (iii) is trivial. Suppose now that  $f$  satisfies statement (iii). Without loss of generality, we may suppose that  $\int_G f(s)ds = 1$ . Then by [4], there exist sequences  $\{a_i\}_{i \in \mathbb{N}}$  and  $\{b_i\}_{i \in \mathbb{N}}$  from  $C_\pi^*$  such that  $\sum_{i=1}^\infty a_i a_i^*$  and  $\sum_{i=1}^\infty b_i^* b_i$  converge in norm, and

$$\Gamma_\pi(f)x = \sum_{i=1}^\infty a_i x b_i$$

for each  $x \in \mathcal{B}(\mathcal{H})$ . But it then follows from Remark 3.6 that

$$I = \int_G f(s)\pi(s)I\pi(s)^*ds = \Gamma_\pi(f)I = \sum_{i=1}^{\infty} a_i b_i \in \mathbf{C}_\pi^*.$$

Hence  $\pi$  is norm continuous by Proposition 3.3.  $\square$

In the next section, we will address the necessity of the assumption that  $f \in L^1(G) \setminus I_0(G)$  in (iii) above.

It is interesting to note that the kernel of  $\Gamma_\pi$  is related to the kernel of a more familiar representation. Below, we will let  $\overline{\mathcal{H}}$  denote the conjugate Hilbert space and  $\overline{\pi} : G \rightarrow \mathcal{U}(\overline{\mathcal{H}})$  denote the conjugate representation. We will also let  $\pi \otimes \overline{\pi} : G \rightarrow \mathcal{U}(\mathcal{H} \otimes_2 \overline{\mathcal{H}})$  be the usual tensor product of representations on the Hilbert space  $\mathcal{H} \otimes_2 \overline{\mathcal{H}}$ .

**Proposition 3.8.**  $\ker \Gamma_\pi = \ker(\pi \otimes \overline{\pi})_1$ .

**Proof.** We have that  $\mu \in \ker \Gamma_\pi$  if and only if

$$0 = \langle \Gamma_\pi(\mu), \omega_{\xi, \eta} \otimes \omega_{\zeta, \vartheta} \rangle$$

for every elementary tensor of vector functionals  $\omega_{\xi, \eta} \otimes \omega_{\zeta, \vartheta}$  in  $\mathcal{B}(\mathcal{H})_* \otimes^h \mathcal{B}(\mathcal{H})_*$ . (Note that we earlier had used the notation  $\omega_{\xi, \eta} = \eta^* \otimes \xi$ .) We may compute

$$\begin{aligned} \langle \Gamma_\pi(\mu), \omega_{\xi, \eta} \otimes \omega_{\zeta, \vartheta} \rangle &= \int_G \langle \pi(s)\xi | \eta \rangle \langle \pi(s)^*\zeta | \vartheta \rangle d\mu(s) \\ &= \int_G \langle \pi(s)\xi | \eta \rangle \overline{\langle \pi(s)\vartheta | \zeta \rangle} d\mu(s) \\ &= \int_G \langle \pi \otimes \overline{\pi}(s) \xi \otimes \overline{\vartheta} | \eta \otimes \overline{\zeta} \rangle d\mu(s) \\ &= \langle (\pi \otimes \overline{\pi})_1(\mu) \xi \otimes \overline{\vartheta} | \eta \otimes \overline{\zeta} \rangle. \end{aligned}$$

Thus it follows that  $\mu \in \ker \Gamma_\pi$  if and only if  $\mu \in \ker(\pi \otimes \overline{\pi})_1$ .  $\square$

In particular, if we let  $F_{\pi \otimes \overline{\pi}}$  be the linear space generated by all of the coefficient functions,  $s \mapsto \langle \pi \otimes \overline{\pi}(s) \xi \otimes \overline{\vartheta} | \eta \otimes \overline{\zeta} \rangle$ , we see that  $\mu \in \ker \Gamma_\pi$  exactly when  $\mu$ , as a functional on  $\mathcal{C}_b(G)$ , annihilates  $F_{\pi \otimes \overline{\pi}}$ .

## 4. ABELIAN GROUPS

For this section we let  $G$  be a locally compact *abelian* group, and we let  $\widehat{G}$  denote its topological dual group. For each  $s$  in  $G$ , we will let  $\hat{s}$  denote the associated unitary character on  $\widehat{G}$ , defined by  $\hat{s}(\sigma) = \sigma(s)$  for each  $\sigma$  in  $\widehat{G}$ .

As above, we will let  $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$  be a strongly continuous unitary representation. We let  $E_\pi$  denote the spectrum of  $C_\pi^*$ . Since  $C_\pi^*$  is a quotient of the enveloping  $C^*$ -algebra  $C^*(G)$ , and  $C^*(G) \cong \mathcal{C}_0(\widehat{G})$ , we may consider  $E_\pi$  to be a closed subset of  $\widehat{G}$ . Moreover, the natural isomorphism  $F_\pi : C_\pi^* \rightarrow \mathcal{C}_0(E_\pi)$  satisfies

$$F_\pi(\pi_1(f)) = \hat{f}|_{E_\pi}$$

for each  $f$  in  $L^1(G)$ , where  $\hat{f}(\sigma) = \sigma_1(f) = \int_G f(s)\sigma(s)ds$  for each  $\sigma$  in  $\widehat{G}$ . We note that our notation  $f \mapsto \hat{f}$ , for the Fourier transform, differs from that of our main reference, [15].

We would like to be able to extend  $F_\pi$  to some suitable map  $\bar{F}_\pi$  on  $\text{VN}_\pi$ . It is not clear that this can be done in general, but it can be done in many cases.

**Lemma 4.1.** *Consider the following conditions for  $\pi$  or  $G$  below:*

- (a)  $\mathcal{H}$  admits a maximal countable family of mutually orthogonal cyclic subspaces for  $\pi$ .
- (b) There is a family  $\{U_i\}_{i \in I}$  of separable open subsets of  $E_\pi$  such that  $E_\pi = \dot{\bigcup}_{i \in I} U_i$ .
- (c)  $\widehat{G}$  has a separable open subgroup.
- (d)  $G$  is compactly generated.

Then, under any one of these conditions there exists a regular Borel measure  $\nu$  on  $E_\pi$ , bounded on compacta, such that there is a normal  $*$ -homomorphism  $\bar{F}_\pi : \text{VN}_\pi \rightarrow L^\infty(E_\pi, \nu)$  which extends  $F_\pi$ .

**Proof. (a)** By standard arguments (see [5, §7], for example),  $\text{VN}_\pi$  admits a faithful normal state  $\omega$ . Then the measure  $\nu$  given by

$$(4.1) \quad \int_{E_\pi} \varphi(\sigma) d\nu(\sigma) = \omega(F_\pi^{-1}\varphi)$$

for each  $\varphi$  in  $\mathcal{C}_0(E_\pi)$ , gives rise to the desired map  $\bar{F}_\pi$ .

**(b)** Since  $E_\pi = \dot{\bigcup}_{i \in I} U_i$ , we have that  $\mathcal{C}_0(E_\pi) = c_0\text{-}\bigoplus_{i \in I} \mathcal{C}_0(U_i)$ . If we let  $\mathcal{C}_i = F_\pi^{-1}(\mathcal{C}_0(U_i))$ , then  $\mathcal{M}_i = \overline{\mathcal{C}_i}^{w^*}$  is an ideal in  $\text{VN}_\pi$ . The ideals  $\mathcal{M}_i$  are mutually orthogonal, and hence if  $\{p_i\}_{i \in I}$  is the family of projections for which  $\mathcal{M}_i = p_i \text{VN}_\pi$  for each  $i$ , then  $\sum_{i \in I} p_i = I$ . Since each  $\mathcal{C}_i$  is separable, each  $\mathcal{M}_i$  is countably generated, and hence there is a normal state  $\omega_i$  on  $\text{VN}_\pi$  with support projection  $p_i$ . Let  $\nu_i$  be the measure on  $E_\pi$  associated with  $\omega_i$  as in (4.1). Then  $\text{supp}(\nu_i) = U_i$  for each  $i$ , and  $\nu = \bigoplus_{i \in I} \nu_i$  is the desired measure.

**(c)** If  $\widehat{G}$  has a separable open subgroup  $X$ , let  $T$  be any transversal for  $X$  in  $\widehat{G}$ , and we have that  $E_\pi = \dot{\bigcup}_{\tau \in T} (E_\pi \cap \tau X)$ , and again we obtain (b).

**(d)** If  $G$  is compactly generated, then by [15, 9.8] there is a topological isomorphism  $G \cong \mathbb{Z}^n \times \mathbb{R}^m \times K$ , where  $K$  is compact. Then  $\widehat{G} \cong \mathbb{T}^n \times \mathbb{R}^m \times \widehat{K}$ , and the subgroup  $X$  corresponding to  $\mathbb{T}^n \times \mathbb{R}^m$  is open and separable, and hence (c) holds.  $\square$

We will need to use an extension of  $F_\pi$  of a different nature than in the lemma above. Since  $\mathcal{C}_\pi^*$  is an essential ideal in  $\mathcal{M}_\pi^*$ , the map  $F_\pi : \mathcal{C}_\pi^* \rightarrow \mathcal{C}_0(E_\pi)$  extends to an injective  $*$ -homomorphism  $\tilde{F}_\pi : \mathcal{M}_\pi^* \rightarrow \mathcal{C}_b(E_\pi)$ , such that  $F_\pi(na) = \tilde{F}_\pi(n)F_\pi(a)$  for each  $n$  in  $\mathcal{M}_\pi^*$  and  $a$  in  $\mathcal{C}_\pi^*$ , by [18, 3.12.8]. We note that for each  $\mu$  in  $\mathcal{M}(G)$ ,

$$(4.2) \quad \tilde{F}_\pi(\pi_1(\mu)) = \hat{\mu}|_{E_\pi}$$

where for each  $\mu$  in  $\mathcal{M}(G)$ ,  $\hat{\mu}(\sigma) = \sigma_1(\mu) = \int_G \sigma(s) d\mu(s)$ . Thus  $\mu \mapsto \hat{\mu}$  is the Fourier-Stieltjes transform. To see the validity of (4.2), observe that for each  $f$  in  $L^1(G)$  we have

$$\hat{\mu} \hat{f}|_{E_\pi} = \widehat{\mu * f}|_{E_\pi} = F_\pi(\pi_1(\mu * f)) = \tilde{F}_\pi(\pi_1(\mu)) F_\pi(f) = \tilde{F}_\pi(\pi_1(\mu)) \hat{f}|_{E_\pi}.$$

Thus it follows that  $\tilde{F}_\pi(\pi_1(\mu))\varphi = \hat{\mu}\varphi$  for each  $\varphi$  in  $\mathcal{C}_0(E_\pi)$ .

If any of the conditions of Lemma 4.1 hold, then there exists a measure  $\nu$  for which there is a normal extension  $\bar{F}_\pi : \text{VN}_\pi \rightarrow L^\infty(E_\pi, \nu)$  of  $F_\pi$ . Then for any  $\mu$  in  $M(G)$ ,

$$(4.3) \quad \bar{F}_\pi(\pi_1(\mu)) = \hat{\mu}|_{E_\pi}$$

where we identify  $\mathcal{C}_b(E_\pi)$  as a closed subspace of  $L^\infty(E_\pi, \nu)$ . To see (4.3), we note that if  $(a_\beta)$  is any bounded approximate identity in  $C_\pi^*$ , then  $\text{weak}^*\text{-}\lim_\beta a_\beta = I$  in  $\text{VN}_\pi$ , thus  $\text{weak}^*\text{-}\lim_\beta F_\pi(a_\beta) = 1_{E_\pi}$ . Hence

$$\begin{aligned} \bar{F}_\pi(\pi_1(\mu)) &= \text{weak}^*\text{-}\lim_\beta \bar{F}_\pi(\pi_1(\mu)a_\beta) = \text{weak}^*\text{-}\lim_\beta F_\pi(\pi_1(\mu)a_\beta) \\ &= \text{weak}^*\text{-}\lim_\beta \tilde{F}_\pi(\pi_1(\mu))F_\pi(a_\beta) = \tilde{F}_\pi(\pi_1(\mu)) = \hat{\mu}|_{E_\pi}. \end{aligned}$$

We will make use of the spaces  $V^b(E)$ ,  $V^0(E)$  and  $V_0(E)$ , which were defined in (2.3). If  $\nu$  is any non-negative measure on  $E$ , we let

$$V^\infty(E, \nu) = L^\infty(E, \nu) \otimes^{eh} L^\infty(E, \nu).$$

Spaces of this type are discussed in [22].

**Theorem 4.2.** *If  $G$  is a locally compact abelian group and  $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$  is a strongly continuous unitary representation, then for any  $\mu$  in  $M(G)$  and  $(\sigma, \tau)$  in  $E_\pi \times E_\pi$  we have that*

$$(\tilde{F}_\pi \otimes \tilde{F}_\pi)\Gamma_\pi(\mu)(\sigma, \tau) = \hat{\mu}(\sigma\tau^{-1}).$$

*In particular, if  $E$  is any closed subset of  $\hat{G}$  and  $\mu \in M(G)$ , then  $u(\sigma, \tau) = \hat{\mu}(\sigma\tau^{-1})$  is an element of  $V^b(E)$ , and  $u \in V^0(E)$  if  $\mu \in L^1(G)$ . Moreover, if  $E$  is compact then  $u \in V_0(E)$ .*

**Proof.** The result will be established in three stages. The first two of these require additional hypotheses and are preparatory for the general case.

**I.** Suppose that any one of the conditions of Lemma 4.1 is satisfied. Let  $\nu$  be the measure on  $E_\pi$  and let  $\bar{F}_\pi : \text{VN}_\pi \rightarrow L^\infty(E_\pi, \nu)$  be the map given there.

If  $\mu \in M(G)$ , we have that

$$\begin{aligned}
 (\tilde{F}_\pi \otimes \tilde{F}_\pi)\Gamma_\pi(\mu) &= (\bar{F}_\pi \otimes \bar{F}_\pi) \int_G \pi(s) \otimes \pi(s)^* d\mu(s) \\
 (4.4) \qquad \qquad \qquad &= \int_G \hat{s}|_{E_\pi} \otimes \bar{\hat{s}}|_{E_\pi} d\mu(s)
 \end{aligned}$$

where the latter integral converges in the weak\* topology of  $V^\infty(E_\pi, \nu)$ .

For  $(\sigma, \tau)$  in  $E_\pi \times E_\pi$  let

$$u(\sigma, \tau) = \hat{\mu}(\sigma\tau^{-1}).$$

Then  $u \in V^b(E_\pi)$ . Indeed, we have that  $\hat{\mu} \in B(\widehat{G})$ , the *Fourier-Stieltjes algebra* which is defined in [10]. Thus there is a Hilbert space  $\mathcal{L}$ , a continuous unitary representation  $\rho : G \rightarrow \mathcal{U}(\mathcal{L})$ , and vectors  $\xi, \eta$  in  $\mathcal{L}$  with  $\|\mu\| = \|\xi\| \|\eta\|$ , such that  $\hat{\mu}(\sigma) = \langle \rho(\sigma)\xi | \eta \rangle$  for each  $\sigma$  in  $\widehat{G}$ . If  $\{\xi_i\}_{i \in I}$  is an orthonormal basis for  $\mathcal{L}$ , then we have, using Parseval's formula, that

$$\hat{\mu}(\sigma\tau^{-1}) = \langle \rho(\sigma\tau^{-1})\xi | \eta \rangle = \sum_{i \in I} \langle \rho(\sigma)\xi | \xi_i \rangle \langle \xi_i | \rho(\tau)\eta \rangle$$

for any  $(\sigma, \tau)$  in  $E_\pi \times E_\pi$ . Hence

$$u = \sum_{i \in I} \langle \rho(\cdot)\xi | \xi_i \rangle \otimes \overline{\langle \rho(\cdot)\eta | \xi_i \rangle} \in V^b(E_\pi)$$

with  $\|u\|_{eh} \leq \|\xi\| \|\eta\| = \|\mu\|$ . (This is similar to the proof of [23, Prop. 5.1].) We note that if  $\mu \in L^1(G)$ , then  $\rho$  can be taken to be the left regular representation and hence each  $\langle \rho(\cdot)\xi | \xi_i \rangle$  and  $\langle \rho(\cdot)\eta | \xi_i \rangle$  is in  $\mathcal{C}_0(E_\pi)$ . Hence, in this case we would have that  $u \in V^0(E_\pi)$ .

We wish to establish that

$$(4.5) \qquad \qquad \qquad u = (\tilde{F}_\pi \otimes \tilde{F}_\pi)\Gamma_\pi(\mu).$$

We will do this by using the dual pairing (2.1). If  $g \otimes h$  is an elementary tensor in  $L^1(E_\pi, \nu) \otimes^h L^1(E_\pi, \nu)$ , then

$$\begin{aligned} \langle u, g \otimes h \rangle &= \int_{E_\pi} \int_{E_\pi} g(\sigma) h(\tau) \hat{\mu}(\sigma\tau^{-1}) d\nu(\sigma) d\nu(\tau) \\ &= \int_{E_\pi} \int_{E_\pi} g(\sigma) h(\tau) \left( \int_G \sigma(s) \overline{\tau(s)} d\mu(s) \right) d\nu(\sigma) d\nu(\tau) \\ &= \int_G \left( \int_{E_\pi} g(\sigma) \hat{s}(\sigma) d\nu(\sigma) \right) \left( \int_{E_\pi} h(\tau) \overline{\hat{s}(\tau)} d\nu(\tau) \right) d\mu(s) \end{aligned}$$

where the version of Fubini's Theorem required is [15, 13.10], noting that  $g$  and  $h$  each have  $\nu$ - $\sigma$ -finite supports. On the other hand, by (4.4),

$$\begin{aligned} \langle (\tilde{F}_\pi \otimes \tilde{F}_\pi) \Gamma_\pi(\mu), g \otimes h \rangle &= \left\langle \int_G \hat{s}|_{E_\pi} \otimes \overline{\hat{s}}|_{E_\pi} d\mu(s), g \otimes h \right\rangle \\ &= \int_G \left( \int_{E_\pi} g(\sigma) \hat{s}(\sigma) d\nu(\sigma) \right) \left( \int_{E_\pi} h(\tau) \overline{\hat{s}(\tau)} d\nu(\tau) \right) d\mu(s) \end{aligned}$$

and this shows that (4.5) holds.

**II.** Suppose that  $\mu$  is supported on a compactly generated open subgroup  $H$  of  $G$ .

Let us first compute the spectrum  $E_{\pi|_H}$  of  $C_{\pi|_H}^*$ . We note that  $\widehat{H} = \widehat{G}|_H$  and that the restriction map  $r : \widehat{G} \rightarrow \widehat{G}|_H$  is a homomorphic topological quotient map by [15, 24.5]. Moreover,  $\ker r$  is compact, by [15, 23.29(a)]. Then  $E_{\pi|_H} = r(E_\pi)$ . To see this, observe that the map  $\iota : L^1(H) \rightarrow L^1(G)$ , which we define to be the inverse of  $f \mapsto f|_H$ , extends to an injective  $*$ -homomorphism  $\iota_\pi : C_{\pi|_H}^* \rightarrow C_\pi^*$ . In particular, then, each multiplicative linear functional on  $C_{\pi|_H}^*$  is necessarily the restriction of such a functional on  $C_\pi^*$ . Let  $r_\pi = r|_{E_\pi}$ . Then, the map  $r_\pi : E_\pi \rightarrow r(E_\pi)$  induces an injective  $*$ -homomorphism  $j_{r_\pi} : \mathcal{C}_0(r(E_\pi)) \rightarrow \mathcal{C}_0(E_\pi)$ , whose image is the subalgebra of all functions which are constant on relative cosets of  $\ker r$  in  $E_\pi$ . Now, if  $g \in L^1(H)$ , and  $\sigma \in E_\pi$  then

$$(4.6) \quad \widehat{\iota g}(\sigma) = \int_G \iota g(s) \sigma(s) ds = \int_H g(s) r(\sigma)(s) ds = \widehat{g}(r_\pi(\sigma)) = j_{r_\pi} \widehat{g}(\sigma)$$

from which it follows that every character on  $C_{\pi|H}^*$  is from  $r_\pi(E_\pi)$ . Moreover, it follows from (4.6) that

$$F_\pi \circ \iota_\pi = j_{r_\pi} \circ F_{\pi|H}.$$

Now, we let  $\tilde{\iota} : M(H) \rightarrow M(G)$  be the homomorphism whose inverse is  $\kappa \mapsto \kappa_H$ , where for any Borel subset  $B$  of  $G$ ,  $\kappa_H(B) = \kappa(B \cap H)$ . Then  $\iota$  induces an injective  $*$ -homomorphism  $\tilde{\iota}_\pi : M_{\pi|H}^* \rightarrow M_\pi^*$ . It follows from the discussion above that  $\tilde{F}_{\pi|H} : M_{\pi|H}^* \rightarrow \mathcal{C}_b(r(E_\pi))$ . Then

$$(4.7) \quad \tilde{j}_{r_\pi} \circ \tilde{F}_{\pi|H} = \tilde{F}_\pi \circ \tilde{\iota}_\pi$$

where  $\tilde{j}_{r_\pi} : \mathcal{C}_b(E_{\pi|H}) \rightarrow \mathcal{C}_b(E_\pi)$  is the map induced by  $r_\pi : E_\pi \rightarrow E_{\pi|H}$ . Indeed, if  $\kappa \in M(H)$ , then for each  $\sigma$  in  $E_\pi$  we have that  $\widehat{\tilde{\iota}\kappa}(\sigma) = \tilde{j}_{r_\pi} \hat{\kappa}(\sigma)$ , by a computation analagous to (4.6), above. Next, we wish to establish that

$$(4.8) \quad \Gamma_\pi \circ \tilde{\iota} = (\tilde{\iota}_\pi \otimes \tilde{\iota}_\pi) \circ \Gamma_{\pi|H}.$$

If  $\kappa \in M(H)$  and  $x \in \mathcal{B}(\mathcal{H}) \otimes^h \mathcal{B}(\mathcal{H})$ , then

$$\begin{aligned} \langle \Gamma_\pi(\tilde{\iota}\kappa), x \rangle &= \int_G \langle \pi(s) \otimes \pi(s)^*, x \rangle d\tilde{\iota}\kappa(s) \\ &= \int_H \langle \pi(s) \otimes \pi(s)^*, x \rangle d\kappa(s) = \langle \Gamma_{\pi|H}(\kappa), x \rangle \end{aligned}$$

whence, as elements of  $\mathcal{B}(\mathcal{H}) \otimes^{eh} \mathcal{B}(\mathcal{H})$ ,  $\Gamma_\pi(\tilde{\iota}\kappa) = \Gamma_{\pi|H}(\kappa)$ . However, in  $\mathcal{B}(\mathcal{H})$ , the inclusion map  $M_{\pi|H}^* \hookrightarrow M_\pi^*$  is the map  $\tilde{\iota}_\pi$ , and thus (4.8) holds.

Now, since  $\mu$  is supported on  $H$ , we have that  $\mu = \tilde{\iota}\kappa$  for some  $\kappa \in M(H)$ .

Then for each  $(\sigma, \tau)$  in  $E_\pi \times E_\pi$  we have that

$$\begin{aligned} \hat{\mu}(\sigma\tau^{-1}) &= \widehat{\tilde{\iota}\kappa}(\sigma\tau^{-1}) = \hat{\kappa}(r(\sigma\tau^{-1})) = \hat{\kappa}(r_\pi(\sigma)r_\pi(\tau^{-1})) \\ &= (\tilde{F}_{\pi|H} \otimes \tilde{F}_{\pi|H})\Gamma_{\pi|H}(\kappa)(r_\pi(\sigma), r_\pi(\tau)), \quad \text{by part I} \\ &= (\tilde{j}_{r_\pi} \otimes \tilde{j}_{r_\pi})(\tilde{F}_{\pi|H} \otimes \tilde{F}_{\pi|H})\Gamma_{\pi|H}(\kappa)(\sigma, \tau) \\ &= (\tilde{F}_\pi \otimes \tilde{F}_\pi)(\tilde{\iota}_\pi \otimes \tilde{\iota}_\pi)\Gamma_{\pi|H}(\kappa)(\sigma, \tau), \quad \text{by (4.7)} \\ &= (\tilde{F}_\pi \otimes \tilde{F}_\pi)\Gamma_\pi(\tilde{\iota}\kappa)(\sigma, \tau), \quad \text{by (4.8)} \\ &= (\tilde{F}_\pi \otimes \tilde{F}_\pi)\Gamma_\pi(\mu)(\sigma, \tau). \end{aligned}$$

**III.** We now cover the case of a general  $\mu$  in  $M(G)$ .

Let  $U$  be a relatively compact symmetric open neighbourhood of the identity in  $G$ . Then  $H = \bigcup_{n=1}^{\infty} U^n$  is a compactly generated open subgroup of  $G$ . We note that if  $T$  is a transversal for  $H$  in  $G$  then

$$\mu = \sum_{t \in T} \mu_{tH}$$

which is an absolutely summable series. For each  $t$  in  $T$  let

$$\mu_t = \delta_{t^{-1}} * (\mu_{tH})$$

so  $\text{supp}(\mu_t) \subset H$  and  $\mu = \sum_{t \in T} \delta_t * \mu_t$ . We then have that

$$\begin{aligned} (\tilde{F}_\pi \otimes \tilde{F}_\pi) \Gamma_\pi(\mu) &= \sum_{t \in T} (\tilde{F}_\pi \otimes \tilde{F}_\pi) \Gamma_\pi(\delta_t * \mu_t) \\ &= \sum_{t \in T} (\tilde{F}_\pi \otimes \tilde{F}_\pi) [(\pi(t) \otimes \pi(t)^*) \Gamma_\pi(\mu_t)] \\ &= \sum_{t \in T} (\hat{t}|_{E_\pi} \otimes \tilde{t}|_{E_\pi}) (\tilde{F}_\pi \otimes \tilde{F}_\pi) \Gamma_\pi(\mu_t). \end{aligned}$$

Hence if  $(\sigma, \tau) \in E_\pi \times E_\pi$ , we obtain

$$\begin{aligned} (\tilde{F}_\pi \otimes \tilde{F}_\pi) \Gamma_\pi(\mu)(\sigma, \tau) &= \sum_{t \in T} \hat{t}(\sigma) \overline{\hat{t}(\tau)} (\tilde{F}_\pi \otimes \tilde{F}_\pi) \Gamma_\pi(\mu_t)(\sigma, \tau) \\ &= \sum_{t \in T} \hat{t}(\sigma) \overline{\hat{t}(\tau)} \hat{\mu}_t(\sigma \tau^{-1}), \quad \text{by part II} \\ &= \sum_{t \in T} \hat{\delta}_t(\sigma \tau^{-1}) \hat{\mu}_t(\sigma \tau^{-1}) \\ &= \sum_{t \in T} \widehat{\delta_t * \mu_t}(\sigma \tau^{-1}) = \hat{\mu}(\sigma \tau^{-1}). \end{aligned}$$

Thus our first claim is established in general.

If  $E$  is any closed subset of  $\widehat{G}$  then by [14, 33.7] there is a representation  $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$  for which  $E_\pi = E$ . Hence

$$u(\sigma, \tau) = \hat{\mu}(\sigma \tau^{-1}) = (\tilde{F}_\pi \otimes \tilde{F}_\pi) \Gamma_\pi(\mu)(\sigma, \tau)$$

defines an element of  $V^b(E)$ , and of  $V^0(E)$  if  $\mu \in L^1(G)$ . If  $E$  is compact, we note that any representation  $\pi$  for which  $E_\pi = E$ , is norm continuous

by Proposition 3.3, since  $C_\pi^* \cong C_0(E)$ , which is unital. Thus  $\Gamma_\pi(M(G)) = \Gamma_\pi(L^1(G)) \subset C_\pi^* \otimes^h C_\pi^*$ . Hence  $u$ , as above, is in  $V_0(E)$ .  $\square$

We can now obtain a generalisation of [24, Prop. 5.7]. This is a straightforward corollary of Theorems 2.4 and 4.2.

**Corollary 4.3.** *If  $G$  is a locally compact abelian group,  $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$  is a strongly continuous representation and  $\mu \in M(G)$ , then the following are equivalent:*

- (i)  $\Gamma_\pi(\mu)$  is positive.
- (ii)  $\Gamma_\pi(\mu)$  is completely positive.
- (iii)  $(\sigma, \tau) \mapsto \hat{\mu}(\sigma\tau^{-1})$  is positive definite on  $E_\pi \times E_\pi$ .

The next result follows directly from Theorem 4.2, but can also be deduced from Proposition 3.8.

**Corollary 4.4.** *If  $G$  is a locally compact abelian group and  $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$  is a strongly continuous representation then*

$$\ker \Gamma_\pi = \{\mu \in M(G) : \hat{\mu}|_{E_\pi E_\pi^{-1}} = 0\}.$$

Let us now address the assumption that  $f \in L^1(G) \setminus I_0(G)$  in Theorem 3.7 (iii). We want to show that having an  $f$  in  $I_0(G)$  for which  $\Gamma_\pi(f) \subset C_\pi^* \otimes^h C_\pi^*$  does not imply that  $\pi$  is norm continuous. First, following Corollary 4.4 we see that that  $\Gamma_\pi(f) = 0$  if the support of  $\hat{f}$  misses the difference set  $E_\pi E_\pi^{-1}$ . Thus it is possible that  $\Gamma_\pi(f) = 0 \in C_\pi^* \otimes^h C_\pi^*$ , though  $\pi$  need not be norm continuous, i.e.  $E_\pi$  need not be compact. Thus we may ask if  $\Gamma_\pi(L^1(G)) \cap (C_\pi^* \otimes^h C_\pi^*) = \{0\}$  when  $\pi$  is not norm continuous. However, this may not happen, as the next example shows.

**Example 4.5.** Let  $G = \mathbb{T}$ , and identify  $\widehat{\mathbb{T}} = \mathbb{Z}$ . Define  $\pi : \mathbb{T} \rightarrow \mathcal{U}(\ell^2(\mathbb{N}))$  for each  $z$  in  $\mathbb{T}$  by

$$\pi(z)(\xi_n)_{n \in \mathbb{N}} = \left( z^{n^2} \xi_n \right)_{n \in \mathbb{N}}.$$

Then  $E_\pi = \{n^2 : n \in \mathbb{N}\}$ , which is not compact in  $\mathbb{Z}$ . Hence  $C_\pi^* \cong c_0(E_\pi)$ , which is not unital, so  $\pi$  is not norm continuous on  $\mathbb{T}$ , by Proposition 3.3. Fix  $k$  in  $\mathbb{Z} \setminus \{0\}$  and let  $\hat{k}(z) = z^k$ . Then for each pair  $n, m$  in  $\mathbb{N}$ , using normalized Haar measure on  $\mathbb{T}$  and Theorem 4.2, we have that

$$(F_\pi \otimes F_\pi)\Gamma_\pi(\hat{k})(\hat{n}, \hat{m}) = \int_{\mathbb{T}} z^k z^{n^2} \bar{z}^{m^2} dz = \begin{cases} 1 & \text{if } m^2 - n^2 = k \\ 0 & \text{otherwise} \end{cases}.$$

The set of solutions to  $m^2 - n^2 = (m - n)(m + n) = k$  is clearly finite; we shall write them  $\{(n_1, m_1), \dots, (n_{l(k)}, m_{l(k)})\}$ . We then see that

$$(F_\pi \otimes F_\pi)\Gamma_\pi(\hat{k}) = \sum_{i=1}^{l(k)} 1_{(n_i, m_i)} = \sum_{i=1}^{l(k)} 1_{n_i} \otimes 1_{m_i} \in V_0(E_\pi).$$

Hence  $\Gamma_\pi(\hat{k}) \in C_\pi^* \otimes^h C_\pi^*$ . In fact, since  $I_0(\mathbb{T}) = \overline{\text{span}} \{\hat{k} : k \in \mathbb{Z} \setminus \{0\}\}$ , we have that  $\Gamma_\pi(I_0(\mathbb{T})) \subset C_\pi^* \otimes^h C_\pi^*$ .

We remark that for a general locally compact abelian group  $G$ , and representation  $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$ ,  $V_0(E_\pi) \subset \mathcal{C}_0(E_\pi \times E_\pi)$ . Thus if  $f$  in  $L^1(G)$  is such that  $\hat{f}(\sigma) \neq 0$  for some  $\sigma$  in  $\widehat{G}$  such that  $E_\pi \cap \sigma E_\pi$  is not compact, then  $\Gamma_\pi(f) \notin C_\pi^* \otimes^h C_\pi^*$  by Theorem 4.2. Thus if  $E_\pi \cap \sigma E_\pi$  is compact for no  $\sigma$  in  $\widehat{G}$  then we have that

$$\Gamma_\pi(L^1(G)) \cap (C_\pi^* \otimes^h C_\pi^*) = \{0\}.$$

Note that  $E_\pi \cap \sigma E_\pi$  is never compact if  $E_\pi = \widehat{G}$ , which occurs, for example when  $\pi$  is the left regular representation  $\lambda$ . It is shown in [24, Cor. 4.7] that  $\Gamma_\lambda$  is an isometry. This was extended to non-abelian groups in [11] and expanded upon in [16], while [17] contains a proof that  $\Gamma_\lambda$  is a complete isometry. An analogue for the Fourier algebra of an amenable group is shown in [23, Cor. 5.4].

**Question 4.6.** If  $G = \mathbb{R}$ , then  $[0, \infty) \cap (s + [0, \infty)) = [\min\{s, 0\}, \infty)$  is never compact. Thus if  $\pi$  is a representation of  $\mathbb{R}$  such that  $E_\pi = [0, \infty)$ , then  $\Gamma_\pi : M(\mathbb{R}) \rightarrow M_\pi^* \otimes^{eh} M_\pi^*$  is injective by Corollary 4.4. Is  $\Gamma_\pi$  isometric?

How about  $\Gamma_\pi|_{L^1(\mathbb{R})}$ ? More generally, under what conditions for an arbitrary abelian group  $G$  and representation  $\pi$  is  $\Gamma_\pi$ , or  $\Gamma_\pi|_{L^1(G)}$ , a quotient map?

## REFERENCES

- [1] S. D. Allen, A. M. Sinclair, and R. R. Smith. The ideal structure of the Haagerup tensor product of  $C^*$ -algebras. *J. Reine Angew. Math.*, 442:111–148, 1993.
- [2] D. P. Blecher. Geometry of the tensor product of  $C^*$ -algebras. *Math. Proc. Cambridge Phil. Soc.*, 104(1):119–127, 1988.
- [3] D. P. Blecher and V. I. Paulsen. Tensor products of operator spaces. *J. Funct. Anal.*, 99:262–292, 1991.
- [4] D. P. Blecher and R. R. Smith. The dual of the Haagerup tensor product. *J. London Math. Soc.*, 45(2):126–144, 1992.
- [5] J. Dixmier. *Les algèbres d'opérateurs dans l'espace Hilbertien*, volume 25 of *Cahiers scientifiques*. Gauthier-Villars, Paris, 1969.
- [6] E. G. Effros and A. Kisimoto. Module maps and Hochschild-Johnson cohomology. *Indiana U. Math. J.*, 36:257–276, 1987.
- [7] E. G. Effros, J. Kraus, and Z.-J. Ruan. On two quantized tensor products. In *Operator algebras, mathematical physics, and low-dimensional topology (Istanbul, 1991)*, pages 125–145, Wellesley, MA, 1993. A K Peters.
- [8] E. G. Effros and Z.-J. Ruan. *Operator Spaces*, volume 23 of *London Math. Soc., New Series*. Clarendon Press, Oxford Univ. Press, New York, 2000.
- [9] E. G. Effros and Z.-J. Ruan. Operator convolution algebras: an approach to quantum groups. To appear in *J. Operator Theory.*, 2002.
- [10] P. Eymard. L'algèbre de Fourier d'un groupe localement compact. *Bull. Soc. Math. France*, 92:181–236, 1964.
- [11] F. Ghahramani. Isometric representations of  $M(G)$  on  $\mathcal{B}(\mathcal{H})$ . *Glasgow Math. J.*, 23:119–122, 1982.
- [12] A. Guichardet. *Symmetric Hilbert Spaces and related topics*, volume 261 of *Lecture Notes in Math*. Springer, Berlin Heidelberg, 1972.
- [13] U. Haagerup. Decomposition of completely bounded maps on operator algebras. Unpublished, 1980.
- [14] E. Hewitt and K. A. Ross. *Abstract Harmonic Analysis II*, volume 152 of *Die Grundlehren der mathematischen Wissenschaften*. Springer, Berlin Heidelberg, 1970.
- [15] E. Hewitt and K. A. Ross. *Abstract Harmonic Analysis I*, volume 115 of *Die Grundlehren der mathematischen Wissenschaften*. Springer, New York, second edition, 1979.
- [16] M. Neufang. Isometric representations of convolution algebras as completely bounded module homomorphisms and a characterization of the measure algebra. Unpublished, 2001.

- [17] M. Neufang, Z.-J. Ruan, and N. Spronk. Completely isometric representations of  $M_{cb}A(G)$  and  $UCB(\hat{G})^*$ . *Trans. Amer. Math. Soc.*, to appear.
- [18] G. K. Pedersen. *C\*-Algebras and their Automorphism Groups*, volume 14 of *London Math. Soc. Monographs*. Academic Press, London, 1979.
- [19] S. Sakai. *C\*-algebras and W\*-algebras*. Classics in Mathematics. Springer, Berlin, 1998. Reprint of the 1971 edition.
- [20] V. Shulman and L. Turowska. Operator synthesis. 1. synthetic sets, bilattices and tensor algebras. *J. Funct. Anal.*, 209(2):293–331, 2004.
- [21] R. R. Smith. Completely bounded module maps and the Haagerup tensor product. *J. Funct. Anal.*, 102(1):156–175, 1991.
- [22] N. Spronk. Measurable Schur multipliers and completely bounded multipliers of the Fourier algebra. *Proc. London Math. Soc.*, 89(3):161–192, 2003.
- [23] N. Spronk and L. Turowska. Spectral synthesis and operator synthesis for compact groups. *J. London Math. Soc.*, 66:361–376, 2002.
- [24] E. Størmer. Regular Abelian Banach algebras of linear maps of operator algebras. *J. Funct. Anal.*, 37:331–373, 1980.
- [25] N. Th. Varopoulos. Tensor algebras and harmonic analysis. *Acta. Math.*, 119:51–112, 1967.

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY,  
COLLEGE STATION, TEXAS 77843-3368, U.S.A.

Current address of the second author:

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO,  
WATERLOO, ONTARIO, CANADA N2L 3G1.

*E-mail addresses:* rsmith@math.tamu.edu, nspronk@uwaterloo.ca