

## Section 1.2: The Dot Product

Let's start with two equivalent definitions of dot product.

DEFINITION 1. The **dot product** of two nonzero vectors  $\mathbf{a}$  and  $\mathbf{b}$  is the number

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta,$$

where  $\theta$  is the angle between the vectors  $\mathbf{a}$  and  $\mathbf{b}$ ,  $0 \leq \theta \leq \pi$ . If either  $\mathbf{a}$  or  $\mathbf{b}$  is  $\mathbf{0}$ , then we define  $\mathbf{a} \cdot \mathbf{b} = 0$ .

DEFINITION 2. The **dot product** of two given vectors  $\mathbf{a} = \langle a_1, a_2 \rangle$  and  $\mathbf{b} = \langle b_1, b_2 \rangle$  is the number

$$\mathbf{a} \cdot \mathbf{b} = a_1 b_1 + a_2 b_2.$$

Note that the formula from Definition 1 is often used not to compute a dot product but instead to find the angle between two vectors. Indeed, it implies:

$$\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}| |\mathbf{b}|} =$$

EXAMPLE 3. Given  $\mathbf{a} = \langle 2, -3 \rangle$  and  $\mathbf{b} = \langle 3, -4 \rangle$ .

(a) Compute the dot product of  $\mathbf{a}$  and  $\mathbf{b}$ .

(b) Determine the angle between  $\mathbf{a}$  and  $\mathbf{b}$ .

Note that

$$\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$$

$$\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$$

$$\mathbf{0} \cdot \mathbf{a} = 0$$

The dot product gives us a simple way for determining if two vectors are perpendicular (or orthogonal), namely,

**Two nonzero vectors  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal if and only if  $\mathbf{a} \cdot \mathbf{b} = 0$ . (Prove it!)**

EXAMPLE 4. Determine whether the given vectors are orthogonal, parallel, or neither.

(a)  $\langle 3, 4 \rangle$ ,  $\langle -8, 6 \rangle$

(b)  $\langle -7, -4 \rangle$ ,  $\langle 28, 16 \rangle$

(c)  $\langle 1, 1 \rangle$ ,  $\langle 2, 3 \rangle$

EXAMPLE 5. What is the dot product of  $12\mathbf{j}$  and  $11\mathbf{i}$ ?

DEFINITION 6. The **work** done by a force  $\mathbf{F}$  in moving an object from point  $A$  to point  $B$  is given by

$$W = \mathbf{F} \cdot \mathbf{D}$$

where  $\mathbf{D} = \overrightarrow{AB}$  is the distance the object has moved (or displacement).

EXAMPLE 7. Find the work done by a force of 50lb acting in the direction  $N30^\circ W$  in moving an object 10ft due west.



EXAMPLE 11. Given  $\mathbf{a} = \langle 4, 3 \rangle$  and  $\mathbf{b} = \langle 1, -1 \rangle$ . Find:

- $\mathbf{a} \cdot \mathbf{b} =$

- $|\mathbf{a}| =$

- $|\mathbf{b}| =$

- $\text{proj}_{\mathbf{b}} \mathbf{a} =$

- $\text{comp}_{\mathbf{a}} \mathbf{b} =$

EXAMPLE 12. Find the distance from the point  $P(-2, 3)$  to the line  $y = 3x + 5$ .

