


5.2: Maximum and Minimum Values

DEFINITION 1. Let D be the domain of a function f .

- A function f has an **absolute maximum** (or **global maximum**) at $x = c$ if $f(c) \geq f(x)$ for all x in D . In this case, we call $f(c)$ the **maximum value**.
- A function f has an **absolute minimum** (or **global minimum**) at $x = c$ if $f(c) \leq f(x)$ for all x in D . In this case, we call $f(c)$ the **minimum value**.


The maximum and minimum values of f on D are called the **extreme values** of f .

DEFINITION 2. A function f has a local maximum at $x = c$ if $f(c) \geq f(x)$ when x is near c (i.e. in a neighborhood of c). A function f has a local minimum at $x = c$ if $f(c) \leq f(x)$ when x is near c .




EXAMPLE 3. Find the absolute and local extrema of f by sketching its graph:

(a) $f(x) = x^2, -1 \leq x \leq 3$




	Local	Absolute	Value
Maximum			
Minimum			

(b) $f(x) = x^2, -3 \leq x \leq 3$




	Local	Absolute	Value
Maximum			
Minimum			

(c) $f(x) = x^2$




	<i>Local</i>	<i>Absolute</i>	<i>Value</i>
<i>Maximum</i>			
<i>Minimum</i>			

(d) $f(x) = x^3$



	<i>Local</i>	<i>Absolute</i>	<i>Value</i>
<i>Maximum</i>			
<i>Minimum</i>			


(e) $f(x) = \frac{1}{x}, 0 < x \leq 3$



	<i>Local</i>	<i>Absolute</i>	<i>Value</i>
<i>Maximum</i>			
<i>Minimum</i>			

(f) $f(x) = \begin{cases} x^4 & \text{if } -1 \leq x < 0 \\ 2 - x^4 & \text{if } 0 \leq x \leq 1 \end{cases}$

	<i>Local</i>	<i>Absolute</i>	<i>Value</i>
<i>Maximum</i>			
<i>Minimum</i>			



Extreme Value Theorem: If f is a continuous function on a closed interval $[a, b]$, then f attains both an absolute maximum and an absolute minimum.

EXAMPLE 4. Graph an example of a continuous function on a non closed interval that does not attain an absolute minimum but does attain an absolute maximum.

EXAMPLE 5. Graph an example of a function that is not continuous at a point in the given interval and yet has both absolute extrema.

DEFINITION 6. A **critical number** of $f(x)$ is a number c is in the domain of f such that either $f'(c) = 0$ or $f'(c)$ does not exist.

Illustration:

EXAMPLE 7. Find the critical numbers of $f(x)$:

(a) $f(x) = x^3 - 3x^2 + 3x$

(b) $f(x) = |4 - x^2|$

(c) $f(x) = x^{2/5}(5 - x)$

(d) $f(x) = x \ln x$

EXAMPLE 8. Find the absolute extrema for $f(x)$ on the interval I where

(a) $f(x) = x^3 - 3x^2 + 3x$, $I = [-1, 3]$

(b) $f(x) = \sqrt{3}x^2 + 2 \cos x^2, I = \left[\frac{\sqrt{\pi}}{2}, \sqrt{\pi} \right]$