

## Section 1.1: Vectors

Quantities that we measure that have magnitude but not direction are called **scalars**.

DEFINITION 1. A **vector** is a quantity that has both magnitude and direction. A 2-dimensional vector is an ordered pair  $\mathbf{a} = \langle a_1, a_2 \rangle$ . The numbers  $a_1$  and  $a_2$  are called the **components** of the vector  $\mathbf{a}$ .

Typical notation to designate a vector is a boldfaced character or a character with an arrow on it (i.e.  $\mathbf{a}$  or  $\vec{a}$ ).

DEFINITION 2. Given the points  $A(a_1, a_2)$  and  $B(b_1, b_2)$ , the vector  $\mathbf{a}$  with representation  $\vec{AB}$  is

$$\vec{AB} = \langle b_1 - a_1, b_2 - a_2 \rangle.$$

The point  $A$  here is initial point and  $B$  is terminal one.

A vector with the initial point located at the origin is called the **position** vector (or we say that a vector is in standard position).

Vectors are equal if they have the same length and direction (same slope).

EXAMPLE 3. Graph the vector with initial point  $A(1, -2)$  and terminal point  $B(2, 1)$ . Find the components of  $\vec{AB}$  and  $\vec{BA}$ .

### Vector operations

- *Scalar Multiplication:* If  $c$  is a scalar and  $\mathbf{a} = \langle a_1, a_2 \rangle$ , then

$$c\mathbf{a} = c \langle a_1, a_2 \rangle = \langle ca_1, ca_2 \rangle.$$

DEFINITION 4. Two vectors  $\mathbf{a}$  and  $\mathbf{b}$  are called **parallel** if  $\mathbf{b} = c\mathbf{a}$  with some scalar  $c$ .

If  $c > 0$  then  $a$  and  $ca$  have the same direction, if  $c < 0$  then  $a$  and  $ca$  have the opposite direction.

- *Vector addition:* If  $\mathbf{a} = \langle a_1, a_2 \rangle$  and  $\mathbf{b} = \langle b_1, b_2 \rangle$  then

$$\mathbf{a} + \mathbf{b} = \langle a_1 + b_1, a_2 + b_2 \rangle.$$

**Triangle Law**

**Parallelogram Law**

$\mathbf{a} + \mathbf{b}$  is called the **resultant vector**

EXAMPLE 5. Let  $\mathbf{a} = \langle -1, 2 \rangle$  and  $\mathbf{b} = \langle 2.1, -0.5 \rangle$ . Then  $3\mathbf{a} + 2\mathbf{b} =$

The **magnitude** or **length** of a vector  $\mathbf{a} = \langle a_1, a_2 \rangle$  is denoted by  $|\mathbf{a}|$ ,

$$|\mathbf{a}| =$$

EXAMPLE 6. Find magnitudes of the following vectors:

(a)  $\langle 3, -8 \rangle$

(b)  $\left\langle \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right\rangle$

(c)  $\mathbf{0}$

(d)  $\langle \cos \alpha, \sin \alpha \rangle$

A **unit** vector is a vector with length one. Any vector can be made into a unit vector by dividing it by its length. So, a unit vector in the direction of  $\mathbf{a}$  is

$$\mathbf{u} = \frac{\mathbf{a}}{|\mathbf{a}|}.$$

Any vector  $\mathbf{a}$  can be fully represented by providing its length,  $|\mathbf{a}|$  and a unit vector  $\mathbf{u}$  in its direction:

$$\mathbf{a} = |\mathbf{a}| \mathbf{u}$$

EXAMPLE 7. Given  $\mathbf{a} = \langle 2, -1 \rangle$ . Find

(a) a unit vector that has the same direction as  $\mathbf{a}$ ;

(b) a vector  $\mathbf{b}$  in the direction opposite to  $\mathbf{a}$  s.t  $|\mathbf{b}| = 7$ .

The **standard basis vectors** are given by the unit vectors  $\mathbf{i} = \langle 1, 0 \rangle$  and  $\mathbf{j} = \langle 0, 1 \rangle$  along the x and y directions, respectively. Using the basis vectors, one can represent any vector  $\mathbf{a} = \langle a_1, a_2 \rangle$  as


$$\mathbf{a} = a_1 \mathbf{i} + a_2 \mathbf{j}.$$

EXAMPLE 8. Given  $\mathbf{a} = 2\mathbf{i} - \mathbf{j}$ ,  $\mathbf{b} = \langle 5, -2 \rangle$ . Find a scalars  $s$  and  $t$  such that  $s\mathbf{a} + t\mathbf{b} = -4\mathbf{j}$ .

**Applications:** Quantities such as force, displacement or velocity that have direction as well as magnitude are represented by vectors.

EXAMPLE 9. *Ben walks due West on the deck of a ship at 5 mph. The ship is moving north at a speed of 25 mph. Find the direction and speed of Ben relative to the surface of the water.*

EXAMPLE 10. An 60 kg weight hangs from two wires as shown. Find the tensions (forces) in both wires and their magnitudes.



EXAMPLE 11. *An airplane, flying due east at an airspeed of 450mph, encounters a 50-mph wind acting in the direction of  $E60^\circ N$  ( $60^\circ$  North of East). The airplane holds its compass heading due East but, because of the wind, acquires a new ground speed (i.e. the magnitude of the resultant) and direction. What are they?*