

Section 1.3: Vector functions

Parametric equations:

$$x = x(t), \quad y = y(t)$$

where the variable t is called a **parameter**. Each value of the parameter t defines a point that we can plot. As t varies over its domain we get a collection of points $(x, y) = (x(t), y(t))$ on the plane which is called the **parametric curve**.


Each parametric curve can be represented as the **vector function**:

$$\vec{r}(t) = \langle x(t), y(t) \rangle.$$

Note that Parametric curves have a **direction of motion** given by increasing of parameter t . So, when sketching parametric curves we also include arrows that show the direction of motion.

EXAMPLE 1. Examine the parametric curve $x = \cos t$, $y = \sin t$, $0 \leq t \leq 3\pi/2$.

t	0	$\pi/4$	$\pi/2$...
(x, y)				




EXAMPLE 2. Given $\mathbf{r}(t) = \langle t + 1, t^2 \rangle$.

(a) Does the point $(4, 3)$ belong to the graph of $\mathbf{r}(t)$?

(b) Sketch the graph of $\mathbf{r}(t)$.

t	$\mathbf{r}(t)$
-2	
-1	
0	
1	
2	



(c) Find the Cartesian equation of $\mathbf{r}(t)$ eliminating the parameter.

EXAMPLE 3. Find the Cartesian equation for $\mathbf{r}(t) = \cos t \mathbf{i} + \cos(2t) \mathbf{j}$

EXAMPLE 4. An object is moving in the xy -plane and its position after t seconds is given by $\mathbf{r}(t) = \langle 1 + t^2, 1 + 3t \rangle$.

(a) Find the position of the object at time $t = 0$.

(b) At what time does the object reach the point $(10, 10)$.

(c) Does the object pass through the point $(20, 20)$?

(d) Find an equation in x and y whose graph is the path of the object.

A **Vector equation of the line** passing through the point (x_0, y_0) and parallel to the vector $\mathbf{v} = \langle a, b \rangle$ is given by

$$\mathbf{r} = \mathbf{r}_0 + t\mathbf{v},$$

where $\mathbf{r}_0 = \langle x_0, y_0 \rangle$.

The vector equation of the line can be rewritten in parametric form. Namely, we have

$$\begin{aligned}\langle x(t), y(t) \rangle &= \mathbf{r} = \mathbf{r}_0 + t\mathbf{v} = \\ &= \langle x_0, y_0 \rangle + t \langle a, b \rangle = \langle x_0, y_0 \rangle + \langle ta, tb \rangle = \\ &= \langle x_0 + ta, y_0 + tb \rangle.\end{aligned}$$

This immediately yields that the **parametric equations of the line** passing through the point (x_0, y_0) and parallel to the vector $\mathbf{v} = \langle a, b \rangle$ are

$$x(t) = x_0 + at, \quad y(t) = y_0 + bt.$$

EXAMPLE 5. Find parametric equations of the line

(a) passing through the point $(1, 0)$ and parallel to the vector $\mathbf{i} - 4\mathbf{j}$;

(b) passing through the point $(-4, 5)$ with slope $\sqrt{3}$;

(c) passing through the points $(7, 2)$ and $(3, 2)$.

EXAMPLE 6. Determine whether the lines $\mathbf{r}(t) = \langle 1 + t, 1 - 3t \rangle$, $\mathbf{R}(s) = \langle 1 + 3s, 12 + s \rangle$ are parallel, orthogonal or neither. If they are not parallel, find the intersection point.