

## Section 5.7: Antiderivatives

DEFINITION 1. A function  $F$  is called an **antiderivative** of  $f$  on an interval  $I$  if  $F'(x) = f(x)$  for all  $x$  in  $I$ .

EXAMPLE 2. (a) Is the function  $F(x) = x \ln(x) - x + \sin x$  is an antiderivative of  $f(x) = \ln(x) + \cos x$ ?

(b) Is the function  $F(x) = x \ln(x) - x + \sin x + 10$  is an antiderivative of  $f(x) = \ln(x) + \cos x$ ?

(c) What is the most general antiderivative of  $f(x) = \ln(x) + \cos x$ ?

THEOREM 3. If  $F$  is an antiderivative of  $f$  on an interval  $I$ , then the most general antiderivative of  $f$  on  $I$  is  $F(x) + C$ , where  $C$  is an arbitrary constant.

EXAMPLE 4. Find the most general antiderivative of  $f = 2x$ .

**Table of Antidifferentiation Formulas**

Function	Particular antiderivative	Most general antiderivative
$k$ ( $k \in \mathbb{R}$ )	$kx$	$kx + C$
$x^n$ ( $n \neq -1$ )	$\frac{x^{n+1}}{n+1}$	$\frac{x^{n+1}}{n+1} + C$
$\frac{1}{x}$	$\ln x $	
$e^x$	$e^x$	
$\cos x$	$\sin x$	
$\sin x$	$-\cos x$	
$\sec^2 x$	$\tan x$	
$\csc^2 x$	$-\cot x$	
$\sec x \tan x$	$\sec x$	
$\csc x \cot x$	$\csc x$	
$\frac{1}{\sqrt{1-x^2}}$	$\arcsin x$	
$\frac{1}{1+x^2}$	$\arctan x$	

EXAMPLE 5. Find the most general antiderivative of  $f$  where

(a)  $f(x) = 5 \sin x + 7x^6 - \sqrt[8]{x^7} + 15$

(b)  $f(x) = \frac{3x + 8 - x^2}{x^3}$

(c)  $f(x) = e^x + (1 - x^2)^{-1/2}$

EXAMPLE 6. Find  $f(x)$  given that  $f'(x) = 4 - 3(1 + x^2)^{-1}$ ,  $f(1) = 0$ .

EXAMPLE 7. Find  $f(x)$  given that  $f''(x) = 3e^x + 5\sin x$ ,  $f(0) = 1$ ,  $f'(0) = 2$ .