

5 FUNCTIONS

5.1 Definition and Basic Properties

DEFINITION 1. Let X and Y be nonempty sets. A **function** f from the set X to the set Y is a correspondence that assigns to each element x in the set X one and only one element y in the set Y , which is denoted by $f(x)$.

We call X the **domain** of f and Y the **codomain** of f .

If $x \in X$ and $y \in Y$ are such that $y = f(x)$, then y is called the **value** of f at x , or the **image** of x under f . We may also say that f **maps** x to y .

Using diagram

DEFINITION 2. Two functions f and g are **equal** if they have the same domain and the same codomain and if $f(x) = g(x)$ for all x in domain.

DEFINITION 3. The **graph** of $f : X \rightarrow Y$ is the set

$$G_f = \{(x, y) \in X \times Y \mid y = f(x)\}.$$

- We can determine a function from its domain, codomain, and graph.
- We can describe a function by formula, by listing its values, or by words.

EXAMPLE 4. Let $X = \{2, 4, 6\}$ and $Y = \{a, b, c, d\}$. Determine in which of the following cases, f is function from X to Y .

(a) $f(2) = b, f(4) = a, f(6) = d$

(b) $f(2) = c, f(4) = c, f(6) = c$

(c) $f(2) = a, f(4) = b, f(6) = c, f(4) = d$

(d) $f(2) = c, f(6) = d$

Some common functions

- **Identity** function $I_X : X \rightarrow X$ maps every element to itself:

- **Polynomial** of degree n with real coefficients a_0, a_1, \dots, a_n is a function from \mathbb{R} to \mathbb{R}

Polynomials of degrees 0,1,2,3 are constant, linear, quadratic, cubic, respectively.

EXAMPLE 5. Let $f : X \rightarrow Y$ be defined by $f(x) = x^3 + 3$. In each of the following cases find its graph and illustrate it .

(a) $X = Y = \mathbb{R}$

(b) $X = \{-1, 0, 1\}, Y = \mathbb{R}$

Range (or Image) of a Function

DEFINITION 6. Let $f : X \rightarrow Y$ be a function. The **range** of f (also called the **image** of f) is the set

$$\{y \in Y \mid y = f(x) \text{ for some } x \in X\}.$$

We denote the range (or image) of the function f by $\text{ran}f$ (or $\text{Im}f$).

EXAMPLE 7. Let $f : X \rightarrow Y$ be a function. Using symbols complete the following

- $\text{ran}f \subseteq$ _____
- $\forall y \in Y, y \in \text{ran}f \Leftrightarrow$ _____
- $y \notin \text{ran}f \Leftrightarrow$ _____

EXAMPLE 8. $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = \cos x$. Find $\text{ran}f$.

EXAMPLE 9. Let $f : [\frac{1}{3}, \infty) \rightarrow \mathbb{R}$ be defined by $f(x) = \sqrt{3x - 1}$ and $S = \{y \in \mathbb{R} \mid y \geq 0\}$. Prove that $\text{ran}f = S$.

5.2 Composition of Functions

DEFINITION 10. Let A , B , and C be nonempty sets, and let $f : A \rightarrow B$, $g : B \rightarrow C$. We define a function

$$g \circ f : A \rightarrow C,$$

called the **composition** of f and g , by

$$(g \circ f)(a) =$$

Using diagram

EXAMPLE 11. Let $A = \{1, 2, 3, 4\}$, $B = \{a, b, c, d\}$, $C = \{r, s, t, u, v\}$ and define the functions $f : A \rightarrow B$, $g : B \rightarrow C$ by their graphs:

$$G_f = \{(1, b), (2, d), (3, a), (4, a)\}, \quad G_g = \{(a, u), (b, r), (c, r), (d, s)\}.$$

Find $g \circ f$. What about $f \circ g$?

EXAMPLE 12. Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = e^x$ and $g(x) = x \sin x$. Find $f \circ g$ and $g \circ f$.

PROPOSITION 13. Let $f : A \rightarrow B$, $g : B \rightarrow C$, and $h : C \rightarrow D$. Then

$$(h \circ g) \circ f = h \circ (g \circ f),$$

i.e. composition of functions is associative.

Proof.

Section 5.3 Surjective (or onto) and Injective (or one-to-one) Functions

Surjective functions (“onto”)

DEFINITION 14. Let $f : X \rightarrow Y$ be a function. Then f is **surjective** (or a surjection) if the range of f coincides with its codomain, i.e.

$$\text{ran } f = Y.$$

Note: surjection is also called “onto”.

Proving surjection:

We know that for all $f : X \rightarrow Y$: _____

Thus, to show that $f : X \rightarrow Y$ is a surjection it is sufficient to prove that _____

In other words,

to prove that $f : X \rightarrow Y$ is a surjective function it is sufficient to show that _____

Question: How to disprove surjectivity?

EXAMPLE 15. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow [0, \infty)$ defined by $f(x) = g(x) = x^4$. Determine whether the following are true

(a) $\text{ran} f = \text{ran}(g)$

(b) $f = g$

(c) f is surjective

(d) g is surjective

EXAMPLE 16. Prove that the function $f : \mathbb{R} - \{2\} \rightarrow \mathbb{R} - \{3\}$ defined by $f(x) = \frac{3x}{x-2}$ is surjective.

Injective functions (“one to one”)

DEFINITION 17. Let $f : X \rightarrow Y$ be a function. Then f is **injective** (or an **injection**) if whenever $x_1, x_2 \in X$ and $x_1 \neq x_2$, we have $f(x_1) \neq f(x_2)$.

In other words, f is injective if and only if the ranges of every two distinct points in the domain of f are distinct.

EXAMPLE 18. Given $X = \{1, 2, 3\}$ and $Y = \{3, 4, 5\}$. Determine whether the following functions are injective. Justify your answer.

(a) $f : X \rightarrow Y$ defined by $G_f = \{(1, 3), (2, 4), (3, 5)\}$

(b) $g : X \rightarrow Y$ defined by $G_g = \{(1, 5), (2, 4), (3, 4)\}$

Proving injection:

Let $P(x_1, x_2) : x_1 \neq x_2$ and $Q(x_1, x_2) : f(x_1) \neq f(x_2)$.

Then by definition f is injective if _____.

Using contrapositive, we have _____.

In other words, to prove injection show that:

Question: How to disprove injectivity?

EXAMPLE 19. Prove or disprove injectivity of the following functions.

(a) $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \sqrt[5]{x}$.

(b) $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = x^4$.

$$(c) f : \mathbb{Z} \rightarrow \mathbb{Z}, f(n) = \begin{cases} n/2 & \text{if } n \in \mathbb{E}, \\ 2n & \text{if } n \in \mathbb{O}. \end{cases}$$

$$(d) f : \mathbb{Z} \rightarrow \mathbb{Z}, f(n) = \begin{cases} n & \text{if } n \in \mathbb{E}, \\ 5n & \text{if } n \in \mathbb{O}. \end{cases}$$

Discussion Exercise.

- Must a strictly increasing or decreasing function be injective?

- Must an injective function be strictly increasing or decreasing?

EXAMPLE 20. Prove or disprove injectivity of the following functions. In each case, $f : \mathbb{R} \rightarrow \mathbb{R}$.

(a) $f(x) = 3x^5 + 5x^3 + 2x + \pi$.

(b) $f(x) = x^{12} + x^8 - x^4 + 12$.

Bijjective functions

DEFINITION 21. A function that is both surjective and injective is called **bijjective** (or bijection.)

Complete the following.

• f is not bijjective \Leftrightarrow _____

• f is surjective $\Leftrightarrow (\text{codom } f \subseteq \text{_____}) \Leftrightarrow (\forall y, y \in \text{codom } f \Rightarrow \text{_____}) \Leftrightarrow$

$\Leftrightarrow (\forall y, y \in \text{codom } f \Rightarrow \exists x \in \text{dom } f \text{_____}) \Leftrightarrow (\forall y \in \text{codom } f, \exists x \in \text{dom } f \text{_____})$

In other words, f is surjective if and only if every point in $\text{codom } f$ has a preimage in the $\text{dom } f$.

If in addition f is injective, then we obtain

• f is bijjective $\Leftrightarrow (\exists! x \in \text{dom } f \text{_____})$

In other words, f is bijjective if and only if every point in $\text{codom } f$ has a unique preimage in the $\text{dom } f$.

EXAMPLE 22. Determine which of the following functions are bijjective.

(a) $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^3$.

(b) $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$.

PROPOSITION 23. *Let $f : A \rightarrow B$ and $g : B \rightarrow C$. Then*

i. *If f and g are surjections, then $g \circ f$ is also a surjection.*

Proof.

ii. *If f and g are injections, then $g \circ f$ is also an injection.*

Proof.

COROLLARY 24. *If f and g are bijections, then $g \circ f$ is also a bijection.*

PROPOSITION 25. Let $f : X \rightarrow Y$. Then $f \circ I_X = f$ and $I_Y \circ f = f$.

5.4 Invertible Functions

Inverse Functions

DEFINITION 26. Let $f : X \rightarrow Y$ be a function. We say that f is **invertible** if there is a function $g : Y \rightarrow X$ such that for all $x \in X$ and for all $y \in Y$,

$$y = f(x) \quad \Leftrightarrow \quad x = g(y).$$

We say that such a function g is an **inverse function** of f .

Question: What is the inverse of g ?

REMARK 27. f is invertible if and only if its inverse is invertible.

EXAMPLE 28. Show that the function $f : \mathbb{R} - \{2\} \rightarrow \mathbb{R} - \{3\}$ defined by $f(x) = \frac{3x}{x-2}$ is invertible and find its inverse function. (Note that the given function is bijective.)

PROPOSITION 29. *A function $f : X \rightarrow Y$ is invertible if and only if there exists a function $g : Y \rightarrow X$ such that*

$$g \circ f = I_X \quad \text{and} \quad f \circ g = I_Y.$$

PROPOSITION 30. *The inverse function is unique.*

Proof.

Notation

When $f : X \rightarrow Y$ is invertible, the unique inverse function is denoted by f^{-1} , and $f^{-1} : Y \rightarrow X$.

REMARK 31. Finding the inverse of a bijective function is not always possible by algebraic manipulations. For example,

if $f(x) = e^x$ then $f^{-1}(x) = \underline{\hspace{2cm}}$

The function $f(x) = 3x^5 + 5x^3 + 2x + 220$ is known to be bijective, but there is no way to find expression for its inverse.

THEOREM 32. *A function $f : A \rightarrow B$ is invertible if and only if f is bijective.*

COROLLARY 33. *If a function $f : A \rightarrow B$ is bijective, so is f^{-1} .*