

## 14.6: Parametric surfaces and their areas


Consider a continuous vector valued function of two variables

$$\mathbf{r}(u, v) = x(u, v)\mathbf{i} + y(u, v)\mathbf{j} + z(u, v)\mathbf{k}, \quad (u, v) \in D.$$

**Parametric surface:**

$$S : x = x(u, v), \quad y = y(u, v), \quad z = z(u, v), \quad (u, v) \in D.$$

In other words, the surface  $S$  is traces out by the position vector  $\mathbf{r}(u, v)$  as  $(u, v)$  moves throughout the region  $D$ .



**EXAMPLE 1.** Determine the surface given by the parametric representation

$$\mathbf{r}(u, v) = \langle u, u \cos v, u \sin v \rangle, \quad 1 \leq u \leq 5, \quad 0 \leq v \leq 2\pi$$

EXAMPLE 2. Give parametric or vector representations for each of the following surfaces:

(a) the cylinder:  $x^2 + y^2 = 9$ ,  $1 \leq z \leq 5$ .

(b) the upper half-sphere:  $z = \sqrt{100 - x^2 - y^2}$ .

(c) elliptic paraboloid:  $x = 3y^2 + z^2 + 1$ .

(d) *elliptic paraboloid*  $y = x^2 + 4z^2$

*CONCLUSION:* To parametrize surface we may use polar, cylindrical or spherical coordinates, or

- $z = f(x, y) \longrightarrow \mathbf{r}(x, y) = x\mathbf{i} + y\mathbf{j} + f(x, y)\mathbf{k}$
- $y = f(x, z) \longrightarrow \mathbf{r}(x, z) = x\mathbf{i} + f(x, z)\mathbf{j} + z\mathbf{k}$
- $x = f(y, z) \longrightarrow \mathbf{r}(y, z) = f(y, z)\mathbf{i} + y\mathbf{j} + z\mathbf{k}$
- **Tangent planes:**

*PROBLEM:* Find a normal vector to the tangent plane to a parametric surface  $S$  given by a vector function  $\mathbf{r}(u, v)$  at a point  $P_0$  with position vector  $\mathbf{r}(u_0, v_0)$ , i.e.  $P_0(x(u_0, v_0), y(u_0, v_0), z(u_0, v_0))$

*Solution:*

### The normal vector

$$\mathbf{N} = \mathbf{N}(u, v) =$$

If a normal vector is not  $\mathbf{0}$  then the surface  $S$  is called **smooth** (it has no "corner").

**EXAMPLE 3.** Find the tangent plane to the surface with parametric equations  $x = uv + 1, y = ue^v, z = ve^u$  at the point  $(1, 0, 0)$ .

*Special Case:* a surface  $S$  given by a *graph*  $z = f(x, y)$ . Then one can choose the following parametrization of  $S$ :

$$\mathbf{r}(x, y) =$$

and then the normal vector is

$$\mathbf{N} =$$

• **Surface Area:**

Consider a smooth surface  $S$  given by

$$\mathbf{r}(u, v) = x(u, v)\mathbf{i} + y(u, v)\mathbf{j} + z(u, v)\mathbf{k}, \quad (u, v) \in D,$$

then

$$dS = |N(u, v)| du dv =$$

and the **surface area**

$$A(S) = \iint_S dS = \iint_D |\mathbf{r}_u \times \mathbf{r}_v| dA.$$

REMARK 4. *Special Case:* a surface  $S$  given by a *graph*  $z = f(x, y)$  we have

$$\mathbf{r}(x, y) = x\mathbf{i} + y\mathbf{j} + f(x, y)\mathbf{k}$$

and

$$dS = |\mathbf{N}(x, y)| dA =$$

EXAMPLE 5. *Find the surface area of the surface*

$$S : \quad x = uv, \quad y = u + v, \quad z = u - v, \quad u^2 + v^2 \leq 1.$$

EXAMPLE 6. *Find the surface area of the part paraboloid  $z = x^2 + y^2$  between two planes:  $z = 0$  and  $z = 4$ .*