

14.9: The Divergence Theorem

Let E be a simple solid region with the boundary surface S (which is a closed surface.) Let S be positively oriented (i.e.the orientation on S is outward that is, the unit normal vector $\hat{\mathbf{n}}$ is directed outward from E).

The Divergence Theorem: Let E be a simple solid region whose boundary surface S has positive (outward) orientation. Let \mathbf{F} be a continuous vector field on an open region that contains E . Then

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iiint_E \operatorname{div}\mathbf{F} \, dV.$$

EXAMPLE 1. Let $E = \{(x, y, z) : x^2 + y^2 \leq R^2, 0 \leq z \leq H\}$. Find the flux of the vector field $\mathbf{F} = \langle 1 + x, 3 + y, z - 10 \rangle$ over ∂E .

REMARK 2. If $\mathbf{F} = \left\langle \frac{x}{3}, \frac{y}{3}, \frac{z}{3} \right\rangle$ then

EXAMPLE 3. Let E be the solid bounded by the paraboloid $z = 4 - x^2 - y^2$ and the xy -plane.

Evaluate $I = \iint_S \langle x^3, 2xz^2, 3y^2z \rangle \cdot d\mathbf{S}$ if

(a) S is the boundary of the solid E .

(B) S is the part of the paraboloid $z = 4 - x^2 - y^2$ between the planes $z = 0$ and $z = 4$.

EXAMPLE 4. Evaluate $I = \iint_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S}$ if S is the boundary of

(a) ellipsoid $E = \left\{ \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} \leq 1 \right\}$ and $\mathbf{F} =$

(b) an arbitrary simple solid region E and F is an arbitrary continuous vector field.