


## 12.7: Maximum and minimum values

Function $y = f(x)$	Function of two variables $z = f(x, y)$
<p>DEFINITION 1. A function <math>f(x)</math> has a local maximum at <math>x = a</math> if <math>f(a) \geq f(x)</math> when <math>x</math> is near <math>a</math> (i.e. in a neighborhood of <math>a</math>). A function <math>f</math> has a local minimum at <math>x = a</math> if <math>f(a) \leq f(x)</math> when <math>x</math> is near <math>a</math>.</p> <p>.</p> <p>If the inequalities in this definition hold for ALL points <math>x</math> in the domain of <math>f</math>, then <math>f</math> has an <b>absolute max</b> (or <b>absolute min</b>) at <math>a</math></p> <p>If the graph of <math>f</math> has a tangent line at a local extremum, then the tangent line is horizontal: <math>f'(a) = 0</math>.</p> 	<p>DEFINITION 2. A function <math>f(x, y)</math> has a local maximum at <math>(x, y) = (a, b)</math> if <math>f(a, b) \geq f(x, y)</math> when <math>(x, y)</math> is near <math>(a, b)</math> (i.e. in a neighborhood of <math>(a, b)</math>). A function <math>f</math> has a local minimum at <math>(x, y) = (a, b)</math> if <math>f(a, b) \leq f(x, y)</math> when <math>(x, y)</math> is near <math>(a, b)</math>.</p> <p>If the inequalities in this definition hold for ALL points <math>(x, y)</math> in the domain of <math>f</math>, then <math>f</math> has an <b>absolute maximum</b> (or <b>absolute minimum</b>) at <math>(a, b)</math>.</p> <p>If the graph of <math>f</math> has a tangent plane at a local extremum, then the tangent PLANE is horizontal.</p>

**THEOREM 3.** If  $f$  has a local extremum (that is, a local maximum or minimum) at  $(a, b)$  and the first-order partial derivatives exist there, then

$$f_x(a, b) = f_y(a, b) = 0 \quad (\text{or, equivalently, } \nabla f(a, b) = 0.)$$

**DEFINITION 4.** A point  $(a, b)$  such that  $f_x(a, b) = 0$  and  $f_y(a, b) = 0$ , or one of this partial derivatives does not exist, is called a **critical point** of  $f$ .

At a critical point, a function could have a local max or a local min, or neither.

We will be concerned with two important questions:

- Are there any local or absolute extrema?
- If so, where are they located?

SETS in  $\mathbb{R}^2$ 

in $\mathbb{R}$	in $\mathbb{R}^2$
close interval $[a, b]$	close set
open interval $(a, b)$	open set
end points of an interval	boundary points

DEFINITION 5. A **bounded set** in  $\mathbb{R}^2$  is one that contained in some disk.

*THE EXTREME VALUE THEOREM:*

Function $y = f(x)$	Function of two variables $z = f(x, y)$
If $f$ is continuous on a closed interval $[a, b]$ , then $f$ attains an absolute maximum value $f(x_1)$ and an absolute minimum value $f(x_2)$ at some points $x_1$ and $x_2$ in $[a, b]$ .	If $f$ is continuous on a closed bounded set $D$ in $\mathbb{R}^2$ , then $f$ attains an absolute maximum value $f(x_1, y_1)$ and an absolute minimum value $f(x_2, y_2)$ at some points $(x_1, y_1)$ and $(x_2, y_2)$ in $D$ .

EXAMPLE 6. Find extreme values of  $f(x, y) = x^2 + y^2$ .

	Local	Absolute
Maximum		
Minimum		

Domain:

EXAMPLE 7. Find extreme values of  $f(x, y) = \sqrt{1 - x^2 - y^2}$ .

	Local	Absolute
Maximum		
Minimum		

Domain:

EXAMPLE 8. Find extreme values of  $f(x, y) = y^2 - x^2$ .

	Local	Absolute
Maximum		
Minimum		

Domain:

REMARK 9. Example 8 illustrates so called **saddle point** of  $f$ . Note that the graph of  $f$  crosses its tangent plane at  $(a, b)$ .

EXAMPLE 10. Find the points on the surface  $z^2 = xy + 1$  that are closest to the origin.

### ABSOLUTE MAXIMUM AND MINIMUM VALUES on a closed bounded set.

#### THE EXTREME VALUE THEOREM:

To find the absolute maximum and minimum values of a continuous function  $f$  on a closed interval  $[a, b]$ :

1. Find the values of  $f$  at the critical points of  $f$  in  $(a, b)$ .
2. Find the extreme values of  $f$  at the endpoints of the interval.
3. The largest of the values from steps 1&2 is the absolute max value; the smallest of the values from steps 1&2 is the absolute min value.

To find the absolute max and min values of a continuous function  $f$  on a closed bounded set  $D$ :

1. Find the values of  $f$  at the critical points of  $f$  in  $D$ .
2. Find the extreme values of  $f$  on the boundary of  $D$ . (This usually involves the Calculus I approach for this work.)
3. The largest of the values from steps 1&2 is the absolute maximum value; the smallest of the values from steps 1&2 is the absolute minimum value.

- The quantity to be maximized/minimized is expressed in terms of variables (as few as possible!)
- Any constraints that are presented in the problem are used to reduce the number of variables to the point they are independent,
- After computing partial derivatives and setting them equal to zero you get purely algebraic problem (but it may be hard.)
- Sort out extreme values to answer the original question.

EXAMPLE 11. A lamina occupies the region  $D = \{(x, y) : 0 \leq x \leq 3, -2 \leq y \leq 4 - 2x\}$ . The temperature at each point of the lamina is given by

$$T(x, y) = 4(x^2 + xy + 2y^2 - 3x + 2y) + 10.$$

Find the hottest and coldest points of the lamina.

### Second derivatives test:

Suppose  $f''$  is continuous near  $a$  and  $f'(c) = 0$  (i.e.  $a$  is a critical point).

- If  $f''(c) > 0$  then  $f(c)$  is a local minimum.
- If  $f''(c) < 0$  then  $f(c)$  is a local maximum.

**NOTE:**

- If  $f''(c) = 0$ , then the test gives no information.

Suppose that the second partial derivatives of  $f$  are continuous near  $(a, b)$  and  $\nabla f(a, b) = \mathbf{0}$  (i.e.  $(a, b)$  is a critical point).

Let  $\mathcal{D} = \mathcal{D}(a, b) = f_{xx}(a, b)f_{yy}(a, b) - [f_{xy}(a, b)]^2$

- If  $\mathcal{D} > 0$  and  $f_{xx}(a, b) > 0$  then  $f(a, b)$  is a local minimum.
- If  $\mathcal{D} > 0$  and  $f_{xx}(a, b) < 0$  then  $f(a, b)$  is a local maximum.
- If  $\mathcal{D} < 0$  then  $f(a, b)$  is not a local extremum (saddle point).

If  $\mathcal{D} = 0$  or does not exist, then the test gives no information. fails.

To remember formula for  $\mathcal{D}$ :

$$\mathcal{D} = f_{xx}f_{yy} - [f_{xy}]^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}$$

**EXAMPLE 12.** Use the Second Derivative Test to confirm that a local cold point of the lamina in the previous Example is  $(2, -1)$ .

EXAMPLE 13. Find the local extrema of  $f(x, y) = x^3 + y^3 - 3xy$ .

*Solution:* Find critical points:

Calculate the second partial derivatives and  $\mathcal{D}$ .

$f_{xx} =$		
$f_{xy} =$		
$f_{yy} =$		
$\mathcal{D}$		

EXAMPLE 14. *The mountain is defined by  $z = xy$  in the elliptical domain*

$$D = \left\{ (x, y) \mid \frac{x^2}{16} + y^2 \leq 1 \right\}.$$

(a) *Find the top of the mountain.*

(b) *Is the critical point found in the previous item the highest or the lowest in its neighborhood?*